2-P-139

Poster Sessions

A dendritic spine shrinkage in response to extracellular high K^+ is inhibited by Co^{2+} or Cd^{2+}

Naoko Konishi, Toshinori Sawano, Jin Nakatani, Hidekazu Tanaka

Lab. Pharm., Grad.sch. Life sci., Ritsumeikan Univ.

In the brain, neurons communicate with each other at excitatory and inhibitory synapses. In each excitatory synapse, a dendritic spine receives the excitatory neurotransmitter released by an axon terminal. An extracellular high K^+ induces robust depolarization and mimics massive synaptic transmission. A dendritic spine transiently shrinks in high K^+ solution. In this study, we investigated the mechanisms of the spine shrinkage using cultured hippocampal neurons. The spine shrinkage observed in response to high K^+ was inhibited in the presence of extracellular Co^{2+} or Cd^{2+} . Then we investigated Ca^{2+} -signaling pathway because these metal ions are supposed to be wide spectrum Ca^{2+} -channel blockers. We preincubated neurons with ROCK inhibitor Y27632 or CaMKII inhibitor KN-93. However, neither of them inhibited the spine shrinkage. Finally, we tested whether myosin is involved in the spine shrinkage. A myosin ATPase inhibitor, blebbistatin, did not inhibit the spine shrinkage in high K^+ . Taken together, extracellular high K^+ -triggered Ca^{2+} -influx seem to induce the spine shrinkage; however, the downstream signaling pathway or the motor machinery has not yet been determined.