1-P2-25

Poster Sessions

A Discussion on the Mechanism of Actions of a gold compound, Auranofin

Masamichi Yamashita

Dept. Biosci. Daily Life, Coll. Biores. Sci., Nihon Univ.

Effect of an antirheumatic gold compound, Auranofin (AF), have been revealed. 1-10 μ M of AF inhibited prostaglandin E₂ (PGE₂) production and nitric oxide (NO) production in macrophages treated with some stimulatory compounds. We found that AF reduced cyclooxygenase (COX)-2 and inducible NO synthase (iNOS) mRNA levels, but not affected their enzyme activities. We then revealed that AF partly inhibited the nuclear-translocation of NF- κ B while PGE₂ production was almost completely inhibited.

In the review on manuscripts, AF is reported to down-regulate the mRNA levels of *SLC47A1*, *SLC22A1*, *ATP1A3*, *ABCG2*, *SLCO1B1*, *PIK3CA*, and up-regulate the mRNA levels of *ABCC3*, *ABCB1*, *SLC22A3*, *ABCC2*, *SLC31A2*, *CLTD*, *CLTB*. PGs are known to be transported by 4 transporters, OATP-PG (SLC22A22), SLC21A2 (OATP2A1, SLCO2), SLC21A11 (OATP3A1), and ABCC4, though I could not identify whether AF reduced PGE₂ concentration in cellular cultured medium as described above via these transporters by downregulation for export of PGE₂.