3-S30-4

Identification of pathological factors for Alzheimer disease using CRISPR/Cas9-based screening system

Taisuke Tomita

Dept Neuropathol Neurosci, Grad Sch Pharm Sci, Univ Tokyo

Alzheimer disease (AD) is pathologically characterized by the deposition of the amyloid- β peptide ($A\beta$) as senile plaques in the brain. $A\beta$ is generated through the sequential cleavage of amyloid precursor protein (APP) by β -site APP cleaving enzyme 1 (BACE1) and γ -secretase. Whereas the aberrant metabolism of $A\beta$ has been implicated in the etiology of AD, the precise regulatory mechanism of $A\beta$ generation remains unclear. In this study, we established a genetic screening based on the CRISPR/Cas9 system to identify novel regulators of $A\beta$ production. And we successfully identified calcium and integrin-binding protein 1 (CIB1) as a negative regulator of $A\beta$ production. CIB1 is a ubiquitously expressed, myristoylated, calcium-binding adaptor protein. We found that CIB1 interacts with the γ -secretase complex. Moreover, the disruption of CIB1 specifically reduced the cell-surface localization of the γ -secretase complex, without changing the intrinsic activity of γ -secretase. Finally, we confirmed using the single-cell RNA-seq data in human that CIB1 mRNA level in neuron was decreased in the early stage of AD. Taken together, our results indicate that CIB1 regulates $A\beta$ production via controlling the subcellular localization of γ -secretase, suggesting CIB1 is involved in the development of AD.