Stability of secondary metabolite-producing aquatic endophytic fungi from Lotus against Methicillin-resistant *Staphylococcus aureus*

Surachai Techaoei¹, Umpava Pinruan², Salilaporn Nuankaew²

¹Thai Traditional Medicine College, Rajamangala University of Technology Thanyaburi, Thailand, ²National Center for Genetic Engineering and Biotechnology (BIOTEC), Khlong Luang Pathum Thani, Thailand

The research was to conduct the stability potential of secondary metabolite of five aquatic endophytic fungi isolated from Lotus in Thailand, *Alternaria longipes*, *Pleosporales* sp., *Setosphaeria monoceras*, *Curvularia lunata* and *Alternaria eichhorniae*. All fungal were cultured in PDA and extract with organic solvent as hexane, ethyl acetate and methanol before tested by using disc diffusion method. The thermal and UV radiation were tested stability for antimicrobial potential. Result showed that the crude extract of *Alternaria longipes* and *Alternaria eichhorniae* (ethyl acetate extracts) showed thermal stability at 55-121° C and UV stability with tested microorganism as Methicillin-resistant *Staphylococcus aureus* (MRSA) at inhibition zone between of 10.00 ± 0.00 mm and 10.33 ± 0.57 mm, respectively. The further study will be focused on chemical characterization and function of antifungal activity to apply as natural drugs in the future.