Upregulation of Ca^{2+} -sensing receptors by PDGF signal in idiopathic pulmonary arterial hypertension

Aya Yamamura, Rie Takahashi, Hisaki Hayashi, Motohiko Sato

Department of Physiology, Aichi Medical University, Japan

Pulmonary arterial hypertension (PAH) is a rare, progressive, and fatal disease of unknown pathogenesis. Sustained vasoconstriction and vascular remodeling are key pathogenic events that lead to early morbidity and mortality. These events have been linked to cytosolic Ca^{2+} handling in pulmonary arterial smooth muscle cells (PASMCs). An increase in cytosolic Ca^{2+} concentration in PASMCs is a major trigger for pulmonary vasoconstriction and an important stimulus for PASMC proliferation. We have previously shown that Ca^{2+} -sensing receptor (CaSR) is upregulated in PASMCs from patients with idiopathic pulmonary arterial hypertension (IPAH) and animal models of pulmonary hypertension. The enhanced CaSR function contributes to augmented Ca^{2+} responses and cell proliferation in IPAH-PASMCs. In this study, we examined to elucidate the upregulation mechanism of CaSR in IPAH-PASMCs. In normal-PASMCs, expression of CaSR was increased by the exposure to 20 ng/ml platelet-derived growth factor (PDGF), which is known as an enhanced endogenous signal associated with IPAH, for 48 h. Expression of PDGF α receptors in IPAH-PASMCs was identical to that in normal-PASMCs, while expression of PDGF β receptors in IPAH-PASMCs was higher than that in normal-PASMCs. In addition, phosphorylation of PDGF β receptors by PDGF stimulation in IPAH-PASMCs was long-lasting compared to in normal-PASMCs. Phosphorylation levels of ERK, p38, and Akt, which are downstream pathways of PDGF receptors, were also higher in IPAH-PASMCs than normal-PASMCs. In normal-PASMCs, PDGF stimulation facilitated both proliferation and migration of PASMCs. Furthermore, siRNA knockdown of PDGF β receptors (50 nM) attenuated the CaSR upregulation in IPAH-PASMCs. Pretreatment with 5 μM imatinib, a blocker of PDGF receptors, or 10 μM NPS2143, an antagonist of CaSR, inhibited the PDGF-induced CaSR upregulation in IPAH-PASMCs. These results suggest that PDGF signal activates the upregulation mechanism of CaSR in IPAH-PASMCs. In conclusion, the crosslink between CaSR and PDGF signals is a novel pathogenic mechanism contributing to the enhanced Ca^{2+} influx and excessive cell proliferation in PAH.