Comparison of the effect of lipopolysaccharide on tumor necrosis factor α (TNF- α) secretion and *TNF* and *TNFR1* mRNA levels in feline endometrium throughout the estrous cycle during pyometra and after medroxyprogesterone acetate treatment

Ewelina JURZA-PIOTROWSKA¹⁾ and Marta J. SIEMIENIUCH¹⁾

¹⁾Department of Reproductive Immunology and Pathology, Institute of Animal Reproduction and Food Research, Polish Academy of Sciences, 10-748 Olsztyn, Poland

Abstract. Endotoxins released by Gram-negative bacteria are potent stimulators of tumor necrosis factor α (TNF- α) production. The objectives of this study were to evaluate plasma levels of TNF- α , TNF- α secretion, and mRNA levels of *TNF* and TNF- α receptor type 1 (*TNFR1*) following exposure to lipopolysaccharide (LPS). For this, we used cultured endometrial cells or organ cultures, throughout the estrous cycle, after hormone treatment with medroxyprogesterone acetate (MPA), and during pyometra. Plasma TNF- α concentrations were increased in animals at estrus ($P < 0.05$) compared to other groups. In the LPS-challenged endometrium, secretion of TNF- α by tissues collected during estrus increased ($P < 0.001$) compared to that of other groups. LPS, alone or combined with TNF- α , upregulated *TNF* gene expression in the feline endometrium at diestrus ($P < 0.001$ for both treatments), in queens treated short-term with MPA ($P < 0.01$ and $P < 0.05$, respectively) and in queens treated long-term with MPA ($P < 0.01$ and $P < 0.001$, respectively). During pyometra, *TNF* and *TNFR1* mRNA were increased only after tissues were challenged with TNF- α and LPS ($P < 0.001$ and $P < 0.01$, respectively). When cultured endometrial cells were challenged with LPS, the concentration of TNF- α increased only in epithelial cells after 4 h and 12 h ($P < 0.05$ and $P < 0.01$, respectively). Since LPS did not affect stromal cells, but TNF- α increased its own transcript after 2 h ($P < 0.01$), 4 h ($P < 0.05$) and 12 h ($P < 0.001$), we assume that stromal cells are not directly involved in pathogen recognition, as was the case for epithelial cells.

Key words: Feline endometrium, Lipopolysaccharide (LPS), Medroxyprogesterone acetate (MPA) treatment, Pyometra, Tumor necrosis factor α (TNF- α)

(J. Reprod. Dev. 62: 385–391, 2016)

The endometrium, consisting of epithelial, stromal, endothelial, smooth muscle, and immune cells, is the first line of defense against invading pathogens. A single layer of columnar epithelium is in constant contact with the uterine lumen, which may be intermittently contaminated by opportunistic bacteria or pathogens. Epithelial cells are able to recognize pathogen-associated molecular patterns (PAMPs) via pattern recognition receptors (PRRs), which include the Toll-Like Receptor family (TLR) [1]. Among nine types of feline TLRs (TLR 1-9) [2, 3], TLR2/4 are the best characterized, particularly with regard to their role in pathogen recognition. We previously confirmed the presence of both receptor types at the gene and protein level in feline endometrial epithelial and stromal cells [4]. In dogs, engagement of TLRs initiates a signaling cascade, resulting in altered expression of genes such as *prostaglandin-endoperoxide synthase 2* (*PTGS2*), *PGE₂-synthase* (*PGES*), *interleukin (IL)-1 β* , and *tumor necrosis factor alpha* (*TNF*) [5]. In cats, lipopolysaccharide (LPS), which is

the main ligand for TLR4, is responsible for upregulation of *PTGS2*, *PGES*, and *prostaglandin F₂ α synthase* (*PGFS*) mRNA in the cultured endometrial cells, and also for increasing prostaglandin secretion [6]. In an earlier study, we showed that treatment of cultured endometrial cells with LPS results in abundant TNF- α secretion; however, this effect was specific to epithelial cells (and not stromal cells) [7]. TNF- α is a multifunctional proinflammatory cytokine that plays a role in the activation of immune cells such as macrophages, granulocytes, and cytotoxic T cells, and in the maturation of dendritic cells [8]. Besides its immunomodulatory function, TNF- α is involved in the control of cell differentiation, tissue renewal, and restructuring [9]. Although TNF- α is predominantly secreted by endotoxin-stimulated monocytes/macrophages [10], it can be synthesized by many different tissues and cell types, including feline endometrium, as recently shown [7]. Generally, concentrations of TNF- α correlate with the severity of illness and outcome [11]. Moreover, 10-fold higher plasma concentrations of TNF- α were observed in dogs with pyometra compared to healthy controls [12]. Pyometra, with the clinical manifestation of pus accumulation in the uterine lumen, is a common reproductive disorder in bitches but is observed less frequently in queens [13, 14]. Hormonal imbalances together with bacterial infections contribute to the development of pyometra. The development of this condition is observed during diestrus or the luteal-like phase and can be a result of treatment with megestrol or medroxyprogesterone acetate (used

Received: January 7, 2016

Accepted: April 8, 2016

Published online in J-STAGE: April 25, 2016

©2016 by the Society for Reproduction and Development

Correspondence: MJ Siemieniuch (e-mail: m.siemieniuch@pan.olsztyn.pl)

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/4.0/>>.

for silencing estrus behavior in dogs and cats). High concentrations of endogenous or exogenous progesterone increase adherence of bacteria to feline and canine endometrium and reduce resistance to local infection. Moreover, high concentrations of progesterone lead to proliferation and cystic dilation of the endometrial glands and excessive mucus secretion [13]. Mucus accumulation in the uterine lumen creates favorable conditions for the growth of bacteria, specifically, Gram-negative *Escherichia coli* (*E. coli*) [15], which is also present in the reproductive tract of healthy queens. Recently we showed that LPS could stimulate epithelial cells to produce TNF- α , without affecting stroma, in the feline endometrium [7]. This study also showed that TNF- α and both of its receptors localized only to superficial and glandular epithelium, with essentially no localization to the stroma; however, changes in gene expression of *TNF* and *TNF receptor type 1* (*TNFR1*) were not shown [7]. However, it is still unclear if the estrous cycle, treatment with MPA, or the presence of uterine inflammation affect the endometrial concentration of TNF- α during the inflammatory response to LPS. Therefore, the aims of the present study were as follows: (1) evaluate TNF- α secretion by feline endometrial organ cultures throughout the estrous cycle after exposure to LPS, after hormone treatment with MPA, and during pyometra; (2) identify the immunological profile of queens throughout the estrous cycle, after treatment with MPA and during pyometra by studying plasma TNF- α ; (3) analyze mRNA levels of *TNF/TNFR1* after treatment with LPS, TNF- α , or both; (4) evaluate *TNF/TNFR1* mRNA levels and TNF- α production in stimulated epithelial and stromal cells *in vitro*.

Materials and Method

Animals

All procedures were approved by the Local Animal Care and Use Committee in Olsztyn, Poland (No. 60/2010/DTN). Adult domestic queens (total $n = 45$, ranging in age from 7 months to 11 years, with an average age of 2.67 years) were used in this study. Based on circulating concentrations of progesterone in peripheral blood and macroscopic observations of ovaries [16], 30 animals were assigned to appropriate estrous cycle stage groups (anestrus, estrus, mid and late diestrus). Animals with a smooth ovarian surface and a basal progesterone (P4) level > 1 ng/ml were classified as being in anestrus phase. Uterine samples from queens in the anestrus phase were collected after breeding season. Estrus phase (day 1–7) was characterized by the presence of ovarian follicles ≥ 2 mm in diameter. Queens in estrus phase were housed individually or in pairs, with or without contact with an intact male. No pharmacological treatment was performed to provoke ovulation in the animals. Queens with the presence of a reddish corpus luteum (CL) on the ovaries and a mean P4 concentration above 20 ng/ml were classified as being in mid diestrus phase (day 15–20), whereas animals with a pale CL and a level of progesterone < 5 ng/ml were assigned to late diestrus (day 30–35). After interviewing owners, 10 additional cats were assigned equally to the following groups: (i) acute MPA-treated group, in which animals had previously been orally treated with MPA (Promon Vet®, Pfizer Animal Health, Louvain-la-Neuve, France; 5 mg/animal/week) for one month to prevent estrus behavior, or (ii) long-term MPA treated group (5 mg/animal/week), in which animals

were treated for four to twelve months. The mean serum level of P4 in the MPA-treated group was > 10 ng/ml. No pharmacological treatment was performed to provoke ovulation in the animals. The last five queens were classified in the pyometra group. These animals were not treated with MPA and had a developing corpora lutea (CL), damage to the uterine tissue, and a fluid-filled uterus. The mean plasma concentration of P4 in the pyometra group ranged from 8 to 14 ng/ml. All the animals were ovariohysterectomized at the owners' request and consent, and cats were anesthetized with ketamine (20 mg/kg) and acepromazine (0.5 mg/kg).

Tissue and blood collection

From all queens, 1–1.5 ml of blood was collected into EDTA-containing tubes for TNF- α analyses, immediately before surgery. Plasma was separated by centrifugation of whole blood at $1500 \times g$ for 10 min, and stored at -20°C . Immediately after ovariohysterectomy (OVH), uteri were rinsed with sterile saline to remove blood contamination, placed into fresh saline at 4°C , and transported on ice to the laboratory. Both horns of each uterus ($n = 35$) were slit longitudinally and the endometrium was separated from the myometrium using a binocular microscope (Olympus SZX7, Tokyo, Japan).

Ex vivo organ cultures of endometrium

To evaluate the effect of LPS and TNF- α on the endometrium and *TNF* and *TNFR1* gene expression, organ cultures of endometrium tissue were generated from each uterus (sixteen explants from each uterus, $n = 35$). Endometrial fragments were prepared as previously described [4, 6]. Using doses established in that study, cultured endometrial tissues were stimulated with 1 ng/ml human TNF- α (Sigma Aldrich, St. Louis, MO, USA), 50 ng/ml LPS purified from *E. coli* O55:B5 (Sigma Aldrich), and TNF- α with LPS (at the same concentrations). As a control, non-stimulated explants ($n = 5$ for each tested group) were treated with the same conditions as the treated cultures. After stimulation for 4 h, explants were placed into cryotubes filled with 1 ml of RNeasy lysis buffer (Qiagen, Crawley, UK). The samples were stored overnight at 4°C ; RNeasy lysis buffer was then removed, and the tissues were stored at -80°C until RNA isolation. The supernatants recovered from the endometrial organ cultures were collected and used for measurement of TNF- α concentrations.

Isolation and culture of feline endometrial cells

Epithelial cells from cats at estrus ($n = 5$), and stromal cells from cats at diestrus ($n = 5$) were isolated from the endometrium according to a published protocol [6, 7]. After reaching 85% confluence, cells were stimulated at doses reported in a previous study [4] as follows: (i) 10 μl of sterile PBS served as a control, (ii) TNF- α (Sigma Aldrich) at 1 ng/ml, (iii) LPS purified from *E. coli* O55:B5 (Sigma Aldrich) at 50 ng/ml, and (iv) TNF- α together with LPS at the same concentrations. After 2, 4, or 12 h of incubation, the supernatant was removed and the cells were suspended in 1 ml of TRIzol Reagent (Gibco-BRL Life Technologies, Karlsruhe, Germany) and stored at -80°C until RNA isolation.

TNF- α ELISA

Concentrations of TNF- α in plasma samples and culture media

Table 1. Primers used for real-time PCR

Gene	Gene name		Primer
<i>RPL17</i>	60S ribosomal protein L17	Forward	CTCTGGTCATTGAGCACATCC
		Reverse	TCAATGTGGCAGGGAGAGC
<i>RPS7</i>	40S ribosomal protein S7	Forward	GTCCCAGAAGCCGCACTTTGAC
		Reverse	CTCTTGCCCACAATCTCGCTCG
<i>TNF</i>	Tumor necrosis factor α	Forward	TGGCCTGCAACTAATCAACC
		Reverse	GTGTGGAAGGACATCCTTGG
<i>TNFR1</i>	TNF- α receptor type 1	Forward	TCTCCTTGACAGTGTACCG
		Reverse	AGGACAGAGGCTGCAGTTTAG

were determined after incubating with LPS for 4 h, using an enzyme immunoassay TNF- α EIA kit (feline TNF- α DuoSet[®] ELISA DEVELOPMENT SYSTEM, R&D Systems, Minneapolis, USA; DY1814) according to the manufacturer's instructions. The standard curve ranged between 15.62 pg/ml and 1000 pg/ml. The intra- and inter-assay coefficients of variation were 7.8% and 12.5%, respectively. Each sample was tested in duplicate. The cross-species reactivity of TNF- α was 34.1% for canine TNF- α , 1.4% for equine TNF- α , 7.5% human TNF- α , and 11.2% for rhesus macaque.

RNA isolation and reverse transcription

TRI-reagent[®] (Sigma Aldrich) was used to isolate total RNA according to the method of total RNA isolation, using a single extraction with an acid guanidinium thiocyanate-phenol-chloroform mixture, described by Chomczynski and Sacchi [17]. RNA concentrations ranged from 200 to 300 ng/ μ l and were quantitated using a NanoDrop ND-1000 spectrophotometer (Thermo Scientific, Wilmington, USA); A_{260/280} ratios ranged from 1.95 to 2.04. DNase treatment was performed using Amplification Grade DNase I (Sigma Aldrich) according to the manufacturer's protocol. Complementary DNA (cDNA) synthesis was performed using the ImProm-II[™] Reverse Transcription System (Promega, Dübendorf, Switzerland) according to the manufacturer's instructions.

Real-Time PCR

All real-time PCR reactions were carried out in duplicate on a Viia 7 machine (Applied Biosystems, Carlsbad, California, USA) as previously described [16]. The reaction mix was set up as follows: 5 μ l GoTaq Green Master Mix (Promega), 1 μ l (0.5 μ M) of each primer, and 3 μ l cDNA (3.33 ng/ μ l). The primers are listed in Table 1; *RPS7* and *RPL17* were chosen as housekeeping genes [18]. All primers were purchased from Genomed S.A. (Warszawa, Poland) (Table 1). Each run included autoclaved water instead of cDNA as a negative control. The amplification was carried out as follows: denaturation for 20 sec at 95°C, followed by 40 cycles of 95°C for 1 sec and 60°C for 20 sec. The presence of product was confirmed by electrophoresis on a 2% agarose gel. Relative mRNA quantification data were then analyzed with the Real-time PCR miner algorithm [19]. According to the instructions supplied by the miner algorithm (<http://www.miner.ewindup.info/>), after determination of average cyclic threshold (Ct) and primer efficiency for each sample, the Ct levels were related to the primer efficiency level (E) using the equation $[1/(1+E)^{Ct}]$.

Thereafter, expression of the target genes was normalized against that of the reference genes and mRNA expression is presented in relation to different conditions in the endometria or to treatments.

Statistics

Data regarding the secretion of TNF- α by the feline endometrium in response to LPS and gene expression of *TNF* and *TNFR1* after stimulation with TNF- α and LPS in endometrial explants were analyzed using a Kruskal-Wallis nonparametric one-way ANOVA followed by the Dunn's Test (GraphPad PRISM v 6.0; GraphPad Software, San Diego, CA, USA). Expression is presented as an n-fold increase or decrease in mRNA level. Statistical significance was defined as $P < 0.05$. The differences in TNF- α in plasma samples, gene expression of *TNF/TNFR1*, and TNF- α secretion by epithelial and stromal cells were analyzed using a Kruskal-Wallis nonparametric one-way ANOVA followed by the Dunn's Test (GraphPad PRISM v 6.0; GraphPad Software) and are presented as the mean ($\pm$ SEM). Statistical significance was defined as $P < 0.05$.

Results

Plasma concentration of TNF- α

Mean TNF- α concentrations were higher in queens at estrus than those determined to be at anestrus or diestrus, or those that were MPA-treated or those that had pyometra ($P < 0.05$) (Fig. 1).

Endometrial response to LPS via TNF- α secretion

LPS-stimulated explants from animals at estrus produced higher amounts of TNF- α than control explants ($P < 0.001$). No changes were found in TNF- α production, after LPS treatment, in endometrial explants from anestrus, mid and late diestrus, or from queens treated short- or long-term with MPA. Differences were also not observed in explants from inflamed uteri (Fig. 2).

Transcription levels of *TNF* and *TNFR1* genes in the feline endometrium

TNF- α ($P < 0.05$) and LPS ($P < 0.001$) treatment increased *TNF* gene expression in endometrial explants during estrus. A higher transcription level of *TNF* mRNA was observed in mid and late diestrus after stimulation with LPS alone or in combination with TNF- α ($P < 0.01$). Expression of *TNF* was higher in explants derived from queens treated short-term with MPA after stimulation with LPS

Fig. 1. Concentration of TNFα in blood plasma of queens in anestrus (A; n = 5), estrus (E; n = 5), and diestrus (D; n = 5), in queens treated with MPA (MPA; n = 5), and in queens with pyometra (P; n = 5). Asterisks indicate statistical differences among groups (* $P < 0.05$). Values are expressed as mean $\pm$ SEM.

($P < 0.01$) or with LPS together with TNF-α ($P < 0.05$). LPS alone ($P < 0.01$) and combined with TNF-α ($P < 0.001$) also increased *TNF* gene expression in endometrial explants from queens treated long-term with MPA. *TNF* gene expression was upregulated by TNF-α upon LPS challenge in inflamed endometrium ($P < 0.001$) (Fig. 3A).

Expression of *TNFR1* was greater in the endometrium during estrus after stimulation with TNF-α ($P < 0.01$) and LPS ($P < 0.001$). LPS decreased *TNFR1* gene expression in endometrial explants from queens at mid diestrus ($P < 0.01$). A higher transcription level of *TNFR1* mRNA was observed in endometria from cats suffering from pyometra after stimulation with LPS combined with TNF-α ($P < 0.01$) (Fig. 3B).

TNF-α production by endometrial epithelial and stromal cells after LPS stimulation

LPS stimulated TNF-α production in epithelial cells after 4 and 12 h ($P < 0.01$), compared to control treated cells (Fig. 4A). However, no changes were found in TNF-α production in stromal cells treated

with LPS (Fig. 4B).

Transcription levels of TNF and TNFR1 genes in cultured cells

Expression of *TNF* was upregulated in epithelial cells after 2 ($P < 0.01$), 4 ($P < 0.05$), and 12 h ($P < 0.01$) of stimulation by LPS compared to control levels (Fig. 4C). Expression of *TNFR1* was increased by LPS treatment in epithelial cells after 2 ($P < 0.05$) and 12 h ($P < 0.01$), but decreased after 4 h ($P < 0.05$), compared to control levels (Fig. 4D). In addition, TNF-α increased *TNFR1* expression in epithelial cells after 4 ($P < 0.05$) and 12 h ($P < 0.01$) of incubation compared to control levels (Fig. 4F).

The expression of *TNFR1* was lower in LPS-treated stromal cells after 2, 4 ($P < 0.01$), and 12 h ($P < 0.001$), compared to control levels (Fig. 4D), whereas expression of *TNF* was increased by TNF-α treatment in stromal cells after 2 ($P < 0.01$), 4 ($P < 0.05$), and 12 h ($P < 0.001$) of incubation (Fig. 4E). No significant changes in *TNFR1* mRNA expression were seen in stromal cells after TNF-α treatment (Fig. 4F).

Discussion

This study found that LPS-challenge increased TNF-α secretion by endometrial tissue solely in estrous queens. However, an elevation of *TNF* mRNA was observed in the endometria of almost every estrous cycle stage (with the exception of anestrus) as well as in the endometria of MPA-treated cats and those suffering from pyometra. Increased cytokine production during estrus could be related to elevated levels of estradiol, enhancing the action of LPS. Treatment of ovariectomized rats with 17β-estradiol previously resulted in an increase in endometrial TNF-α mRNA and protein expression [20]. Furthermore, in human endometrial cells, TNF-α enhanced the intracellular estrogenic milieu by coordinating the expression of enzymes involved in local estrogen biosynthesis and metabolism [21]. A positive feedback loop between estradiol and TNF-α could be crucial for the enhanced proliferation of endometrial cells during the estrous cycle. Estradiol promotes endometrial proliferation of luminal and glandular epithelial cells and regulates the endometrial expression of cytokines as well as growth factors [22]. In turn, TNF-α,

Fig. 2. TNFα secretion by feline endometrium derived from queens in anestrus (n = 5), estrus (n = 5), mid (n = 5) and late diestrus (n = 5), and queens treated short- (n = 5) and long-term with MPA (n = 5) and queens with pyometra (n = 5) in response to LPS. Asterisks indicate statistical differences between treatment and control (*** $P < 0.001$).

Fig. 3. Expression of *TNF* (A) and *TNFR1* (B) in endometrial explants from queens in anestrus ($n = 5$), estrus ($n = 5$), mid ($n = 5$) and late diestrus ($n = 5$), queens treated short-term with MPA ($n = 5$), queens treated long-term with MPA ($n = 5$), and queens with pyometra ($n = 5$) after 4 h incubation with TNF α , LPS, or TNF α together with LPS. Asterisks indicate statistical differences between expression levels depending on the treatment (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

via its two receptors, mediates different cellular responses. TNFR1 is related to cell death and apoptosis, whereas TNFR2 is associated with growth and proliferation [23]. Increased growth of glands, located in the endometrium, and under the influence of estradiol and TNF- α /TNFR2, could be a starting point for the development of cystic endometrial hyperplasia (CEH).

In this study, we confirmed the fundamental role of feline endometrial epithelial cells in pathogen recognition and the subsequent secretion of TNF- α . When activated by exposure to an infectious agent, macrophages and neutrophils recognize pathogen-associated molecular patterns via TLRs, resulting in intracellular signaling, activation of transcription factors, and production of pro-inflammatory cytokines [TNF- α , interleukin 1 (IL-1), and interleukin 6 (IL-6)] [24, 25] as well as secondary inflammatory mediators (prostaglandins, nitric oxide, and oxygen free radicals). Previously it was shown that an increased number of immune cells, including monocytes and neutrophils, in the blood of cats with pyometra, were able to produce TNF- α [26]. However, in our study no changes were observed in plasma TNF- α concentrations. A possible explanation for the lack of increased TNF- α plasma concentration in cats suffering from pyometra may be its short half-life [27]. Moreover, TNF- α has been considered as an early marker of acute endotoxin exposure in experimental animal sepsis [28]. Another study showed that TNF- α

tends to precede the acute-phase response, and its detection was impossible as soon as 6 to 24 h after the induction of inflammation [29]. However, a study of 53 bitches with pyometra found plasma TNF- α concentrations to be significantly different between dogs with pyometra and healthy dogs [12]. The dogs showed a mean onset of pyometra approximately 1 week before veterinary examination; hence, these results were puzzling in the context of the experimental studies [12]. One possible explanation for this observation may be the continuous or intermittent systemic release of inflammatory mediators or bacterial compounds, including LPS. We assume that similar levels of TNF- α found in queens with diagnosed uterine inflammation compared to that of all other groups examined (with the exception of cats at estrus), may be due to the inherent characteristics of pyometra in cats, which is a severe disease. In this regard, our data are in accordance with earlier experimental studies [27–29]. We conclude this part of our study with the assumption that plasma TNF- α concentrations, as opposed to endometrial content, is not a reliable marker of pyometra.

Pyometra is associated with the upregulation of proinflammatory cytokines and chemokines in the inflamed uterus [30]. In a previous study, we clearly demonstrated that TNF- α is expressed mostly in epithelial cells of the feline endometrium. In the inflamed uterus, strong TNF- α immunolabeling was observed in the endometrial

Fig. 4. A, B) Time course of TNFα production by endometrial epithelial (n = 5) and stromal (n = 5) cells after LPS stimulation. C–F) Time course of *TNF* and *TNFR1* mRNA expression in feline epithelial (n = 5) and stromal (n = 5) cells after LPS and TNFα treatment. Values were calculated in relation to controls (CTR) (separately for stromal and epithelial cells taking into account stimulation time) and presented as fold change in gene expression (C–F). Asterisks indicate statistical differences between expression levels depending on the treatment (* P < 0.05, ** P < 0.01, *** P < 0.001).

glands, whereas moderate to weak labeling was shown in the surface epithelium and weak or negative staining was found in the stroma [4]. However, in bitches with pyometra, the endometrial *TNF*-transcript level was not significantly altered [30]. Moreover, in the present study, LPS alone did not affect endometrial *TNF* gene expression or the secretion of TNF-α by feline endometrial explants from inflamed uteri. Only LPS enhanced by TNF-α led to an increase in the endometrial expression of *TNF* and *TNFR1*. A possible explanation for this result may be the severity of endometrial damage during the course of pyometra. In pyometra, serious endometrial epithelium impairment can be observed. However, LPS, alone or in combination with TNF-α, increased *TNF* transcription in the endometria of cats during diestrus and after hormone treatment with MPA. Nevertheless, in the study presented here, LPS had no effect on TNF-α production by endometrial explants, either from diestrus or from MPA-treated endometria. An elevation in *TNF* mRNAs, without a subsequent increase in protein levels may however suggest that the period of incubation (4 h) was not enough to observe changes in secreted TNF-α. This might be

supported by an earlier observation that LPS-challenge led to TNF-α secretion in the endometrial epithelium, after 24 h of incubation [7]. Importantly, by taking advantage of endometrial epithelial or stromal cell cultures (as opposed to studying whole tissues containing other cell types such as immune cells), we could clearly show how these specific cells respond to LPS. The observed increases in *TNF* mRNA levels and TNF-α production by LPS-treated epithelial cells confirms our preliminary data using ELISpot. Therein we clearly demonstrated that feline epithelial cells, in response to LPS, release markedly greater amounts of TNF-α compared to stromal cells [7]. Endometrial stromal cells seem to play only a secondary role in the endotoxin challenge response, as LPS stimulation did not affect TNF-α concentration in these cells when levels were compared to those of untreated control cells. In fact, LPS negatively influenced *TNFR1* expression in stromal cells and did not affect *TNF* mRNA when compared to levels of untreated cells. Since the only effect observed in stromal cells was the TNF-α-induced upregulation of *TNF* and *TNFR1*, one may speculate that stromal cells play a supporting

role in the production of proinflammatory cytokines and chemokines; however, they are not involved in an acute pro-inflammatory response.

To conclude, we provided further support for a fundamental role of endometrial epithelial cells in pathogen recognition and subsequent upregulation of TNF- α production also evident in endometrial tissues obtained at estrus. Endometrial stromal cells seem to play only a secondary role in tissue responses to endotoxin-challenge.

Conflict of interest: The authors declare that they have no competing interests.

Acknowledgments

The present study was performed with support of the Polish Ministry of Scientific Research and High Education (N308 560 740). We thank Dr Anna Rapacz-Leonard and Piotr Socha from the Department of Animal Reproduction with Clinic, Faculty of Veterinary Medicine, University of Warmia & Mazury Olsztyn, Poland, and the Dogmed veterinary clinic from Olsztyn for providing materials used in this study.

References

1. Soboll G, Schaefer TM, Wira CR. Effect of toll-like receptor (TLR) agonists on TLR and microbicidal expression in uterine and vaginal tissues of the mouse. *Am J Reprod Immunol* 2006; **55**: 434–446. [Medline] [CrossRef]
2. Asahina Y, Yoshioka N, Kano R, Morimoto T, Hasegawa A. Full-length cDNA cloning of Toll-like receptor 4 in dogs and cats. *Vet Immunol Immunopathol* 2003; **96**: 159–167. [Medline] [CrossRef]
3. Ignacio G, Nordone S, Howard KE, Dean GA. Toll-like receptor expression in feline lymphoid tissues. *Vet Immunol Immunopathol* 2005; **106**: 229–237. [Medline] [CrossRef]
4. Jursza E, Kowalewski MP, Boos A, Skarzynski DJ, Socha P, Siemieniuch MJ. The role of toll-like receptors 2 and 4 in the pathogenesis of feline pyometra. *Theriogenology* 2015; **83**: 596–603. [Medline] [CrossRef]
5. Silva E, Henriques S, Brito S, Ferreira-Dias G, Lopes-da-Costa L, Mateus L. Oestrous cycle-related changes in production of Toll-like receptors and prostaglandins in the canine endometrium. *J Reprod Immunol* 2012; **96**: 45–57. [Medline] [CrossRef]
6. Jursza-Piotrowska E, Socha P, Skarzynski DJ, Siemieniuch MJ. Prostaglandin release by cultured endometrial tissues after challenge with lipopolysaccharide and tumor necrosis factor α , in relation to the estrous cycle, treatment with medroxyprogesterone acetate, and pyometra. *Theriogenology* 2016; **85**: 1177–1185. DOI [CrossRef]. [Medline]
7. Jursza E, Szóstek AZ, Kowalewski MP, Boos A, Okuda K, Siemieniuch MJ. LPS-challenged TNF α production, prostaglandin secretion and TNF α /TNFRs expression in the endometrium of domestic cats in estrus or diestrus, and in cats with pyometra or receiving medroxyprogesterone acetate. *Med Inflamm* 2014; 689280.
8. Carramolino L, Kremer L, Goya I, Varona R, Buesa JM, Gutiérrez J, Zaballos A, Martínez-A C, Márquez G. Down-regulation of the beta-chemokine receptor CCR6 in dendritic cells mediated by TNF- α and IL-4. *J Leukoc Biol* 1999; **66**: 837–844. [Medline]
9. Vassalli P. The pathophysiology of tumor necrosis factors. *Annu Rev Immunol* 1992; **10**: 411–452. [Medline] [CrossRef]
10. Carswell EA, Old LJ, Kassel RL, Green S, Fiore N, Williamson B. An endotoxin-induced serum factor that causes necrosis of tumors. *Proc Natl Acad Sci USA* 1975; **72**: 3666–3670. [Medline] [CrossRef]
11. Blackwell TS, Christman JW. Sepsis and cytokines: current status. *Br J Anaesth* 1996; **77**: 110–117. [Medline] [CrossRef]
12. Fransson BA, Lagerstedt AS, Bergstrom A, Hagman R, Park JS, Chew BP, Evans MA, Ragle CA. C-reactive protein, tumor necrosis factor α , and interleukin-6 in dogs with pyometra and SIRS. *J Vet Emerg Crit Care* 2007; **17**: 373–381. [CrossRef]
13. Dow C. The cystic hyperplasia-pyometra complex in the bitch. *J Comp Pathol* 1959; **69**: 237–250. [Medline] [CrossRef]
14. Dow C. The cystic hyperplasia-pyometra complex in the cat. *Vet Rec* 1962; **74**: 141–147.
15. Lawler DF, Evans RH, Reimers TJ, Colby ED, Monti KL. Histopathologic features, environmental factors, and serum estrogen, progesterone, and prolactin values associated with ovarian phase and inflammatory uterine disease in cats. *Am J Vet Res* 1991; **52**: 1747–1753. [Medline]
16. Siemieniuch MJ, Jursza E, Kowalewski MP, Majewska M, Skarzynski DJ. Prostaglandin endoperoxide synthase 2 (PTGS2) and prostaglandins F2 α and E2 synthases (PGFS and PGES) expression and prostaglandin F2 α and E2 secretion following oestrogen and/or progesterone stimulation of the feline endometrium. *Reprod Domest Anim* 2013; **48**: 72–78. [Medline] [CrossRef]
17. Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987; **162**: 156–159. [Medline] [CrossRef]
18. Jursza E, Skarzynski DJ, Siemieniuch MJ. Validation of reference genes in the feline endometrium. *Reprod Biol* 2014; **14**: 302–306. [Medline] [CrossRef]
19. Zhao S, Fernald RD. Comprehensive algorithm for quantitative real-time polymerase chain reaction. *J Comput Biol* 2005; **12**: 1047–1064. [Medline] [CrossRef]
20. Russo LA, Peano BJ, Trivedi SP, Cavalcanto TD, Olenchok BA, Caruso JA, Smolock AR, Vishnevsky O, Gardner RM. Regulated expression of matrix metalloproteinases, inflammatory mediators, and endometrial matrix remodeling by 17 β -estradiol in the immature rat uterus. *Reprod Biol Endocrinol* 2009; **7**: 124. [Medline] [CrossRef]
21. Salama SA, Kamel MW, Diaz-Arrastia CR, Xu X, Veenstra TD, Salih S, Botting SK, Kumar R. Effect of tumor necrosis factor- α on estrogen metabolism and endometrial cells: potential physiological and pathological relevance. *J Clin Endocrinol Metab* 2009; **94**: 285–293. [Medline] [CrossRef]
22. Tabibzadeh S. Cytokines and the hypothalamic-pituitary-ovarian-endometrial axis. *Hum Reprod* 1994; **9**: 947–967. [Medline]
23. Hsu H, Xiong J, Goeddel DV. The TNF receptor 1-associated protein TRADD signals cell death and NF-kappa B activation. *Cell* 1995; **81**: 495–504. [Medline] [CrossRef]
24. Robertson CM, Coopersmith CM. The systemic inflammatory response syndrome. *Microbes Infect* 2006; **8**: 1382–1389. [Medline] [CrossRef]
25. Dickson S. Sepsis and multiple organ failure. *Anaesth Intensive Care* 2009; **10**: 165–168. [CrossRef]
26. Hagman R, Karlstam E, Persson S, Kindahl H. Plasma PGF 2 alpha metabolite levels in cats with uterine disease. *Theriogenology* 2009; **72**: 1180–1187. [Medline] [CrossRef]
27. Ruaux CG. Pathophysiology of organ failure in severe acute pancreatitis in dogs. *Compend Contin Educ Pract Vet* 2000; **22**: 531–542.
28. Otto CM, Rawlings CA. Tumor necrosis factor production in cats in response to lipopolysaccharide: an *in vivo* and *in vitro* study. *Vet Immunol Immunopathol* 1995; **49**: 183–188. [Medline] [CrossRef]
29. Yamashita K, Fujinaga T, Miyamoto T, Hagio M, Izumisawa Y, Kotani T. Canine acute phase response: relationship between serum cytokine activity and acute phase protein in dogs. *J Vet Med Sci* 1994; **56**: 487–492. [Medline] [CrossRef]
30. Hagman R, Rönnerberg E, Pejler G. Canine uterine bacterial infection induces upregulation of proteolysis-related genes and downregulation of homeobox and zinc finger factors. *PLoS ONE* 2009; **4**: e8039. [Medline] [CrossRef]