Advance Publication by J-STAGE
Journal of Reproduction and Development

Accepted for publication: August 2, 2021

Advanced Epub: August 26, 2021

The I510V mutation in *KLHL10* in a patient with oligoasthenoteratozoospermia

Running title: The I510V mutation in *KLHL10*

Zicong Huang^{1,#}, Feilong Chen^{2,#}, Minyu Xie¹, Hanbin Zhang¹, Yuge Zhuang¹, Chuyu Huang³,
Xuemei Li⁴, Hong liu^{4,*}, Zhenguo Chen^{1,*}

¹Department of Cell Biology, School of Basic Medical Sciences, Southern Medical
University, Guangzhou 510515, P. R. China.

²Department of Pathology, Panyu Maternal and Child Care Service Centre of Guangzhou,
Guangzhou 511499, P. R. China.

³Center for Reproductive Medicine, Department of Obstetrics and Gynecology, Nanfang
Hospital, Southern Medical University, Guangzhou 510515, P. R. China.

⁴Reproductive center, Affiliated Shenzhen Maternity & Child Healthcare Hospital, Southern
Medical University (Shenzhen Maternity & Child Healthcare Hospital), Shenzhen 518017, P.
R. China.

[#]These authors contributed equally to this work.

^{*}Correspondence: **Hong Liu**, Reproductive center, Affiliated Shenzhen Maternity & Child
Healthcare Hospital, Southern Medical University (Shenzhen Maternity & Child Healthcare
Hospital) (email: zllhhong@126.com); **Zhenguo Chen**, Department of Cell Biology, School
of Basic Medical Sciences, Southern Medical University. (Tel: 86-20-62789095, Fax:
86-20-61648208, email: czg1984@smu.edu.cn).

Acknowledgments

This work was supported by the National Natural Science Foundation of China (31972910,
31571185), the Science, Technology and Innovation Commission of Shenzhen Municipality

23 (JCYJ20190809172209453), and the Health & Family Planning System Research Project of
24 Shenzhen City (SZXJ2017002).

25 **Disclosure Statement:** All the authors have no conflicts of interest.

26

Abstract

Oligoasthenoteratozoospermia is a human infertility syndrome caused by defects in spermatogenesis, spermiogenesis, and sperm maturation, and its etiology remains unclear. Kelch-like 10 (KLHL10) is a component of ubiquitin ligase E3 10 (KLHL10) and plays an important role in male fertility. Deletion or mutation of the *Klhl10* gene in *Drosophila* or mice results in defects in spermatogenesis or sperm maturation. However, the molecular mechanisms by which KLHL10 functions remain elusive. In this study, we identified a missense mutation (c.1528A→G, p.I510V) in exon 5 of *KLHL10*, which is associated with oligoasthenoteratozoospermia in humans. To investigate the effects of this mutation on KLHL10 function and spermatogenesis and/or spermiogenesis, we generated mutant mice duplicating the amino acid conversion using the clustered regularly interspaced palindromic repeat/cas9 (CRISPR/Cas9) system and designated them *Klhl10*^{I510V} mice. However, the *Klhl10*^{I510V} mice did not exhibit any defects in testis development, spermatogenesis, or sperm motility at ten-weeks-of-age, suggesting that this mutation does not disrupt the KLHL10 function, and may not be the cause of male infertility in the affected individual with oligoasthenoteratozoospermia.

Keywords: oligoasthenoteratozoospermia; KLHL10; missense mutation; CRISPR/Cas9; spermatogenesis

1 Introduction

Infertility is a major health problem worldwide, with a prevalence in developed and developing countries ranging from 3.5% to 16.7%, and male factors account for approximately half of the cases[1, 2]. Disorders in spermatogenesis are the principal causes of male infertility. Spermatogenesis is a process in which spermatogonial stem cells undergo a series of mitosis, meiosis, and morphological changes to form mature sperm, involving more than 2000 genes[3]. In the late stage of spermatogenesis, the process of round spermatozoa forming and growing to sperm for release is called spermiogenesis, including nuclear condensation, acrosome formation, tail formation, and residual body loss[4-6]. Spermiogenesis plays a crucial role in sperm maturation and motility, as well as in final sperm-egg fertilization. To date, dozens of genes have been reported to regulate this process. For example, previous studies have shown that SEPTIN12 regulates the morphogenesis of sperm heads and the elongation of sperm tails, probably implicating its association with α - and β -tubulins[7], and its family is required for the structural integrity and motility of the sperm tail during postmeiotic differentiation[8]. Another study identified a protein located on the plasma membrane of sperm flagellum, CatSper, as a “sperm related cation channel” protein, which plays an important role in sperm motility[9, 10]. However, the mechanisms underlying sperm deformation are still not completely understood.

Ubiquitination plays a key role in eliminating unnecessary proteins in sperm during the middle and late stages of sperm deformation[11, 12]. Kelch-like 10 (KLHL10), characterized by an amino terminal BTB/POZ domain and kelch repeats at the carboxyl terminus, is a member of a large BTB (Bri-a-brac, Tramtrack, and Broad-Complex)-kelch protein

superfamily[12]. A previous study showed that male mice with a null mutation in one allele of *Klh10* were infertile, displaying asynchronous spermatid maturation, degeneration of late spermatids, and significant reduction in the number of late spermatids, indicating that normal KLHL10 levels are essential for sperm maturation and male fertility[13]. As a testis-specific protein, KLHL10 is expressed in elongated and elongated sperm. In *Drosophila*, it interacts functionally and physically with Cul3_{Testis} and Roc1b to form ubiquitin ligase E3 to promote caspase activation and spermatid individualization[14], and specific point mutations altering conserved amino acid residues in the Kelch domain of Klhl10 blocked caspase activation and spermatid individualization, causing excessive sperm death[14, 15]. In addition, a number of point mutations in *KLHL10* have been identified in clinical patients with azoospermia or oligoasthenospermia[16, 17]. These studies demonstrated that KLHL10 plays an essential role in spermatogenesis and spermiogenesis. However, the molecular mechanisms by which KLHL10 functions are still largely unknown, except for a model in which KLHL10 complexes with CUL3 act as a testis-specific ubiquitin E3 ligase, in which KLHL10 is postulated to orientate substrate protein for degradation[12].

In this study, we performed the exome analysis of male infertility-related candidate genes and identified a heterogeneous mutation (c.1528A→G, p.I510V) in exon 5 of the *KLHL10* gene in a patient with oligoasthenoteratozoospermia. To explore the clinical and biological significance of this point mutation in male fertility, we duplicated this point mutant in mice; however, the mutant mice did not exhibit any disorder in male fertility at the age of ten weeks.

2 Materials and Methods

2.1 Clinical information

A couple (both aged 34) suffering from infertility for four years turned to our reproductive center for assistance with reproductive technology. Laboratory examinations showed that the wife had normal hormone levels and regular menstrual cycles, and ultrasonic examinations also indicated normal size and morphology of the uterus, normal follicular development, and ovulation. However, the husband demonstrated oligoasthenoteratozoospermia by sperm quality analysis (Table 1). He had no history of diseases that may affect male fertility and had normal somatic karyotypes. To better diagnose the potential genetic cause, the husband approved to participate in exome analysis of the candidate genes (Table 2) that are known to be associated with male infertility. Informed consent was obtained from the patient to report this case, and the study was approved by the Ethics Committee of Shenzhen Maternity and Child Healthcare Hospital and Southern Medical University.

2.2 Human blood sampling and exome for candidate genes

Blood samples were obtained from the patient (5 ml taken into ethylenediaminetetraacetic acid (EDTA) by venipuncture) and immediately stored at -80°C until detection. Genomic DNA isolation, polymerase chain reaction (PCR), and subsequent high-throughput sequencing combined with bioinformatics analysis were performed by Yikon Genomics (Shanghai, China) to screen for potential mutations in the male infertility-related genes (Table 2).

2.3 Targeted point mutation of *Klhl10* (I510V) in mice using the clustered regularly interspaced palindromic repeat/caspase 9 (CRISPR/Cas9) system

Klhl10 mutant mice (c.1528A→G, p.I510V) were generated by Shanghai Model Organisms (Shanghai, China), using CRISPR/Cas9 technology to introduce the target point mutations by

homologous recombination repair. Briefly, the Cas9 mRNA and guide RNA were obtained by in vitro transcription, oligo donor DNA was obtained by synthesis, and cas9 mRNA, gRNA, and donor DNA were microinjected into one-cell stage fertilized eggs of C57BL/6J mice to obtain F0 generation mice. A total of 225 eggs were injected, and seven chimeric F0 mice were identified by two-round genotyping by PCR and subsequent sequencing. F1 mice were obtained by mating F0 mice with C57BL/6J mice, followed by mating F1 mice to generate homozygous F2 generation. DNA was isolated from tail biopsies and used to genotype the mice, followed by PCR product sequencing (Sanger sequencing, Sangon, Shanghai). The gRNA sequence was 5'-TACGAGGGTTAAACATAGTG-3'. The primer sequences for genotyping were as follows: forward, 5'-ACCTGTACCTGTCTTCTGTCTA-3', reverse: 5'-ATCCCCTTCCTTACTCCCTCATT-3'. The forward primer was also used as the sequencing primer. Male mice with complete amino acid substitutions were designated as Khl10^{I510V} mice, and littermate wild-type (WT) males were used as controls and were designated as Khl10^{WT} mice. All mice were housed in a clean, temperature-controlled environment with a 12-h light/dark cycle, and were given free access to regulate laboratory chow diet and water for housing in regular cages.

2.4 Tissue collection and histological analysis

Following euthanasia, the testes and epididymides of mice were removed, weighed, fixed in modified Davidson fixative, and processed in paraffin according to standard methods. At least three sections from each testis and epididymis (5 μ m, taken 100 μ m apart) were stained with hematoxylin and eosin (H&E) for regular histological examination.

2.5 Sperm concentration and motility analysis

Human sperms were collected from the patient via masturbation and incubated at 37 ° C for 20 min for liquefaction. For mice, one epididymis from each mouse was removed and minced in M2 medium (0.5 mL; Sigma, M7167-100ML) at 37 ° C for 15 min to allow the sperm to be released into the medium. Sperm concentration was assessed with a hemocytometer, and sperm motility was analyzed using a microscope and CASA (Hamilton Thorne IVOS software) under parameters for humans and mice, respectively.

2.6 Statistical analysis

All experiments were performed in triplicate. Data were expressed as the mean values $\pm$ standard deviation (SD), and the between-group differences were analyzed using the *t*-test for measurable data and the Chi-square test for percentage data (SPSS 13.0). Statistical significance was set at $p < 0.05$.

3 Result

3.1 Identification of a missense mutation in *KLHL10* in a patient with oligoasthenoteratozoospermia

Sperm quality analysis showed that the husband exhibited markedly decreased sperm concentration and motility and 100% sperm with abnormal morphology (Table 1). H&E staining of the smeared sperm further confirmed the abnormal morphology in sperm, characterized by round or irregular head, and/or short tail (Figure 1), indicative of oligoasthenoteratozoospermia in the patient. To explore the potential genetic cause for this affected individual, we performed exome combined with bioinformatics analysis on a panel of 75 candidate genes [18-20], which have been demonstrated to be strongly associated with

male infertility (Table 2). Interestingly, we identified a heterogeneous point mutation at nucleotide 1528(A→G) (located in exon 5) in *KLHL10*, according to GenBank entry NM_152467. The transition generated a missense amino acid substitution, I510V, located in the kelch-5 domain of the KLHL10 protein (Table 3; Figure 2A). However, due to the patient's family reasons, it was impossible to trace the hereditary mode. Point mutations in the human *KLHL10* gene have been reported in the Genome Aggregation Database (gnomAD) in 0.015% of the normal East Asian individuals tested. However, we hypothesized that this may be the cause of oligoasthenoteratozoospermia in the affected individual.

*3.2 Generation of *Klhl10* mutant mice duplicating the missense mutation (c.1528A→G, p.I510V) by CRISPR/Cas9*

We then blasted the amino acid sequences of KLHL10 among the species and found that the amino acid sequences adjacent to the mutant site were highly conserved among species. Thus, to elucidate the effect of this missense mutation on the function of KLHL10 as well as on spermatogenesis in vivo, we duplicated the mutation (c.1528A→G, p.I510V) in mice using CRISPR/Cas9 (Figure 2B). By breeding the F1 mice, followed by PCR and sequencing, we obtained homozygous *Klhl10*^{I510V} mutant mice and their littermate controls (*Klhl10*^{WT}). (Figure 2C-E). These data indicated the successful generation of *Klhl10* mutant mice with target point mutations (1528A→G, I510V), and this point mutation did not cause any splicing abnormality in *Klhl10*.

*3.3 *Klhl10*^{I510V} mice do not exhibit any defects in testis development, spermatogenesis, or sperm motility at ten-weeks-of-age*

The *Klhl10*^{I510V} mice had no obvious difference in physical appearance or body weight at

any age during the experiments, compared with their control littermates (data not shown). Similarly, there was no significant difference in testis weight between the $Klhl10^{I510V}$ and $Klhl10^{WT}$ males at ten-weeks-of-age (Figure 3A). Further morphological examination showed that the $Klhl10^{I510V}$ and $Klhl10^{WT}$ males had indistinguishable seminiferous tubular development, manifested as comparable tubular size, seminiferous epithelium thickness, and cell architecture (Figure 3B). These observations indicated that the I510V transversion in $Klhl10$ may not disturb tubular or testis development in mice. Accordingly, the $Klhl10^{I510V}$ and $Klhl10^{WT}$ males had parallel sperm concentrations (Figure 3C). In line with the observations of the seminiferous epithelium, the $Klhl10^{I510V}$ males exhibited normal spermatogenesis (Figure 3D). Moreover, the $Klhl10^{I510V}$ mice also showed normal sperm motility and morphology (Figure 3E and F), and had similar litter sizes compared with $Klhl10^{WT}$ mice (Figure G). Taken together, these results suggest that the I510V substitution in $Klhl10$ does not impair testis development, spermatogenesis, or sperm motility in mice at ten-weeks-of-age.

4 Discussion

Spermatogenesis encompasses a complex network of processes that occur in the seminiferous tubules and culminates in the production of mature male gamete[21]. Thus, spermatogenesis is strictly controlled and requires tight coordination of numerous genes. There are dozens of genes with abnormal expression levels or mutations in humans or mice have been reported to cause male reproductive disorders. Therefore, screening potential mutations of candidate genes in the pedigree of patients with reproductive disorders is

important, not only for the etiologic and prenatal diagnosis, but also for the methods used for assisted reproductive technology. In this study, we screened potential mutation(s) in 75 candidate genes in a patient with oligoasthenoteratozoospermia and identified a heterogeneous *KLHL10* gene at nucleotide 1528 (A→G) (located in exon 5), which produces a missense amino acid substitution (I510V) located in the kelch-5 domain of KLHL10 protein. Thus, we speculated that this missense mutation in exon 5 of KLHL10 is associated with oligoasthenoteratozoospermia in humans.

As a testis-specific gene, KLHL10 is expressed in elongating and elongated sperm, and has been demonstrated to be essential for male fertility as haploinsufficiency of *Klhl10* gene in mice disrupted spermatogenesis and caused male infertility[13]. Subsequent studies showed that deletion or mutations of *Klhl10* resulted in male infertility in *Drosophila*[14, 15], and mechanistically, Klhl10 regulated spermatogenesis via ubiquitination-mediated target protein degradation[12, 14, 15, 22]. In addition, a high frequency of mutations in *KLHL10* was also identified in infertile men with azoospermia or oligozoospermia [16, 17]. Therefore, KLHL10 plays a critical role in spermatogenesis and male fertility.

It has been well evidenced that KLHL10 and CUL3 form a complex, in which KLHL10, specifically, its six kelch repeats, acts to recognize substrate proteins for ubiquitination-dependent degradation during sperm maturation[14, 15, 22]. In the present study, we identified a missense mutation in an infertile male with oligoasthenoteratozoospermia, which produces an I510V substitution located in the Kelch 5 domain in the KLHL10 protein. However, the mutant mice harboring such an amino acid transversion failed to replicate the infertile phenotypes in the patient, without any defects observed in testis development,

spermatogenesis, or sperm maturation. These observations are contradictory to a previous study in *Drosophila* showing that five single mutations located in each kelch domain in *Drosophila* Klhl10 protein blocked spermatid individualization[14]. We hypothesized that this discrepancy may be due to species differences. Another explanation is that the remaining intact five-kelch repeats may compensate for the inactivation of kelch 5, or the I510V conversion was not sufficient to disrupt the function of kelch 5 in mice. Previously, an A313T mutation located in Kelch 1 of KLHL10 identified in an oligozoospermic man was found to fail to form a homodimer via a yeast two-hybrid assay; however, the significance of this homodimer formation for KLHL10 function was not illustrated in that study[17]. In addition, as both Ile and Val are hydrophobic-aliphatic amino acids, the substitution of Ile with Val may have little effect on protein conformation. Interestingly, chicken KLHL10 has V508, which corresponds to I510 of human KLHL10 and mouse Klhl10. Thus, in light of evolution, replacement of Ile/Val at this residue seems to be acceptable for the function of KLHL10 protein. The last explanation is that there may be other genes that are not included in the panel of 75 candidate genes that are responsible for infertility in the affected individual. Our study suggests that generating a mouse model mimicking gene mutations in patients is an important step in evaluating the significance of variants associated with human diseases. Nevertheless, our report does not deny the importance of KLHL10 in male fertility, but rather calls for further in-depth studies to unveil the mechanism by which KLHL10 recognizes the substrate proteins for degradation as well as its targets.

References

1. **Jacky B, Laura B, Collins JA, Nygren KG.** International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Human Reproduction* 2007; 1506-12.
2. **Zorrilla M, Yatsenko AN.** The Genetics of Infertility: Current Status of the Field. *Current Genetic Medicine Reports* 2013; 1: 247-260.
3. **O'Brien K, Varghese AC, Agarwal A.** The genetic causes of male factor infertility: A review. *Fertility and Sterility* 2010; 93: 1-12.
4. **Chu DS, Shakes DC.** Spermatogenesis. *Adv Exp Med Biol* 2013; 757: 171-203.
5. **Gadella BM, Luna C.** Cell biology and functional dynamics of the mammalian sperm surface. *Theriogenology* 2014; 81: 74-84.
6. **Hao SL, Ni FD, Yang WX.** The dynamics and regulation of chromatin remodeling during spermiogenesis. *Gene* 2019; 706:201-210.
7. **Kuo PL, Chiang HS, Wang YY, Kuo YC, Chen MF, Yu IS, Teng YN, Lin SW, Lin YH.** SEPT12-microtubule complexes are required for sperm head and tail formation. *Int J Mol Sci* 2013; 14: 22102-22116.
8. **Yung-Che, Kuo, Yi-Ru, Shen, Hau-Inh, Chen, Ying-Hung, Lin, Ya-Yun, Wang.** SEPT12 orchestrates the formation of mammalian sperm annulus by organizing core octameric complexes with other SEPT proteins. *Journal of Cell Science* 2015;128:923-934.
9. **Struener T, Goodwin N, Brenker C, Kashikar ND, Weyand I, Seifert R, Kaupp UB.** The CatSper channel mediates progesterone-induced Ca²⁺ influx in human sperm. *Nature* 2011; 471: 382-386.
10. **Sun XH, Zhu YY, Wang L, Liu HL, Ling Y, Li ZL, Sun LB.** The Catsper channel and its roles in male fertility: a systematic review. *Reproductive Biology & Endocrinology Rb & E* 2017; 15: 65.
11. **Upadhyay RD, Kumar AV, Ganeshan M, Balasinor NH.** Tubulobulbar complex: Cytoskeletal remodeling to release spermatozoa. *Reproductive Biology & Endocrinology* 2012; 10: 27.
12. **Wang S, Zheng H, Yumiko E, Fiona K, Wei Y.** Cullin3 Is a KLHL10-Interacting Protein Preferentially Expressed During Late Spermiogenesis. *Biology of Reproduction* 2006: 102-108.
13. **Yan W, Ma L, Burns, Kathleen H, Matzuk, Martin M.** Haploinsufficiency of keich-like protein homolog 10 causes infertility in male mice. *Proceedings of the National Academy of Sciences of the United States of America* 2004;101:7793-7798.
14. **Arama E, Bader M, Rieckhof GE, Steller H.** A ubiquitin ligase complex regulates caspase activation during sperm differentiation in Drosophila. *PLoS Biology* 2007; 5: e251-.
15. **Kaplan Y, Gibbs-Bar L, Kalifa Y, Feinstein-Rotkopf Y, Arama E.** Gradients of a ubiquitin E3 ligase inhibitor and a caspase inhibitor determine differentiation or death in spermatids. *Developmental Cell* 2010; 19: 160-173.
16. **Qiu QM, Gang L, Li WN, Shi QW, Lu GX.** [Mutation of KLHL-10 in idiopathic infertile males with azoospermia, oligospermia or asthenospermia]. *Zhonghua nan ke xue = National journal of andrology* 2009; 15: 974-979.

- 287 17. **Yatsenko AN, Roy A, Chen R, Lang M, Murthy LJ, Wei Y, Lamb DJ, Matzuk, M. M.**
288 Non-invasive genetic diagnosis of male infertility using spermatozoal RNA: KLHL10
289 mutations in oligozoospermic patients impair homodimerization. *Human Molecular Genetics*
290 2006; 15: 3411.
- 291 18. **Cannarella R, Condorelli RA, Duca Y, La Vignera S, Calogero AE.** New insights into the
292 genetics of spermatogenic failure: a review of the literature. *Human Genetics*
293 2019;138:125-140.
- 294 19. **Krausz C, Casamonti E.** Spermatogenic failure and the Y chromosome. *Human Genetics*
295 2017; 136: 637-655.
- 296 20. **Ray, P., F., Coutton, C., Toure, A., Mitchell, M., J.** Single gene defects leading to sperm
297 quantitative anomalies. *Clinical Genetics An International Journal of Genetics in Medicine*
298 2017;91:208-216.
- 299 21. **Neto F, Bach PV, Najari BB, Li PS, Goldstein M.** Spermatogenesis in humans and its
300 affecting factors. *Seminars in Cell & Developmental Biology* 2016; 59: 10-26.
- 301 22. **Aram L, Braun T, Braverman C, Kaplan Y, Ravid L, Levin-Zaidman S, Arama E.** A
302 Krebs Cycle Component Limits Caspase Activation Rate through Mitochondrial Surface
303 Restriction of CRL Activation. *Developmental Cell* 2016: 15-33.

304

Figure legends

Figure 1. Morphology of sperm via hematoxylin and eosin (H&E) staining. (A) Normal human sperm; (B) Representative images showing the irregular sperm head or tail in the patient. Scale bars: 10 μm for (A) and 5 μm for (B).

Figure 2. Targeted point mutation (1528A→G, I510V) of Kelch-like 10 (*Klhl10*) in mice by the clustered regularly interspaced palindromic repeat/caspase 9 (CRISPR/Cas9) system. (A) Alignment of the KLHL10 amino acid sequence. (B) Schematic diagram of the mutant mouse construction strategy. *Klhl10* consists of five exons. (C) Genotyping of the F2 generations by polymerase chain reaction (PCR). (D) Nucleoside acid base peak diagram for PCR products sequencing. (E) Blasting for the PCR products sequence demonstrates homogeneous target mutation in the *Klhl10*^{I510V} mice. Red square indicates the mutation site.

Figure 3. The I510V substitution in *Klhl10* does not impair spermatogenesis or sperm motility in mice at ten-weeks-of-age. (A) Similar *Klhl10*^{WT} and *Klhl10*^{I510V} testis weight. Data are presented as the mean $\pm$ standard deviation (SD), $n = 5$. (B) Testes morphology by H&E staining. (C) No significantly reduced epididymal sperm (intact spermatozoa plus sperm flagella) concentration in *Klhl10*^{I510V} mice compared with wild-type (WT) males. Data are presented as the mean $\pm$ SD, $n = 5$. (D) Epididymis tubular morphology by H&E staining. (E) No significantly reduced total motility of *Klhl10*^{I510V} epididymal headless spermatozoa. Data are presented as the mean $\pm$ SD, $n = 5$. (F) Analysis of litter size. Comparison of *Klhl10*^{WT} with *Klhl10*^{I510V} mice, where no significant difference in the average number of pups per litter.

327 Data are presented as the mean $\pm$ SD, $n = 15$. (G) Detailed morphology of the smeared sperm
328 by H&E staining. Scale bars: 50 μ m for (B) and (D) 20 μ m for (G).
329

330 Table 1. Sperm parameters of the individual with oligoasthenoteratozoospermia.

Sperm characteristics	Parameter	WHO5 criteria
Sperm volume (ml)	2.5	≥ 1.5
Sperm concentration (10^6 / ml)	2.9	≥ 15
Motility (%)	16.7	≥ 40
Precession rate (PR, %)	16.7	≥ 32
Norprecession rate (NR, %)	0	
Immotility (%)	83.3	
Abnormal sperm morphology (%)	100	
Vitality (eosin staining, %)	27	$\geq 58\%$

331

332

Table 2. A panel of human male infertility genes.

<i>AR</i>	<i>ARMC4</i>	<i>AURKC</i>	<i>AZF</i>	<i>BMP15</i>	<i>CATSPER1</i>	<i>CCDC103</i>
<i>CCDC114</i>	<i>CCDC151</i>	<i>CCDC39</i>	<i>CCDC40</i>	<i>CFTR</i>	<i>CHD7</i>	<i>DNAAF1</i>
<i>DNAAF2</i>	<i>DNAH11</i>	<i>DNAH5</i>	<i>DNAI1</i>	<i>DNAI2</i>	<i>DPY19L2</i>	<i>DUSP6</i>
<i>EIF2B2</i>	<i>EIF2B4</i>	<i>EIF2B5</i>	<i>ERCC6</i>	<i>FEZF1</i>	<i>FGF17</i>	<i>FGF8</i>
<i>FGFR1</i>	<i>FIGLA</i>	<i>FII</i>	<i>FLRT3</i>	<i>FMR1</i>	<i>FOXL2</i>	<i>FSHR</i>
<i>FV</i>	<i>GNRH1</i>	<i>GNRHR</i>	<i>HFM1</i>	<i>IL17RD</i>	<i>INSL3</i>	<i>KAL1</i>
<i>KISS1</i>	<i>KISS1R</i>	<i>KLHL10</i>	<i>LHB</i>	<i>LHCGR</i>	<i>LRRC6</i>	<i>MCM8</i>
<i>NANOS1</i>	<i>NOBOX</i>	<i>NR5A1</i>	<i>POF1B</i>	<i>POR</i>	<i>PROK2</i>	<i>PROKR2</i>
<i>PSMC3IP</i>	<i>RSPH1</i>	<i>RSPH4A</i>	<i>SLC26A8</i>	<i>SPAG1</i>	<i>SPATA16</i>	<i>SPRY4</i>
<i>STAG3</i>	<i>SYCE1</i>	<i>SYCP3</i>	<i>TAC3</i>	<i>TACR3</i>	<i>TAF4B</i>	<i>TEX11</i>
<i>TUBB8</i>	<i>WDR11</i>	<i>ZMYND10</i>	<i>ZP1</i>	<i>GAS8</i>		

Abbreviation: AR, Androgen receptor; ARMC4, Armadillo repeat containing 4; AURKC, Aurora kinase C; AZF, Azoospermia factor 1; BMP15, Bone morphogenetic protein 15; CATSPER1, Cation channel, sperm associated 1; CCDC103, Coiled-coil domain containing 103; CCDC114, Coiled-coil domain containing 114; CCDC151, Coiled-coil domain containing 151; CCDC39, Coiled-coil domain containing 39; CCDC40, Coiled-coil domain containing 40; CFTR, Cystic fibrosis transmembrane conductance regulator; CHD7, Chromodomain helicase DNA binding protein 7; DNAAF1, Dynein, axonemal assembly factor 1; DNAAF2, Dynein, axonemal assembly factor 2; DNAH11, Dynein, axonemal, heavy chain 11; DNAH5, Dynein, axonemal, heavy chain 5; DNAI1, Dynein intermediate chain 1, axonemal; DNAI2, Dynein intermediate chain 2, axonemal; DPY19L2, Probable C-mannosyltransferase DPY19L2; DUSP6, Dual specificity protein phosphatase 6; EIF2B2, Translation initiation factor eIF-2B subunit beta; EIF2B4, Translation initiation factor eIF-2B subunit delta; EIF2B5, Translation initiation factor eIF-2B subunit epsilon; ERCC6, DNA excision repair protein ERCC-6; FEZF1, Fez family zinc finger protein 1; FGF17, Fibroblast growth factor 17; FGF8, Fibroblast growth factor 8; FGFR1, Fibroblast growth factor receptor 1; FIGLA, Factor in the germline alpha; FII, Coagulation factor II; FLRT3, Leucine-rich repeat transmembrane protein FLRT3; FMR1, Synaptic functional regulator FMR1; FOXL2, Forkhead box protein L2; FSHR, Follicle-stimulating hormone receptor; FV,

352 Friend virus susceptibility; GNRH1, Progonadoliberin-1; GNRHR, Gonadotropin-releasing
 353 hormone receptor; HFM1, Probable ATP-dependent DNA helicase HFM1; IL17RD,
 354 Interleukin-17 receptor D; INSL3, Insulin-like 3; KAL1, Anosmin-1; KISS1,
 355 Metastasis-suppressor KiSS-1; KISS1R, kiSS-1 receptor; KLHL10, Kelch-like protein 10;
 356 LHB, Lutropin subunit beta; LHCGR, Lutropin-choriogonadotropic hormone receptor;
 357 LRRC6, Protein tilB homolog; MCM8, DNA helicase MCM8; NANOS1, Nanos homolog 1;
 358 NOBOX, Homeobox protein NOBOX; NR5A1, Steroidogenic factor 1; POF1B, Protein
 359 POF1B; POR, NADPH--cytochrome P450 reductase; PROK2, Prokineticin-2; PROKR2,
 360 Prokineticin receptor 2; PSMC3IP, Homologous-pairing protein 2 homolog; RSPH1, Radial
 361 spoke head 1 homolog; RSPH4A, Radial spoke head protein 4 homolog A; SLC26A8, Testis
 362 anion transporter 1; SPAG1, Sperm-associated antigen 1; SPATA16,
 363 Spermatogenesis-associated protein 16; SPRY4, Protein sprouty homolog 4; STAG3, Cohesin
 364 subunit SA-3; SYCE1, Synaptonemal complex central element protein 1; SYCP3,
 365 Synaptonemal complex protein 3; TAC3, Tachykinin-3; TACR3, Neuromedin-K receptor;
 366 TAF4B, Transcription initiation factor TFIID subunit 4B; TEX11, Testis-expressed protein 11;
 367 TUBB8, Tubulin beta-8 chain; WDR11, WD repeat-containing protein 11; ZMYND10, Zinc
 368 finger MYND domain-containing protein 10; ZP1, Zona pellucida sperm-binding protein 1;
 369 GAS8, Dynein regulatory complex subunit 4.

370

Table 3. Mutation information of the affected individual.

Gene	Chromosome position	Transcript	Nucleotide change	Amino acid change	Zygote type
<i>KLHL10</i>	chr17: 40004260	NM_152467	c.1528A>G	p.I510V	Heterozygous

A

<i>Homo sapiens</i>	481	HVYAVGGFDGANRLRS AEAYSPVANTWR	TIP	TMFNPRSNFGIEVVDD	527
<i>Mus musculus</i>	481	HVYAVGGFDGANRLRS AEAYSPVANTWR	TIP	TMFNPRSNFGIEVVDD	527
<i>Rattus norvegicus</i>	481	HVYAVGGFDGANRLRS AEAYSPVANTWR	TIP	TMFNPRSNFGIEVVDD	527

B

C

D

E

Query	698	TCAACCTTGTTTCTTTTGCTCTTCACAGGTGGGTGGCTTTGATGGAGCCAATCGACTGAG	757
Sbjct	63	TCAACCTTGTTTCTTTTGCTCTTCACAGGTGGGTGGCTTTGATGGAGCCAATCGACTGAG	122
Query	758	GAGTGCAGAAAGCCTACAGCCAGTGGCTAATACTTGGCGCACGATCCCACTATGTTTAA	757
Sbjct	123	GAGTGCAGAAAGCCTACAGCCAGTGGCTAATACTTGGCGCACGATCCCACTATGTTTAA	182
Query	818	CCCTCGTAGCAATTTTGGGATCGAGGTAGTGGACGACCTCTTGTTTGTGGTGGGTGGCTT	757
Sbjct	183	CCCTCGTAGCAATTTTGGGATCGAGGTAGTGGACGACCTCTTGTTTGTGGTGGGTGGCTT	242
Query	878	TAATGGCTTTACCACCACCTTTAACGTTGAGTGCTATGATGAAAAGACCGATGAGTGGTA	757
Sbjct	243	TAATGGCTTTACCACCACCTTTAACGTTGAGTGCTATGATGAAAAGACCGATGAGTGGTA	302

A

B

C

D

E

F

G

Khl10^{I510V}