An Ecological Study on *Edwardsiella tarda* in Flounder Farms

M. Mamnur Rashid, Kazumitsu Honda, Toshihiro Nakai
and Kiyokuni Muroga*

Faculty of Applied Biological Science, Hiroshima University,
Kagamiyama, Higashi-Hiroshima 724, Japan

(Received June 16, 1994)

An ecological investigation on *Edwardsiella tarda* was carried out in two Japanese flounder (*Paralichthys olivaceus*) farms in Fukuyama City from April to December 1992. *E. tarda* was isolated from 86% of water, 44% of sediment and 14% of fish samples in one pond, and 22% of water, 0% of sediment and 2% of fish samples in the other pond, although epizootics caused by *E. tarda* were not observed in either pond. Serological studies proved that all the isolates of *E. tarda* from flounder and its environment shared a common heat stable O-antigen. Almost all the isolates tested exhibited pathogenicity to flounder by intramuscular injection. In survivability tests of selected isolates in sea water and freshwater, they perished within 7 days in sea water but survived longer period in freshwater.

These results suggest that there is only one serotype of *E. tarda* which has pathogenicity to flounder in the environment of flounder farms.

Edwardsiella tarda has been known to be pathogenic for a wide variety of animals including several species of fishes (Austin and Austin, 1993). In Japan, the disease (edwardsiellosis) caused by this pathogen constitutes an economical threat particularly in eel culture in freshwater (Hoshina, 1962; Wakabayashi and Egusa, 1973a, b) and flounder culture in sea water (Yasunaga *et al.*, 1982; Nakatsugawa, 1983; Kodama *et al.*, 1987).

Ecological studies on this pathogen in eel culture ponds were made (Minagawa *et al.*, 1983; Park *et al.*, 1983) and *E. tarda* was detected from healthy eels and their culture environment. Distribution of the pathogen in a flounder farm was also studied (Kanai *et al.*, 1988), but the pathogen was not detected from flounder culture environment except for the period when edwardsiellosis prevailed.

In the present work, an ecological survey of the pathogen in apparently healthy flounder (*Paralichthys olivaceus*) and their culture environment was made. Physiological, serological and pathological characteristics of isolated *E. tarda* were also studied.

Materials and Methods

Field sampling

Monthly samplings were done in two flounder farms (Pond A and Pond B) in Fukuyama City, Hiroshima Prefecture, from April through December 1992. In both farms, fish were reared in land based ponds throughout that period. Fish (10 fish), pond waters, supply waters (sand-filtered coastal sea water), sediments and foods for the fish were sampled from each farm. Fish were carried to the laboratory in Hiroshima University (Higashi-Hiroshima City) in oxygen-filled plastic bags or in ice boxes spending 2 or 3 h. Water, mud and food samples were also carried in ice boxes.

Bacterial isolation

Fish: At the first sampling in April, when average body weight of fish was 0.1 g, fish were homogenized individually with 2 ml of sterilized physiological saline (0.85% NaCl: PS). A 0.1 ml of each original homogenate, 10- and 100-fold dilutions with PS were inoculated in duplicate onto *Salmonella-Shigella* (SS) agar (Eiken) for colony count (direct plating culture). From the second sampling in May, with larger fish, the above method was continued for the intestine samples instead of the whole body. In

* Author to whom correspondence should be addressed.

addition, 1 ml of the intestine, liver, spleen, and kidney homogenates (sample:PS=1:2) was put into 10 ml double-strength SS broth (DSSS) as a selective enrichment medium for *E. tarda* (Wyatt *et al.*, 1979) (enrichment culture).

Water: Waters, undiluted, 10- and 100-fold diluted, were inoculated onto SS agar for colony count. One ml of undiluted water samples were subjected to enrichment culture with DSSS. From the third sampling in June, each water sample of 100 ml was passed through a membrane filter (0.45 μ m, Millipore) and that filter was subjected to enrichment culture.

Sediment and food: Direct plating and enrichment culture were done for sediment samples and only enrichment culture was performed for foods (pelleted artificial feed).

Post-incubation works: Inoculated SS agar were incubated at 25°C for 2 days and *E. tarda*-like colonies were selected by their characteristic appearance, transparent colonies with black center. For selective enrichment culture, after 2 days of incubation in DSSS at 25°C, one loopful of each culture was streaked onto SS agar, followed by incubation for 2 days at the same temperature. *E. tarda*-like colonies were again streaked onto SS agar to obtain pure culture, and stocked in 1/3 strength heart infusion (HI) agar (Eiken) for further examinations.

Characterization of isolates

Morphological and biochemical characteristics: Rapid identification of the isolated bacteria was performed using triple sugar iron (TSI), sulfide-indole-motility (SIM), lysine, and mannitol test media, and ONPG disks, as suggested by Minagawa *et al.* (1983). Final identification of presumptive *E. tarda* isolates was done based on detailed morphological and biochemical characterization. All biochemical tests were performed at 25°C unless otherwise specified.

Serological analysis: Bacteria were cultured on nutrient agar (NA, Eiken) at 25°C for 48 h. Slide agglutination tests for isolated *E. tarda* were carried out with 10- and 20-fold diluted rabbit serum which was raised against *E. tarda* NUF251, a reference strain isolated from diseased flounder in Nagasaki Prefecture. The agglutination titers of anti-NUF251 serum were measured in micro-titer plates using both formalin-killed (formalin 0.3%) and heat-killed (100°C, 2.5 h) cells. In addition to NUF251, E22

which was isolated from diseased eel and classified to serotype A (Park *et al.*, 1983) was also used as a reference strain.

Pathogenicity test: Selected 16 isolates of *E. tarda* were tested on their pathogenicity. *E. tarda* NUF251 was used as a positive control. A 2-day-culture of each isolate on NA at 25°C was diluted in sterilized PS to a concentration of 10⁵–10⁶ CFU/ml. Five flounder (average body weight: 13.1 g) were injected intramuscularly with 0.05 ml of each bacterial suspension or of PS as a control group. No food was given during the experiment. The fish were observed everyday for 15 days at about 22°C of water temperature and dead fish were submitted to bacterial isolation with NA and SS agar from kidney.

Survival of *E. tarda* in waters

Two isolates from water and healthy fish were selected to study their survivabilities in sea water, freshwater (from a freshwater pond in the campus of Hiroshima University), and distilled water. Two reference strains (NUF251 and E22) were also used for this study. One hundred ml of each kind of water were autoclaved and kept at 20°C overnight. Two day-culture of 4 strains at 25°C on NA was diluted in PS to make a concentration of 10⁷ CFU/ml, and 0.1 ml of the suspension was inoculated into each of the water media. The inoculated water were kept at 20°C and viable count was made on NA at appropriate intervals. Slide agglutination tests were performed using anti-*E. tarda* NUF251 rabbit serum to confirm that the grown colonies are of *E. tarda*.

Results

Ecological survey

Water temperature and fish body weight at each sampling are shown in Table 1. The growth rate of fish of pond A was much higher than that of pond B.

There were no apparent outbreaks of *E. tarda* infection in the two farms investigated during the study period. Results of *E. tarda* detection by direct plating (DP) and enrichment culture (EC) methods are shown in Table 2 and 3, respectively. EC method gave a much higher detection rate than DP method and filter paper method was highly effective for the detection of *E. tarda* from waters. By EC method (Table 3), *E. tarda* was detected from pond water (filter paper method) and/or sediment samples from June to September, and from July the pathogen was

Table 1. Water temperature and fish body weight at each sampling in two flounder farms

Date (1992)	Water temperature (°C)		Mean fish body weight ¹⁾ (g)	
	Pond A	Pond B	Pond A	Pond B
Apr. 30	19.5	16.3	0.1	0.1
May 27	19.5	17.2	2.1	2.0
Jun. 30	20.2	19.5	14.7	13.6
Jul. 28	23.5	24.2	42.2	34.9
Aug. 31	27.4	27.1	97.9	61.7
Sep. 28	22.4	22.8	140.5	96.6
Oct. 30	21.5	21.2	196.0	137.2
Nov. 26	15.9	15.0	247.3	143.9
Dec. 22	13.5	13.2	273.1	160.7

¹⁾ Average body weight of 10 fish.**Table 2.** Results of *E. tarda* detection by direct plating method from apparently healthy flounder and their culture environment in two flounder farms

Materials for isolation	Detection of <i>E. tarda</i>								
	Apr. 30	May 27	Jun. 30	Jul. 28	Aug. 31	Sep. 28	Oct. 30	Nov. 26	Dec. 22
Pond water	-/- ¹⁾	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
Supply water	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
Sediment	-/-	-/-	-/-	+/-	-/-	-/-	-/-	-/-	+/-
Fish									
whole body	-/-								
intestine	ND ²⁾	-/-	-/-	-/-	+(1) ³⁾ -	-/-	+(1)/(1)	-/-	+(1)-

¹⁾ Results in A pond (A farm)/B pond (B farm); ²⁾ Not done; ³⁾ Number of fish from which *E. tarda* was isolated.**Table 3.** Results of *E. tarda* detection by enrichment culture method from apparently healthy flounder and their culture environment in two flounder farms

Materials for isolation	Detection of <i>E. tarda</i>								
	Apr. 30	May 27	Jun. 30	Jul. 28	Aug. 31	Sep. 28	Oct. 30	Nov. 26	Dec. 22
Pond water ¹⁾	ND ³⁾	ND	-/- ⁴⁾	-/-	-/-	-/-	-/-	-/-	-/-
²⁾	ND	ND	+/-	+/+	+/+	+/-	-/-	+/-	+/-
Supply water 1	ND	ND	-/-	-/-	-/-	-/-	-/-	-/-	-/-
2	ND	ND	-/-	+/-	-/-	+/-	-/-	-/-	-/-
Sediment	-/-	-/-	+/-	+/-	-/-	-/-	+/-	-/-	+/-
Food	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-	-/-
Fish									
whole body	-/-								
intestine	ND	-/-	-/-	-/(1) ⁵⁾	+(1)-	-/-	+(2)-	+(3)-	+(4)-
liver	ND	-/-	-/-	-/-	+(1)-	-/-	+(1)-	-/-	+(1)-
spleen	ND	-/-	-/-	-/-	-/-	-/-	+(1)-	-/-	+(1)-
kidney	ND	-/-	-/-	-/-	-/-	-/-	+(1)-	-/-	+(1)-

¹⁾ Enrichment culture of 1 ml/ water sample; ²⁾ Enrichment culture of filter paper passed with 100 ml/ water sample; ³⁾ Not done; ⁴⁾ Results in A pond (A farm)/B pond (B farm); ⁵⁾ Number of fish from which *E. tarda* was isolated.

Table 4. Characteristics of 28 isolates of *E. tarda* obtained in this study with those of *E. tarda* reported by Ewing *et al.* (1965)

Characteristics	Present isolates	Ewing <i>et al.</i>	Characteristics	Present isolates	Ewing <i>et al.</i>
Gram stain	—	—	Acid and gas from glucose	+	+
Motility	+	+	Acid from:		
Oxidase	—	—	arabinose	—	—
Catalase	+	+	cellobiose	—	—
OF test	F ¹⁾	F	galactose	+	+
Nitrate reduction	+	+	mannitol	—	—
Gelatin liquefaction	—	—	mannose	+	+
Indole	+	+	sucrose	—	—
H ₂ S	+	+	Growth in NaCl:		
MR	+	+	0%	+	ND ³⁾
VP	—	—	0.5%	+	ND
β -galactosidase (ONPG)	—	—	3%	+	ND
Arginine decomposition	—	—	6%	—	ND
Lysine decarboxylation	+	+	Growth at:		
Ornithine decarboxylation	+	+	4°C	—	ND
Citrate utilization	+	—	40°C	+	ND
	(60%) ²⁾				

¹⁾ F: Fermentative; ²⁾ Percentage of strains showing positive result; ³⁾ No data.

also detected from fish except September. Based on EC method, *E. tarda* detection rates from pond water (filter paper method), sediment and fish samples in pond A and B were 86% (6/7), 44% (4/9), 16% (14/90); and 29% (2/7), 0% (0/9), 2% (2/90), respectively. Food samples were found to be free from *E. tarda* contamination in either farm.

Number of viable cells of *E. tarda* detected by DP method was from 8 to 62 CFU/g in fish intestine, and 5 to 420 CFU/g in sediment.

Characteristics of isolates

Morphological and biochemical characteristics: In total 40 isolates were picked up based on the colony appearance on SS agar plates, out of which 28 isolates were identified as *E. tarda* by rapid identification; 8 from pond water, 2 from supply water, 4 from sediment, 11 from intestine, 1 from liver, 1

from spleen and 1 from kidney of flounder. All these 28 isolates were confirmed to be *E. tarda* by detailed characterization. Morphological and biochemical characteristics of the isolates are shown in Table 4. All results coincided with those of *E. tarda* reported by Ewing *et al.* (1965) except citrate utilization.

Serology: All the 28 isolates reacted positively in slide agglutination tests with 10- and 20-fold dilutions of anti-*E. tarda* NUF251 rabbit serum. Distribution of agglutination titers of antiserum against formalin-killed and heat-killed cells of the present 28 isolates and 2 reference strains (NUF251, E22) are shown in Table 5. In most of the isolates, the titer was 160 for heat-killed cells and 320 or 640 for formalin-killed cells.

Pathogenicity: Results of pathogenicity test are shown in Table 6. Except one intestinal isolate (AFI 10-7), all the isolates examined were proved to be highly virulent in flounder (mortality 80 or 100% within 6-8 days). Kidney samples of all dead fish showed growth of pure *E. tarda* colonies.

Table 5. Agglutination titers of anti-*E. tarda* NUF251 rabbit serum against the present 28 isolates and two reference strains of *E. tarda*

Strain	Titer	
	Formalin-killed cell	Heat-killed cell
Present 28 isolates	320-1280	80-320
Reference strains: NUF251	1280	320
E22	320	160

Survivability in waters

Results of survivability tests of the two present isolates (APW 08, AFK 12) (Table 6) and two reference strains (NUF251, E22) are shown in Fig. 1. The 4 strains survived for the entire 35 days of experimental period in freshwater (pond water) and for 14 to 21 days in distilled water, but in sea water,

Table 6. Pathogenicity to Japanese flounder of the present 16 isolates and NUF251 (reference strain) of *E. tarda*

Strain	Source			Dose CFU/fish ($\times 10^4$)	Mortality ¹⁾ (%)
	Pond	Date	Sample		
APW 06	A	Jun.	Pond water	4.0	100
BPS 06	B	Jun.	Sediment	7.0	100
ASW 07	A	Jul.	Supply water	4.0	80
BPW 07	B	Jul.	Pond water	7.0	100
APW 08	A	Aug.	Pond water	5.5	100
AFI 08	A	Aug.	Fish intestine	3.3	100
AFL 08	A	Aug.	Fish liver	11.0	100
APW 09	A	Sep.	Pond water	6.5	80
AFI 10-7	A	Oct.	Fish intestine	5.5	0
AFS 10	A	Oct.	Fish spleen	7.5	100
APS 10	A	Oct.	Sediment	10.0	100
BFI 10	B	Oct.	Fish intestine	4.3	100
AFI 12-5	A	Dec.	Fish intestine	7.5	100
AFK 12	A	Dec.	Fish kidney	4.2	100
APW 12	A	Dec.	Pond water	7.0	100
APS 12	A	Dec.	Sediment	8.0	100
NUF251				1.3	100
Control (Physiological saline)					0

¹⁾ Five fish were injected intramuscularly with each strain at 22°C.

Fig. 1. Survival of the present 2 isolates and reference 2 strains of *E. tarda* in 3 different waters at 20°C.
 —○— Sea water, —●— Pond water, —□— Distilled water

they perished within 3 to 7 days.

Discussion

During the 9 months' sampling from April to December 1992, *E. tarda* was isolated first from the water or sediment samples in June. In July, it was also detected from the supply water and fish. Food was found to be free from *E. tarda* contamination throughout the survey. This finding may indicate that the original source of *E. tarda* was sea water in these two farms. These two farms suffered from edwardsiellosis in the previous year, thus the pathogen may have survived in sea water environments, probably in biotic elements, of coastal area where the used water from the farms are drained.

E. tarda was detected from the intestine most often with highest density among the tested organs. This seems to indicate that the intestine is the primary organ for the bacterial settlement in flounder as also reported by Kanai *et al.* (1988). Kanai *et al.* (1988) detected the pathogen from kidney only when its number in the intestines rose higher than 10^3 CFU/g. But in the present work *E. tarda* was detected from the liver, spleen and kidney even when density of the pathogen in the intestine was lower than 10^3 CFU/g. This will imply that the pathogen may multiply simultaneously in several organs in certain cases of the infection.

The present study confirmed that *E. tarda* existed in environment of flounder even when the disease did not occur. Kanai *et al.* (1988) could not detect *E. tarda* from flounder culture environment except for one sampling made during an outbreak of the disease, though they also used enrichment culture technique. This difference seems to be due to the difference in flounder culture system between our sampling farms and their sampling farms: that is, our study was made in land based ponds and their study was made in net cages in sea water.

There were no great differences in the reaction against the antiserum in heat-killed states, indicating that all the isolates belong to one O-serotype. All the isolates examined except one was virulent to Japanese flounder. Park *et al.* (1983) and Minagawa *et al.* (1983) reported that there were some different serotypes among *E. tarda* strains isolated from healthy eels and their environment. Their isolates included avirulent or low virulent strains. On the contrary, *E. tarda* isolated from flounder and their

environment consisted of homogeneous isolates in serological and pathological features not only in our study but also in the study made by Kanai *et al.* (1988). Also Mamnur Rashid *et al.* (1994) examined 28 strains of *E. tarda* isolated from diseased flounder in different Prefectures of Japan and found that they also belong to one O-serotype which is identical to type A of eel strains. Therefore it is proved that *E. tarda* isolates either from diseased or healthy flounder or their environment belong to one O-serotype (type A of eel strains). This will imply that only some strains, either having pathogenicity or affinity for flounder and belonging to one serotype, could have existed in sea water environment. Therefore, the ecology of *E. tarda* in flounder farms is different from that of *E. tarda* in eel farms.

E. tarda is frequently isolated from cold-blooded animals and their environment, particularly freshwater (Farmer III and McWhorter, 1984). All the strains tested here died within a week in sea water, indicating that they can not live long in sea water itself. Thus, as mentioned above, only pathogenic strains have been able to remain in flounder or other animals in or around culture environment, although various types of the bacterium could have opportunity to enter into marine environment.

Acknowledgements

We thank Mr. Jun Kaneda and Mr. Masatoki Matsuki for facilitating samplings. The first author is grateful to the Monbusho for scholarship support.

References

- Austin, B. and D. A. Austin (1993): Enterobacteriaceae representatives. In "Bacterial fish pathogens: Disease in farmed and wild fish", 2nd ed., Ellis Horwood Ltd., Chichester, pp. 188-226.
- Ewing, W. H., A. C. McWhorter, M. R. Escobar and A. H. Lubin (1965): *Edwardsiella*, a new genus of Enterobacteriaceae based on a new species, *E. tarda*. *Int. Bull. Bacteriol. Nomencl. Taxon.*, **15**, 33-38.
- Farmer III, J. J. and A. C. McWhorter (1984): Genus X. *Edwardsiella* Ewing and McWhorter 1965. In "Bergey's manual of systematic bacteriology, Vol. 1" (ed. by N. R. Krieg). Williams and Wilkins, Baltimore, pp. 486-491.
- Hoshina, T. (1962): Studies on red disease of eels. *J. Tokyo Univ. Fish.*, Spec. Rep., **6**, 1-104. (In Japanese)
- Kanai, K., S. Tawaki and Y. Uchida (1988): An ecological study of *Edwardsiella tarda* in flounder farm. *Fish*

- Pathol.*, **23**, 41–47.
- Kodama, H., T. Murai, Y. Nakanishi, F. Yamamoto, T. Mikami and H. Izawa (1987): Bacterial infection which produces high mortality in cultured Japanese flounder (*Paralichthys olivaceus*) in Hokkaido. *Jpn. J. Vet. Res.*, **35**, 227–234.
- Mamnur Rashid, M., T. Mekuchi, T. Nakai and K. Muroga (1994): A serological study on *Edwardsiella tarda* strains isolated from diseased Japanese flounder (*Paralichthys olivaceus*). *Fish Pathol.*, **29**, 277.
- Minagawa, T., T. Nakai and K. Muroga (1983): *Edwardsiella tarda* in eel culture environment. *Fish Pathol.*, **17**, 243–250.
- Nakatsugawa, T. (1983): *Edwardsiella tarda* isolated from cultured young flounder. *Fish Pathol.*, **18**, 99–101.
- Park, S.-I., H. Wakabayashi and Y. Watanabe (1983): Serotype and virulence of *Edwardsiella tarda* isolated from eel and their environment. *Fish Pathol.*, **18**, 85–89.
- Wakabayashi, H. and S. Egusa (1973a): Seasonal changes of bacterial infections among pond-cultured eels (*Anguilla japonica*). *Fish Pathol.*, **8**, 91–97.
- Wakabayashi, H. and S. Egusa (1973b): *Edwardsiella tarda* (*Paracolobactrum anguillimortiferum*) associated with pond-cultured eel disease. *Nippon Suisan Gakkaishi*, **39**, 931–936.
- Wyatt, L. E., R. Nickelson II and C. Vanderzant (1979): *Edwardsiella tarda* in freshwater catfish and their environment. *Appl. Environ. Microbiol.*, **38**, 710–714.
- Yasunaga, N., S. Ogawa and K. Hatai (1982): Characteristics of the fish pathogen *Edwardsiella* isolated from several species of cultured marine fishes. *Bull. Nagasaki Prefect. Ins. Fish.*, **8**, 57–65. (In Japanese)