Original Article

A Family Predisposition to Adult T-Cell Leukemia

Kouichiro Nomura¹⁾, Atae Utsunomiya²⁾, Hironori Furushou²⁾, Mitsutoshi Tara³⁾, Masashi Hazeki⁴⁾, Masayoshi Tokunaga⁵⁾, Kimiharu Uozumi⁶⁾, Shuichi Hanada⁶⁾, Shinji Yashiki⁷⁾, Kazuo Tajima⁸⁾ and Shunro Sonoda⁷⁾

We report here the rare case of a family predisposed to adult T-cell leukemia (ATL). Six of seven siblings developed ATL with ages of onset of 77, 48, 60, 64, 72, and 62 years old. Although virological tests for human T-lymphotropic virus type 1 were unavailable for two of the six patients, all were diagnosed with ATL based on their clinical, hematological, and histopathological features. Two of the six patients were tested for HLA haplotypes using fresh blood samples, and both were carriers of the HLA-A*26 allele known in the southern Japanese population to be susceptible to ATL. This series of genetic traits may help explain the familial predisposition to ATL. [*J Clin Exp Hematopathol* 46(2) : 67-71, 2006]

Keywords: adult T-cell leukemia, familial ATL, HTLV-1, HLA allele

INTRODUCTION

Adult T-cell leukemia (ATL) is a T-cell neoplasia with specific clinical and epidemiological features, such as circulating leukemia cells with peculiar nuclear polymorphisms (convoluted or lobulated nuclei), lymphadenopathy, hepatosplenomegaly, skin manifestation, hypercalcemia, and a specific geographic distribution¹⁻³. Human T-lymphotropic virus type 1 (HTLV-1) is the causative of ATL⁴⁻⁶ and is transmitted within families from mothers to infants via breast-feeding and between spouses via sexual intercourse⁷. The infection rate of children of HTLV-1 carrier mothers is estimated at 10-30%⁸, and the HTLV-1 seropositivity rate is as high as 65% in ATL related siblings^{9,10}. Although a high incidence of familial clusters of ATL would therefore be expected, few family pedigrees of ATL have been discovered in which 2 of 4-7 siblings from a single family developed ATL¹¹⁻¹³. We report here a rare case of highly accumulated familial ATL that exhibits the HLA alleles and haplotypes conferring susceptibility to ATL.

DIAGNOSTIC CRITERIA FOR ATL

ATL was diagnosed according to the criteria of the Japanese Lymphoma Study Group organized by Shimoyama *et al*¹⁴. These criteria were the following : (i) histologically and/or cytologically proven lymphoid malignancy with T-cell surface antigens ; (ii) abnormal T-lymphocytes present in the peripheral blood, except lymphoma type. Abnormal T-lymphocytes include both typical ATL called “flower cells,” and the small and mature T-lymphocytes with incised nuclei that are characteristic of chronic and smoldering type ATL ; (iii) antibodies to HTLV-1 present in the sera at the time of diagnosis. ATL clinical subtypes were also classified according to the criteria of the Japanese Lymphoma Study Group¹⁴.

CASE REPORT

Case 1 (Sibling 1, the proband) was a 77-year-old woman who developed swelling of the peripheral lymph nodes (LNs) and skin lesions in November 1989. Her white blood cell count (WBC) was $100 \times 10^9/L$ and 95% of these were abnormal lymphocytes ($CD3^+CD4^+CD8^-$) with convoluted or lobulated nuclei (Fig. 1A). Examination of her LNs revealed diffuse, pleomorphic type non-Hodgkin's lymphoma (NHL). The lymphoma cells were positive for CD2, CD3, CD4, CD25, and CD30, but not CD8. A serological test for anti-HTLV-1 antibodies was positive and the antibody titer was useful at greater than a 40-fold dilution in an immunofluorescence (IF) assay. Based on these results, she was diagnosed with ATL and treated with cyclophosphamide, vindesine, and prednisolone. She went into partial remission but

Received : Oct 21, 2005

Revised : Feb 20, 2006

Accepted : Mar 23, 2006

¹⁾Internal Medicine, Kagoshima Prefectural Hokusatsu Hospital, ²⁾Department of Hematology, Imamura Bun-in Hospital, ³⁾Kagoshima Municipal Hospital, ⁴⁾Hazeki Hospital, ⁵⁾Kodama Hospital, ⁶⁾Second Department of Internal Medicine and ⁷⁾Department of Virology, Faculty of Medicine, Kagoshima University, Kagoshima, and ⁸⁾Division of Epidemiology and Prevention, Aichi Cancer Center Research Institute, Nagoya, Japan

Address correspondence and reprint request to Atae Utsunomiya, Department of Hematology, Imamura Bun-in Hospital, Kamoikeshinmachi 11-23, Kagoshima 890-0064, Japan.

Fig. 1. Abnormal circulating lymphocytes with convoluted or lobulated nuclei. A, Sibling 1 (Case 1); B, Sibling 2 (Case 5); C, Sibling 4 (Case 3); D, Sibling 5 (Case 4); E, Sibling 6 (Case 6); F, Sibling 7 (Case 2). All images are of May-Giemsa stains, x1000.

later died of tumor progression. Her survival period was 17 months after beginning chemotherapy. According to her family history, five of her six siblings had hematological malignancies diagnosed as either lymphoma or leukemia. Hematological malignancies containing ATL, however, were not recognized in the other relatives.

Case 2 (Sibling 7) was a 48-year-old man admitted to our hospital in 1978 due to coughing and dyspnea. He had lymphadenopathy and a WBC count of $49.6 \times 10^9/L$, 90% of which were abnormal lymphocytes with convoluted nuclei (Fig. 1F). As the concept of ATL was unknown in 1978, he was diagnosed with leukemic reticulosarcoma. The typical morphological features of ATL cells found during our review of his peripheral blood specimen caused us to strongly suspect ATL. However, he was treated with combination chemotherapy and died three months after diagnosis. His cause of death was listed as unknown.

Case 3 (Sibling 4) was a 60-year-old woman with swelling of her peripheral LNs and a large number of abnormal lymphocytes with convoluted nuclei in her blood in 1982 (Fig. 1C). Examination of her LNs revealed NHL of diffuse, medium cell type. The surface phenotype of her lymphoma cells was found to be that of T cells by immunohistochemical staining with monoclonal antibodies for UCHL-1, MT-1, and DF-T1. A test for anti-HTLV-1 antibodies was not performed, and her survival period after chemotherapy was three months. Her cause of death was also unknown.

Case 4 (Sibling 5) was a 64-year-old woman with swelling of her cervical LNs in February 1988. She developed multiple skin tumors throughout her clinical course. Her WBC count was $7.7 \times 10^9/L$, of which 33% were abnormal lymphocytes with convoluted nuclei (Fig. 1D). Examination of the skin tumors revealed NHL of diffuse, pleomorphic

type. The lymphoma cells stained positively for CD2, CD4, CD25, and CD30. A serological test for HTLV-1 antibodies was positive. She died of septicemia, and her survival period after chemotherapy was nine months.

Case 5 (Sibling 2) was a 72-year-old woman who noticed swelling of her cervical and inguinal LNs in July 1988. Her WBC count was $3.2 \times 10^9/L$, 10% of which were abnormal lymphocytes with indented nuclei (Fig. 1B). Examination of her LNs showed diffuse, pleomorphic type NHL. The surface phenotype of the lymphoma cells was that of T cells, according to immunohistochemical staining with monoclonal antibodies against UCHL-1, MT-1, and DF-T1. A serological test for HTLV-1 antibodies was positive, and the antibody titer was useful in an IF test at greater than an 80-fold dilution. She was diagnosed with ATL and treated with combination chemotherapy, but she died from tumor progression ten months after beginning chemotherapy.

Case 6 (Sibling 6) was a 62-year-old man with abdominal lymphadenopathy, ascites, and hypercalcemia in 1989. His WBC count was $68.7 \times 10^9/L$ and 84% of these were abnormal lymphocytes with convoluted nuclei (Fig. 1E). Examination of his LNs showed NHL of diffuse, large cell type. The lymphoma cells had a T-cell phenotype, and he tested positively for anti-HTLV-1 antibodies in his serum. He was diagnosed with ATL and survived only three months after beginning chemotherapy. The cause of his death was unknown.

Sibling 3 had died during World War II at the age of 22.

The clinical signs and symptoms and the hematological and pathological findings of the six patients are summarized in Table 1.

Hematological and pathological examination of biopsy specimens

Abnormal circulating lymphocytes from the six cases are shown in Figure 1 (A to F). Convoluted and/or lobulated nuclei were present in all six cases.

The pathological findings of five cases showed diffuse proliferation of lymphoma cells. The surface phenotype of these lymphoma cells was found by immunohistochemical staining to be that of T cells (Table 1).

HLA allele types and haplotypes

The HLA alleles and haplotypes were examined in two (Siblings 1 and 5) of the seven siblings and in three blood relatives (one son and two grandchildren of Sibling 1). The HLA-A*, HLA-C*, HLA-B*, HLA-DRB1*, and HLA-DQB1* alleles were determined by the PCR-SSO method as previously described^{15,16}. The patients' HLA haplotypes were inferred based on known patterns of linkage disequilibrium at five loci^{17,18}. According to pattern analysis of the HLA alleles

of the ATL family members, specific allele types relating to ATL predisposition (HLA-A*26, B*4002, and B*4006)¹⁶ were found in the ATL-affected patients (Fig. 2).

DISCUSSION

We report here a very rare familial predisposition to ATL. Although HTLV-1 tests for Siblings 4 and 7 and surface phenotyping for leukemia cells in Sibling 7 were not performed, we were able to diagnose ATL in all six patients, including Siblings 4 and 7, based on their specific clinical characteristics and blood cytologies. HTLV-1 is the causative

agent of ATL⁴⁻⁶, and vertical transmission from mother to child via HTLV-1-infected lymphocytes in breast milk is a major route of infection¹⁹⁻²¹. The overall infection rate of children by HTLV-1 carrier mothers in Southern Japan is estimated at 10-30%⁸. Furthermore, the HTLV-1 seropositivity rate of family members of ATL sufferers is about 50%⁹, and the rate for their siblings is as high as 65%^{9,10}. This evidence suggests that the mother of siblings reported here was probably infected by HTLV-1 and that the virus transmitted vertically to the siblings. The high rate of HTLV-1 infection in this family might have related to a high viral load in the breast milk, or an HLA allele-based susceptibility to

Fig. 2. Family pedigree showing HLA haplotypes predisposed to ATL. Siblings 1, 2, 4, 5, 6, and 7 died of ATL, while Sibling 3 died in World War II. IF, immunofluorescence ; PA, gelatin particle agglutination.

Table 1. Hematological and pathological findings of familial ATL cases

Order of siblings	WBC ($\times 10^9/L$)	Abnormal lymphocyte (%)	Age at onset	Sex	Subtype of ATL	Pathological diagnosis (LN)	Surface phenotype	anti-HTLV-1
1 (case 1)	100	95	77	F	Acute	Diffuse, pleomorphic	CD3(+) CD4(+), CD25(+)(PB, LN)	> 40 $\times$ (IF)
2 (case 5)	3.3	10	72	F	Acute	Diffuse, pleomorphic	UHL-1(+), MT-1(+), DF-T1(+)(LN)	> 80 $\times$ (IF)
4 (case 3)	LOI	LOI	60	F	Acute	Diffuse, pleomorphic	UHL-1(+), MT-1(+), DF-T1(+)(LN)	NT
5 (case 4)	7.7	33	64	F	Acute	Diffuse, pleomorphic	CD4(+), CD25(+)(skin)	512 $\times$ (PA)
6 (case 6)	68.7	84	62	M	Acute	Diffuse, pleomorphic	UHL-1(+)(LN)	> 128 $\times$ (PA)
7 (case 2)	49.6	90	48	M	Acute	NT	NT	NT

LOI: lack of information, NT: not tested, PB: peripheral blood, LN: lymph node, IF: immunofluorescence method, PA: gelatin particle agglutination method

HTLV-1 in all six siblings. We previously reported that the annual incidence of ATL in Kyusyu was 0.60, based on studies of 1,000 HTLV-1 carriers over 20 years old⁸. Hisada *et al.* reported that HTLV-1 carriers with a higher anti-HTLV-1 titer and a lower anti-Tax reactivity might be at an increased risk for ATL²². We could not, of course, examine the anti-HTLV-1 titers or anti-Tax reactivities in our patients before the onset of ATL. In regard to the genetic risk for ATL, we also reported a relationship between specific HLA alleles and a predisposition to ATL¹⁶. We demonstrated that lower reactivity of anti-Tax cytotoxic T lymphocytes is associated with the HLA-A*26, B*4002, and B*4006 alleles, and higher reactivity is associated with the HLA-A*24 alleles¹⁶. Although HLA haplotypes were examined in only two cases of this family (Siblings 1 and 5), they both had HLA-A*26 alleles. The haplotypes of Siblings 1 and 5 are exclusive, as Sibling 1 has one A*26+haplotype and Sibling 5 has two A*26+haplotypes. From this distribution of HLA haplotypes, one of their parents should be homozygous for A*26+/A*26+, as 50% of the children had homozygous A*26+haplotypes. The A*26+haplotype carriers are low responders against HTLV-1-Tax antigen and are susceptible to HTLV-1 infection and ATL manifestation. This rare family, with six of seven siblings diagnosed with ATL, is predisposed by the ATL-susceptible HLA-A*26 gene, and consequently the risk of ATL was extremely high.

On the other hand, the HTLV-1 viral factor might relate to the incidence of ATL in the patients reported here. Furukawa *et al.* reported that the phylogenetic subgroup of HTLV-1 in the *tax* gene was associated with the occurrence of HTLV-1-associated myelopathy/tropical spastic paraparesis, but not with the incidence of ATL²³. We could not examine the *tax* sequence in our ATL patients, so there might be a relationship between HTLV-1 subgroup, *tax* mutation, deletion, and the incidence of ATL.

In conclusion, immunogenetic features—including specific HLA alleles—may be related to this family's high predisposition to ATL. To our knowledge, this is the most significant ATL cluster in a single family yet reported - a cluster in which six of seven siblings developed ATL.

REFERENCES

- 1 Takatsuki K, Uchiyama T, Sagawa K, Yodoi J: Adult T cell leukemia in Japan. In Seno S, Takaku F, Irino S (eds). Topics in Hematology. Amsterdam-Oxford: Excerpta Medica, pp73-77, 1977
- 2 Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H: Adult T-cell leukemia: Clinical and hematological features of 16 cases. Blood 50: 481-492, 1977
- 3 Matsumoto M, Nomura K, Matsumoto T, Nishioka K, Hanada S, Furushou H, Kikuchi H, Kato Y, Utsunomiya A, Uematsu T, Hashimoto S, Yunoki K: Adult T-cell leukemia-lymphoma in Kagoshima district, southwestern Japan: Clinical and hematological characteristics. Jpn J Clin Oncol 9 (suppl.): 325-336, 1979
- 4 Poiesz BJ, Ruscetti FW, Gazdar AF, Bunn PA, Minna JD, Gallo RC: Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. Proc Natl Acad Sci USA 77: 7415-7419, 1980
- 5 Hinuma Y, Nagata K, Hanaoka M, Nakai M, Matsumoto T, Kinoshita K, Shirakawa S, Miyoshi I: Adult T-cell leukemia: Antigen in an ATL cell line and detection of antibodies to the antigen in human sera. Proc Natl Acad Sci USA 78: 6476-6480, 1981
- 6 Yoshida M, Miyoshi I, Hinuma Y: Isolation and characterization of retrovirus from cell lines of human adult T-cell leukemia and its implication in the disease. Proc Natl Acad Sci USA 79: 2031-2035, 1982
- 7 Kajiyama W, Kashiwagi S, Ikematsu H, Hayashi J, Nomura H, Okochi K: Intrafamilial transmission of adult T cell leukemia virus. J Infect Dis 154: 851-857, 1986
- 8 Tajima K, Hinuma Y: Epidemiology of HTLV-I/II in Japan and the world. Gann Monogr Cancer Res 39: 129-149, 1992
- 9 Nomura K: A study anti-ATLA (antibodies to adult T-cell leukemia virus associated antigens) in healthy subjects. A sero-epidemiological study on residents of Kagoshima prefecture and a family study of patients with adult T-cell leukemia-lymphoma and other diseases. Igaku Kenkyu 54: 750-781, 1984 (*in Japanese*)
- 10 Momita S, Ikeda S, Amagasaki T, Soda H, Yamada Y, Kamihira S, Tomonaga M, Kinoshita K, Ichimaru M: Survey of anti-human T-cell leukemia virus type I antibody in family members of patients with adult T-cell leukemia. Jpn J Cancer Res 81: 884-889, 1990
- 11 Miyamoto Y, Yamaguchi K, Nishimura H, Takatsuki K, Motoori T, Morimatsu M, Yasaka T, Ohya I, Koga T: Familial adult T-cell leukemia. Cancer 55: 181-185, 1985
- 12 Ratner L, Heyden NV, Paine E, Frei-Lahr D, Brown R, Petruska P, Reddy S, Lairmore MD: Familial adult T-cell leukemia/lymphoma. Am J Hematol 34: 215-222, 1990
- 13 Wilks RJ, LaGrenade L, Hanchard B, Campbell M, Murphy J, Cranston B, Blattner WA, Manns A: Sibling adult T-cell leukemia/lymphoma and clustering of human T-cell lymphotropic virus type I infection in a Jamaican family. Cancer 72: 2700-2704, 1993
- 14 Shimoyama M: Diagnostic criteria and classification of clinical subtypes of adult T-cell leukaemia-lymphoma. A report from the Lymphoma Study Group (1984-87). Br J Haematol 79: 428-437, 1991
- 15 Lou H, Li HC, Kuwayama M, Yashiki S, Fujiyoshi T, Suehara M, Osame M, Yamashita M, Hayami M, Gurtsevich V, Ballas M, Imanishi T, Sonoda S: HLA class I and class II of the Nivkhi, an indigenous population carrying HTLV-I in Sakhalin, far eastern Russia. Tissue Antigens 52: 444-451, 1998
- 16 Yashiki S, Fujiyoshi T, Arima N, Osame M, Yoshinaga M, Nagata Y, Tara M, Nomura K, Utsunomiya A, Hanada S, Tajima K, Sonoda S: HLA-A*26, B*4002, B*4006 and B*4801 alleles pre-

- dispose to adult T-cell leukemia : The limited recognition of HTLV-I Tax peptide anchor motifs and epitopes to generate anti-HTLV-I Tax CD8+CTL. *AIDS Res Hum Retrov* 17 : 1047-1061, 2001
- 17 Akaza T, Imanishi T, Fujiwara K, Tokunaga K, Yashiki S, Fujiyoshi T, Sonoda S, Juji T : HLA alleles and haplotypes of Japanese population. *MHC IRS 1 (Suppl.1)* : S219-S226, 1994
- 18 Imanishi T, Akaza T, Kimura A, Tokunaga K, Gojobori T : Allele and haplotype frequencies for HLA and complement loci in various ethnic group. *In* Tsuji K, Aizawa M, and Sasazuki T, (eds) : *HLA 1991 Vol. I*. Oxford University Press, Tokyo, Japan, 1992, pp1065-1220
- 19 Kinoshita K, Hino S, Amagasaki T, Ikeda S, Yamada Y, Suzuyama J, Momita S, Toriya K, Kamihira S, Ichimaru M : Demonstration of adult T-cell leukemia virus antigen in milk from three sero-positive mothers. *Jpn J Cancer Res* 5 : 103-105, 1984
- 20 Nakano S, Ando Y, Ichijo M, Moriyama L, Saito S, Sugimura K, Hinuma Y : Search for possible routes of vertical and horizontal transmission of adult T-cell leukemia virus. *Jpn J Cancer Res* 75 : 1044-1045, 1984
- 21 Hino S, Yamaguchi K, Katamine S, Sugiyama H, Amagasaki T, Kinoshita K, Yoshida Y, Doi H, Tsuji Y, Miyamoto T : Mother-to-child transmission of human T-cell leukemia virus type-I. *Jpn J Cancer Res* 76 : 474-480, 1985
- 22 Hisada M, Okayama A, Shioiri S, Spiegelman DL, Stuver SO, Mueller NE : Risk factors for adult T-cell leukemia among carriers of human T-lymphotropic virus type I. *Blood* 92 : 3557-3561, 1998
- 23 Furukawa Y, Yamashita M, Usuku K, Izumo S, Nakagawa M, Osame M : Phylogenetic subgroups of human T cell lymphotropic virus (HTLV) type I in the *tax* gene and their association with different risks for HTLV-I-associated myelopathy/tropical spastic paraparesis. *J Infect Dis* 182 : 1343-1349, 2000