Contact YAG Laser Palatopharyngoplasty

Hwoe Young Ann

Department of Otolaryngology, College of Medicine Kyung Hee University, Seoul, Korea

Snoring, the lay term for obstructive breathing during sleep, is one of the most prevalent of obnoxious human habits. In a 30- to 35-year-old population, 20% of men and 5% of women will snore; by age 60, 60% of men, and 40% of women will snore habitually.

Most of the heavy snorer with or without sleep apnea have symptoms of 1) loud, irregular snoring, and snorts, gasps and other unusual breathing sounds during sleep, 3) pauses in breathing during sleep, 4) excessive day time sleepiness and 5) fatigue, 6) obesity, 7) changes in alertness, memory, personality, or behavior, 8) impotence, 9) morning headache, and 10) bed wetting.

Common anatomical factors contributing to snoring are 1) incompetent tone of palatal, pharyngeal, and glossal muscles, 2) space occupying masses 3) excessive length of the soft palate and uvula, and 4) obstructive nasal breathing.

Various methods have been utilized in order to improve snoring and/or obstructive sleep apnea syndrome; weight reduction, change of sleep position, stop or reduce alcohol or sedatives, some medications, nasal continuous positive airway pressure (CPAP), and uvulopalatopharyngoplasty (UPPP). The surgical procedure, UPPP has been known the most effective method, with very satisfactory result; elimination or improvement of snoring was reported by Pelausa in 75.9% of patients and by Chouard in 86.3% of patients.

But the conventional UPPP has disadvantages of general anesthesia, several days of hospital stay, severe post-operative pain and high cost for the patients. So, couple of years ago, Dr. Kamami Yves-Victor of Foch Hospital in Paris introduced the CO₂ laser resection of the palatopharynx (LRPP).

LRPP of Dr. Kamami was a new technique, safe and effective, ambulatory by local anesthesia, performed for the treatment of snoring and in some cases of obstructive sleep apnea syndrome (OSAS). The LRPP was based on the progressive enlargement of the air space in the oropharynx, to eliminate or reduce obstructions that occur during sleep, by successive "vaporizations" of the vibrating soft palate, wide posterior tonsil pillars and redundant posterior pharyngeal mucosa. According to Dr. Kamami, he had the 85% of success rate, and Dr. Jack A.

Coleman, Vanderbilt University, Who learned this technique from Kamami and introduced it to U.S., produced 85-95% of success rate.

But this technique LRPP with CO₂ laser needs 3 to 7 operations, usually 3 to 5 for the women, 4 to 5 for the men, "non-OSAS", interval with a fortnight to 3 or 4 weeks between each. That means 4 to 8 months are necessary until the procedures are completed, although the procedures are simple, reliable, non-invasive and painless procedure under local anesthesia without hospitalization and also operator must prepare the new special type of handpiece to protect the posterior pharyngeal wall from CO₂ laser beam.

So I tried more simple, one stage operation with same effectiveness, using the contact YAG laser. Under the local anesthesia with 10% lidocaine spray into the oropharynx first, then injection of 2 % lidocaine with epinephrine, I began to resect the part of posterior pillar, soft palate and uvula, using 7-9 watt of contact YAG laser. After excision of redundant soft tissue of the soft palate and pharynx. I sutured the anterior and posterior resected margins of the mucosa with 4-0 vicryl. After the procedure I gave analgesics and antibiotics to the patient.

Compare to CO₂ LRPP, my contact YAG LPPP was safer, more simple, more convenient, easier to handle, and cheaper economical procedure.

I have done 10 to 15 procedures each since last year. Usually, about 8.3% of patients were satisfied with this procedure. That means elimination or improvement of snoring.

Side effects were post-operative pain and nasal regurgitation. Post-operative swallowing pain was followed 5 to 7 days, but was well controlled by analgesics medication and 2% lidocaine viscous gargling solution. Mild nasal regurgitation was complained in first week in some patients, but improved spontaneously in most patients. There were no patients who complain the change of tone of voice.