N₂O Emission from Degraded Soybean Nodules Depends on Denitrification by *Bradyrhizobium japonicum* and Other Microbes in the Rhizosphere

SHOKO INABA¹, FUMIO IKENISHI¹, MANABU ITAKURA¹, MASAKAZU KIKUCHI¹, SHIMA EDA¹, NAOHIKO CHIBA², CHIE KATSUYAMA², YUICHI SUWA², HISAYUKI MITSUI¹, and KIWAMU MINAMISAWA^{1*}

¹Graduate School of Life Sciences, Tohoku University, 2–1–1 Katahira, Aoba-ku, Sendai 980–8577, Japan; and

²Department of Biological Sciences, Faculty of Science and Engineering, Chuo University, 1–13–27 Kasuga, Bunkyo-ku, Tokyo 112–8551, Japan

(Received May 6, 2012—Accepted May 30, 2012—Published online October 5, 2012)

A model system developed to produce N₂O emissions from degrading soybean nodules in the laboratory was used to clarify the mechanism of N₂O emission from soybean fields. Soybean plants inoculated with *nosZ*-defective strains of *Bradyrhizobium japonicum* USDA110 ($\Delta nosZ$, lacking N₂O reductase) were grown in aseptic jars. After 30 days, shoot decapitation (D, to promote nodule degradation), soil addition (S, to supply soil microbes), or both (DS) were applied. N₂O was emitted only with DS treatment. Thus, both soil microbes and nodule degradation are required for the emission of N₂O from the soybean rhizosphere. The N₂O flux peaked 15 days after DS treatment. Nitrate addition markedly enhanced N₂O emission. A ¹⁵N tracer experiment indicated that N₂O was derived from N fixed in the nodules. To evaluate the contribution of bradyrhizobia, N₂O emission was compared between a *nirK* mutant ($\Delta nirK\Delta nosZ$, lacking nitrite reductase) and $\Delta nosZ$. The N₂O flux from the $\Delta nirK\Delta nosZ$ rhizosphere was significantly lower than that from $\Delta nosZ$, but was still 40% to 60% of that of $\Delta nosZ$, suggesting that N₂O emission is due to both *B. japonicum* and other soil microorganisms. Only *nosZ*-competent *B. japonicum* (*nosZ*⁺ strain) could take up N₂O. Therefore, during nodule degradation, both *B. japonicum* and other soil microorganisms release N₂O from nodule N via their denitrification processes (N₂O source), whereas *nosZ*-competent *B. japonicum* exclusively takes up N₂O (N₂O sink). Net N₂O flux from soybean rhizosphere is likely determined by the balance of N₂O source and sink.

Key words: Nitrous oxide, *Bradyrhizobium japonicum*, Denitrification, *nosZ* gene, Soybean rhizosphere

Nitrous oxide (N₂O) is a key atmospheric greenhouse gas that contributes to global warming and the destruction of stratospheric ozone (14, 46, 47). Agricultural land is a major source of N₂O through the microbial transformation of nitrogen in the soil (13, 24, 58), and contributes significantly to the net increase in atmospheric N₂O (46). Legume crops emit more N₂O than non-legumes (10, 15, 32).

Yang and Cai (55) reported that the emission of N₂O from a soybean field greatly increased in the late growth period, suggesting that senescence and the decomposition of roots and nodules contributed to emissions. Ciampitti *et al.* (7) also reported marked N₂O emissions from a soybean field in the late growth period regardless of N fertilization. N₂O emission from a field with nodulating soybeans was several times higher than that from a field with non-nodulating soybeans (27). N₂O was emitted directly from degraded nodules of field-grown soybeans in the late growth period (20). Thus, soybean nodules emit N₂O under field conditions, although the mechanism remains unresolved.

Microorganisms might be involved, as N₂O can be generated by several microbial processes (4, 13). Using microbial community analysis, Inaba *et al.* (20) nominated potential N₂O producers that increased in abundance in degraded nodules. Among them, *Bradyrhizobium japonicum* was one of the dominant microbes as endosymbionts of soybean nodules and rhizosphere soil bacteria (9, 29, 30, 33,

35, 39). It reduces nitrogen oxides during denitrification as

where each step is catalyzed by specific reductases. These reductases are encoded, respectively, by *napA* (encoding periplasmic nitrate reductase), *nirK* (Cu-containing nitrite reductase), *norCB* (nitric oxide reductase), and *nosZ* (nitrous oxide reductase) (5). The aim of this study was to clarify the involvement of *B. japonicum* in the emission of N₂O from the soybean rhizosphere. The N₂O flux from denitrification mutants of *B. japonicum* was compared in the laboratory.

Materials and Methods

Bacterial strains, plasmids, and media

The bacterial strains and plasmids are listed in Table 1. *Bradyrhizobium japonicum* cells were grown at 30°C in HM salt medium (8) supplemented with 0.1% arabinose and 0.025% (w/v) yeast extract (Difco, Detroit, MI, USA). *Escherichia coli* cells used in transformation were grown at 37°C in Luria–Bertani medium (40). Antibiotics were added to the media at the following concentrations: for *B. japonicum*, 100 µg tetracycline (Tc) mL^{−1}, 100 µg spectinomycin (Sp) mL^{−1}, 100 µg streptomycin (Sm) mL^{−1}, 100 µg kanamycin (Km) mL^{−1}, and 100 µg polymyxin B mL^{−1}; for *E. coli*, 50 µg Tc mL^{−1}, 50 µg Sp mL^{−1}, 50 µg Sm mL^{−1}, 50 µg Km mL^{−1}, and 50 µg ampicillin mL^{−1}.

Construction of *B. japonicum* mutants

Isolation of plasmids, DNA ligation, and transformation of *E. coli* were performed as described previously (40). DNA was prepared as described previously (43). A 5.6-kb *Bam*HI/*Eco*RI fragment

* Corresponding author. E-mail: kiwamu@ige.tohoku.ac.jp;
Tel: +81–22–217–5684; Fax: +81–22–217–5684.

Table 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Relevant characteristics ^a	Source or reference
Strains		
<i>Bradyrhizobium japonicum</i>		
USDA110	Wild type, <i>nosZ</i> ⁺	25
USDA110Δ <i>nosZ</i>	USDA110 derivative, <i>nosZ</i> ::del/ins Tc cassette; Tc ^r	18
USDA110Δ <i>napA</i> Δ <i>nosZ</i>	USDA110 derivative, <i>napA</i> ::Ω cassette, <i>nosZ</i> ::del/ins Tc cassette; Sp ^r , Sm ^r , Tc ^r	18
USDA110Δ <i>nirK</i>	USDA110 derivative, <i>nirK</i> ::Ω cassette; Sp ^r , Sm ^r	This study
USDA110Δ <i>nirK</i> Δ <i>nosZ</i>	USDA110 derivative, <i>nirK</i> ::Ω cassette, <i>nosZ</i> ::del/ins Tc cassette; Sp ^r , Sm ^r , Tc ^r	This study
T9	Field isolate in Tokachi, Hokkaido, Japan, <i>nosZ</i> ⁻	42
<i>Escherichia coli</i>		
DH5a	<i>recA</i> ; cloning strain	Toyobo
Plasmids		
brp01958	pUC18 carrying <i>nirK</i>	25
pHP45Ω	Plasmid carrying 2.1-kb Ω cassette; Sp ^r , Sm ^r , Ap ^r	37
pK18mob	Cloning vector; pMB1 <i>ori</i> oriT; Km ^r	44
pK18mob- <i>nirK</i>	pK18mob carrying 5.6-kb <i>nirK</i> fragment; Km ^r	This study
pK18mob- <i>nirK</i> ::Ω	pK18mob carrying <i>nirK</i> ::Ω cassette; Km ^r , Sp ^r , Sm ^r	This study
pRK2013	ColE replicon carrying RK2 transfer genes; Km ^r	12

^a Ap^r, ampicillin resistant; Tc^r, tetracycline resistant; Km^r, kanamycin resistant; Sp^r, streptomycin resistant; Sm^r, spectinomycin resistant.

containing *nirK* was isolated from brp01958, a clone from the pUC18 library of the sequences of *B. japonicum* USDA110, and inserted into the *Bam*HI/*Eco*RI site of pK18mob (Fig. 1). The Ω-cassette was isolated from pHP45Ω at the *Sma*I site (37) and inserted into pK18mob-*nirK* at the *Psh*AI site, yielding pK18mob-*nirK*::Ω (Fig. 1). pK18mob-*nirK*::Ω was introduced into *B. japonicum* USDA110 by triparental mating using pRK2013 as a helper plasmid (44). A USDA110Δ*nirK*Δ*nosZ* double mutant was constructed by the introduction of pK18mob-*nirK*::Ω into USDA110Δ*nosZ*::Tc (18) (Table 1). Double-crossover events of *nirK* mutation were verified by PCR.

Preparation of soil suspension

Soil was collected from an experimental field at Tohoku University (Kashimadai, Miyagi, Japan). This gray lowland soil had pH[H₂O] 5.6, pH[KCl] 4.2, total C 1.37%, total N 0.132%, and Truog P 48 mg P₂O₅ kg⁻¹. Fresh soil (10 g) was extracted twice with 30 mL distilled water to remove nitrate and nitrite. The suspension was shaken for 10 min in centrifuge tubes and then centrifuged at 5,555×g for 15 min (Himac CR20E; Hitachi, Tokyo, Japan). The pellet was resuspended in 30 mL distilled water.

Inoculation and plant cultivation

Surface-sterilized soybean seeds (*Glycine max* cv. Enrei) were

germinated in sterile vermiculite for 2 days at 25°C. The seedling was then transplanted into a Leonard jar pot (one plant per pot) (28, 53, 56), which contained sterile vermiculite and nitrogen-free nutrient solution (31, 34) (Fig. S1). The seedlings were then inoculated with *B. japonicum* cells at 1×10⁷ cells per seedling. Plants were grown in a phytotron (Koito Industries, Tokyo, Japan) providing 270 μmol photons m⁻² s⁻¹ of photosynthetically active radiation (PAR, 400–700 nm) for 30 days at 25/20°C with a 16-h light/8-h dark cycle. A nitrogen-free sterilized nutrient solution (34) was periodically supplied to the pots. Thirty days after inoculation, a soil suspension (10 g in 30 mL) was added to the vermiculite in the pot (soil addition, S), or the aboveground parts of plants were excised (decapitation, D), or both treatments were performed (DS) (Fig. 2). The aim of the S treatment was to introduce soil microbes into the aseptic pot. That of the D treatment was to stop photosynthate supply to the soybean roots; because field N₂O emission occurred more than 100 days after sowing (55), shoot decapitation was used to promote nodule senescence and degradation. The pots were left in the phytotron until N₂O determination for 15 days except otherwise indicated.

N₂O determination

N₂O flux was determined with a gas chromatograph (GC-14BpsE; Shimadzu, Kyoto, Japan) equipped with a ⁶³Ni electron capture

Fig. 1. Construction of a *nirK* insertion mutant of *Bradyrhizobium japonicum* USDA110. Cloned fragments in pK18mob derivatives are shown alongside the physical map of the *nir* gene cluster of *B. japonicum* USDA110. See text for details.

Fig. 2. Soybean seedlings were inoculated with USDA110 (*nosZ*⁺), denitrification mutants of USDA110, or T9 (*nosZ*⁻). Plants were aseptically grown in Leonard jar pots (Fig. S2) for 30 days after inoculation (DAI). At 30 DAI, treatments were applied. At 40 or 45 DAI, gas phase was sampled for N₂O analysis. In the long term monitoring, gas sampling was continued until 93 DAI. In the tracer experiment, ¹⁵N₂ tracer gas was supplied for 8 h at 29 DAI. See text for details.

detector and tandem columns packed with Porapak Q (80/100 mesh; 3.0 mm×1.0 m and 3.0 mm×2.0 m).

Model system for N₂O emission from degraded nodules

USDA110 (*nosZ*⁺), USDA110Δ*nosZ* (*nosZ*⁻), and T9 (*nosZ*⁻) were used as inoculants. Thirty days after inoculation, treatments were applied (Fig. 2). Ten days later, nodules were collected from soybean roots, washed with sterilized water, and weighed. The nodules were introduced into a 19-mL airtight vial. Gas in the vial was sampled 1, 2, and 3 h after the vials were sealed to determine the N₂O concentration. This was the “excised nodule method” (Fig. 2).

Long-term N₂O monitoring

T9 was used as inoculum. Thirty days after inoculation, the D or DS treatment was applied. The N₂O flux from the pot was intermittently monitored during 2 months. On each measurement day, the pot was sealed with a lid with a gas sampling port (Fig. 2). After 5 h, the gas was sampled to determine N₂O concentration. After the gas sampling, the pot was returned to the phytotron. This was the “sealed jar method” (Fig. 2).

¹⁵N₂ feeding and ¹⁵N determination

At 29 days after inoculation, a gas mixture (30% [v/v] ¹⁵N₂, 20% O₂, 50% Ar; SI Sciences, Tokyo, Japan) containing 32.2 atom% ¹⁵N₂ was supplied to the root zone of soybeans inoculated with USDA110Δ*nosZ* in seven pots for 8 h (Fig. 2 and S1). The nodules from three plants were separately collected and dried at 80°C for 3 days. The ¹⁵N concentrations of the powdered nodules were determined by mass spectrometer (EA 1110 DeltaPlus Advantage ConFlo III; Thermo Fisher Scientific, Bremen, Germany). The other four pots received the DS treatment. Fifteen days later, the gas phase was sampled by the sealed jar method (Fig. 2). The ¹⁵N concentrations were determined by gas chromatography/mass spectrometry (GC/MS-QP2010 Plus; Shimadzu) (21, 22).

N₂O emission from degraded nodules with denitrification mutants

USDA110, USDA110Δ*nosZ*, USDA110Δ*napA*Δ*nosZ*, and USDA110Δ*nirK*Δ*nosZ* were used as inoculants. Thirty days after inoculation, D or DS treatment was applied (Fig. 2). Fifteen days later, the N₂O flux from the nodules was determined by the excised nodule method.

N₂O flux from soybean rhizosphere with denitrification mutants

USDA110 and its Δ*nosZ*, Δ*nirK*, and Δ*nirK*Δ*nosZ* mutants were used as inoculants to evaluate the effect of the *nirK* and *nosZ* genes

on N₂O emission from the rhizosphere. The *nirK* mutation was selected as a nitrate-to-N₂O denitrification mutation, because the *nirK* mutant is not able to denitrify both nitrate and nitrite that exist in the rhizosphere (4). Thirty days after inoculation, DS treatment was applied. Fifteen days later, the N₂O flux from each pot was determined by the sealed jar method, 3 h after the pot was sealed. In addition, 50 mL of 5 mM KNO₃ solution was applied to each pot, the pots were immediately sealed, and the N₂O flux was determined as above.

Results

N₂O emission from degraded nodules

When *B. japonicum* USDA110 (*nosZ*⁺) was used as the inoculum, N₂O was not emitted in any treatment (Fig. 3A). When USDA110Δ*nosZ* or T9 (each *nosZ*⁻) was used, the DS treatment induced marked N₂O emission, whereas the D and S treatments alone did not induce N₂O emission (Fig. 3B and C). Indeed, the nodules in the DS treatment were clearly degraded (Fig. S2), similar to those of field-grown soybean in the late growth period (20). On the other hand, the nodules in the S treatment stayed intact, and those in the D treatment looked slightly degraded (Fig. S2). These results indicate that both soil microbes and nodule degradation are required for N₂O emission. In addition, N₂O was emitted only from DS-treated nodules with *nosZ*⁻ strains, suggesting that the *B. japonicum* *nosZ* gene is critical in the emission of N₂O from degraded nodules.

Long-term monitoring of N₂O flux from the soybean rhizosphere

Substantial N₂O was emitted from the rhizosphere of soybeans inoculated with T9 (*nosZ*⁻) in DS treatment, but none was emitted in D treatment throughout the experimental period (5–63 days) (Fig. 4). This result is similar to the results in the excised nodule method (Fig. 3B and C). As the N₂O flux in the DS treatment peaked 15 days after the treatment was applied and then gradually decreased (Fig. 4), we measured N₂O flux at 15 days in later experiments.

Fig. 3. N_2O flux (excised nodule method) from nodules of soybean inoculated with (A) USDA110, (B) USDA110Δ*nosZ*, or (C) T9 15 days after decapitation (D), soil addition (S), or both (DS). Bars indicate standard error with triplicate biological replications.

Fig. 4. Long-term profile of N_2O flux (sealed jar method) from the rhizosphere of soybean inoculated with T9 (*nosZ*⁻) after decapitation (D) or decapitation plus soil addition (DS). Bars indicate standard error with four biological replications.

Origin of N_2O -N

The profile of N_2O flux (Fig. 4) suggests that the source of N_2O was limited. Thus, we examined whether N_2O is derived from N fixed in the nodules by using ¹⁵N-labeled dinitrogen. The supply of ¹⁵N₂ to the root zone of USDA110Δ*nosZ* plants just before DS treatment produced ¹⁵N concentration in N_2O emitted 15 days later of 1.32±0.42 atom% excess (mean ± SD), similar to the concentration of nodule N (1.13±0.08 atom% excess). This result clearly indicates that the N_2O -N emitted from the soybean rhizosphere was derived from N fixed symbiotically in the nodules.

Fig. 5. N_2O flux (excised nodule method) from nodules of soybean inoculated with (A) USDA110, (B) Δ*nosZ*, (C) Δ*napA*Δ*nosZ*, or (D) Δ*nirK*Δ*nosZ* 15 days after decapitation (D) or decapitation + soil addition (DS). Bars indicate standard error with five biological replications.

N_2O emission from degraded nodules with denitrification mutants

N_2O emissions from the nodules formed with USDA110 and its mutants were determined by the excised nodule method to reveal the involvement of bradyrhizobial denitrification (Fig. 5). Nodules inoculated with Δ*nosZ*, Δ*napA*Δ*nosZ*, and Δ*nirK*Δ*nosZ* emitted marked amounts of N_2O in DS treatment. Nodules inoculated with USDA110 emitted negligible N_2O even in DS treatment (Fig. 5A).

Because the *nosZ* gene is responsible for the reduction of N_2O to N_2 (18, 43), and no N_2O was emitted from *nosZ*⁺ nodules (Figs. 3A and 5A), N_2O reductase encoded by *nosZ* is likely a sink for N_2O in the soybean rhizosphere. In the absence of *nosZ*, N_2O emission from nodules inoculated with double mutants (Δ*napA*Δ*nosZ* and Δ*nirK*Δ*nosZ*) was lower than that from nodules with Δ*nosZ*, although there was no significant difference (Fig. 5B, C, and D, *t*-test [*P*<0.05]).

N_2O flux from the soybean rhizosphere with denitrification mutants

When soybean plants were inoculated with USDA110 and Δ*nirK*, a small quantity of N_2O was released (1.9–2.6 nmol h⁻¹ per pot; Fig. 6A). When plants were inoculated with Δ*nosZ* and Δ*nirK*Δ*nosZ*, N_2O emission was significantly higher (16.7 and 9.9 nmol h⁻¹ per pot, respectively). These results strongly suggest that the *nosZ* gene of *B. japonicum* is involved in the uptake of N_2O that is released from degraded nodules. In Fig. 6A, the relative contribution of the

Fig. 6. N₂O emission (sealed jar method) from the rhizosphere of soybean inoculated with USDA110 or its denitrification mutants ($\Delta nosZ$, $\Delta nirK$, $\Delta nirK\Delta nosZ$) in (A) the absence and (B) the presence of KNO₃. Bars indicate standard error with five biological replications. Differences in N₂O flux are shown as follows: CZ1 and CZ2, contribution of *nosZ* in *B. japonicum*; CK1 (41%) and CK2 (60%), relative contribution of *nirK* under a $\Delta nosZ$ mutant background; CS1 (59%) and CS2 (40%), relative contribution of other soil organisms. Bars labeled with the same letter within a graph are not significantly different (*t*-test, $P < 0.05$).

nosZ gene to N₂O flux is shown as “CZ1”. In the absence of *nosZ*, there was a significant difference in N₂O flux between $\Delta nosZ$ and $\Delta nirK\Delta nosZ$ (CK1 in Fig. 6A). This difference is due to the loss of nitrite reductase in the denitrifying pathway of *B. japonicum*. Therefore, the N₂O flux from soybeans inoculated with $\Delta nosZ$ could have had two distinct sources; denitrification up to N₂O by *B. japonicum* (CK1 [41%] in Fig. 6A), and other soil microbes (CS1 [59%] in Fig. 6A).

KNO₃ was added to the rhizosphere to clarify whether NO₃⁻ is a precursor of N₂O. When KNO₃ was supplied before N₂O determination, the N₂O flux from the pots with each inoculant was markedly enhanced, particularly from pots with $\Delta nosZ$ (78.1 nmol h⁻¹ per pot) and $\Delta nirK\Delta nosZ$ (31.3 nmol h⁻¹ per pot; Fig. 6B). This result confirms that N₂O was produced from NO₃⁻ through microbial denitrification. KNO₃ application also enhanced the contribution of *B. japonicum* to N₂O flux (60% [CK2, Fig. 6B] cf. 41% [CK1, Fig. 6A]). These results suggest that *B. japonicum* prefers nitrate as a substrate for N₂O production.

Discussion

The term “rhizosphere” was first coined in 1904 by Lorenz Hiltner in Germany, who had a special interest in complicated N transformations around leguminous nodules with higher N

Fig. 7. Schematic representation of N₂O metabolism in the soybean rhizosphere induced from the present study. *Bradyrhizobium japonicum* and other soil microorganisms generate N₂O during nodule degradation. *nosZ*⁺ strains of *B. japonicum* are exclusively able to take up N₂O via N₂O reductase. The relative contributions of N₂O emission (CK1 and CK2 in Fig. 6 and text) are shown as percentages at arrows of *B. japonicum* and soil microorganisms. Net N₂O flux is determined by the balance between source and sink. NAP, NO₃⁻ reductase; NIR, NO₂⁻ reductase; NOR, NO reductase; NOS, N₂O reductase.

contents in fields (16). In a sense, the present study advances such historical work on leguminous rhizospheres.

The results show that N₂O emission from degraded nodules in the soybean rhizosphere is due to *B. japonicum* and other soil microbes. When plants were inoculated with *B. japonicum nosZ*⁻ strains and treated with shoot decapitation and soil addition (DS), N₂O was markedly produced (Figs. 3, 4, 5, and 6). On the other hand, when plants were inoculated with a *nosZ*⁺ strain, almost no N₂O was emitted, even in DS treatment. These results suggest that N₂O emission from degrading nodules formed with *nosZ*⁻ strains was due to denitrification by both *B. japonicum* (*nosZ*⁻) and other soil microbes (Fig. 7). It is likely that N₂O produced by soil microbes was offset by *nosZ*-competent *B. japonicum* with its N₂O reductase. In other words, both *B. japonicum* and other soil microorganisms release N₂O during nodule degradation (N₂O source), and *nosZ*-competent *B. japonicum* (*nosZ*⁺ strains) takes up N₂O (N₂O sink) (Fig. 7).

What are these other soil microorganisms that emit N₂O from degraded nodules? Prokaryotic denitrification, fungal denitrification, ammonium oxidation, and nitrate ammonification have been nominated as soil microbial sources of N₂O (1, 14, 26, 38, 49, 50, 57). Community analysis specific to degrading nodules that emit N₂O found many microorganisms that potentially produce N₂O (20), including denitrifying bacteria such as *Acidovorax* (19) and *Enterobacter* (2); *Bradyrhizobium* (25), *Salmonella* (48), *Xanthomonas* (52), and *Pseudomonas* (36), which have functional genes and/or activities for denitrification; and *Fusarium*, a denitrifying fungus (45). Since *Fusarium* species generally lacks N₂O reductase (51), it might be one of the key sources of N₂O from degrading nodules.

The decline in N₂O emission after the peak (Fig. 4) indicates that the source of N in the rhizosphere is limited. Indeed, the ¹⁵N tracer experiment showed that nodule N is a major source of N₂O emission from the soybean rhizosphere. Thus, complicated N transformation in the soybean rhizosphere would involve ammonification, nitrification, and denitrification.

KNO₃ addition enhanced N₂O emission (Fig. 6), supporting the idea that NO₃⁻ is a precursor of N₂O. When NH₄Cl was preliminarily added to the rhizosphere, the addition did not change N₂O emission (Inaba *et al.*, unpublished data), suggesting that it is unlikely to be due to nitrification. KNO₃ addition also enhanced the contribution of *B. japonicum* to N₂O emission in relation to the other soil microbes (Fig. 6). Nitrate might be more available to *B. japonicum*, whereas other microbes might prefer other substrates. In fact, nitrite is a better substrate for denitrifying fungi to produce N₂O (45). New approaches are needed to understand soil N₂O-producing microorganisms and N transformation from fixed nitrogen in the rhizosphere (4).

In soybean fields, it is likely that soybean roots are infected with multiple strains that differ in denitrifying activity. *nosZ*-strains of *B. japonicum* that produce N₂O as the denitrification end product often dominate in agricultural fields (3, 6, 11, 41, 42, 54). Both N₂- and N₂O-producing strains occurred in paddy-upland rotation fields (3). Similarly, both *nosZ*⁺ and *nosZ*⁻ strains of *B. japonicum* were isolated from soybean fields (41, 42). Thus, it is easily conceivable that both N₂- and N₂O-producing strains of *B. japonicum* coexist in soybean fields. Consequently, the flux of N₂O from soybean fields during the late growth period may be partly determined by biotic factors, namely the balance between N₂O emission due to soil microbes and *B. japonicum* (*nosZ*⁻) and N₂O uptake by *B. japonicum* (*nosZ*⁺) (Fig. 7).

The use of *nosZ*⁺ strains of *B. japonicum* as inoculants has been expected to reduce N₂O emissions from soybean fields (42, 43). Indeed, *nosZ*⁺ strains produced no N₂O and were able to take up N₂O from degraded nodules (Fig. 7). Recently, N₂O reduction by *nosZ*-carrying inoculants was shown in a soil-filled pot planted with soybeans (17). Thus, *B. japonicum* mutants with increased N₂OR activity (23) might be more effective to reduce net N₂O flux from soybean rhizosphere.

Acknowledgements

This work was supported in part by the Program for Promotion of Basic Research Activities for Innovative Biosciences (BRAIN), by a grant from the Ministry of Agriculture, Forestry and Fisheries of Japan (Development of Mitigation and Adaptation Techniques to Global Warming, and Genomics for Agricultural Innovation, PMI-0002), and by Grants-in-Aid for Scientific Research (A) 23248052 and for Challenging Exploratory Research 23658057 from the Ministry of Education, Science, Sports and Culture of Japan.

References

1. Amano, T., I. Yoshinaga, T. Yamagishi, C. Van Thuoc, P. The Thu, S. Ueda, K. Kato, Y. Sako, and Y. Suwa. 2011. Contribution of anammox bacteria to benthic nitrogen cycling in a mangrove forest and shrimp ponds, Haiphong, Vietnam. *Microbes Environ.* 26:1–6.
2. Apostoli, P., M. Gelmi, L. Alessio, and A. Turano. 1996. Interferences of urinary tract infection in the measurement of urinary nitrous oxide. *Occup. Environ. Med.* 53:591–594.
3. Asakawa, S. 1993. Denitrifying ability of indigenous strains of *Bradyrhizobium japonicum* isolated from fields under paddy-upland rotation. *Biol. Fertil. Soils* 15:196–200.
4. Baggs, E.M. 2011. Soil microbial sources of nitrous oxide: Recent advances in knowledge, emerging challenges and future direction. *Curr. Opin. Environ. Sustain.* 3:321–327.
5. Bedmar, E.J., E.F. Robles, and M.J. Delgado. 2005. The complete denitrification pathway of the symbiotic, nitrogen-fixing bacterium *Bradyrhizobium japonicum*. *Biochem. Soc. Trans.* 33:141–144.
6. Breitenbeck, G.A., and J.M. Bremner. 1989. Ability of free-living cells of *Bradyrhizobium japonicum* to denitrify in soils. *Biol. Fertil. Soils* 7:219–224.
7. Ciampitti, I.A., E.A. Ciarlo, and M.E. Conti. 2008. Nitrous oxide emissions from soil during soybean (*Glycine max* (L.) Merrill) crop phenological stages and stubbles decomposition period. *Biol. Fertil. Soils* 44:581–588.
8. Cole, M.A., and G.H. Elkan. 1973. Transmissible resistance to penicillin G, neomycin, and chloramphenicol in *Rhizobium japonicum*. *Antimicrob. Agents Chemother.* 4:248–253.
9. Dao, T., M. Nomura, R. Hamaguchi, *et al.* 2008. NAD-malic enzyme affects nitrogen-fixing activity of *Bradyrhizobium japonicum* USDA110 bacteroids in soybean nodules. *Microbes Environ.* 23:215–220.
10. Duxbury, J.M., D.R. Bouldin, R.E. Terry, and R.L. Tate III. 1982. Emissions of nitrous oxide from soils. *Nature* 298:462–464.
11. Fernández, L.A., E.B. Perotti, M.A. Sagardoy, and M.A. Gómez. 2008. Denitrification activity of *Bradyrhizobium* sp. isolated from Argentine soybean cultivated soils. *World J. Microbiol. Biotechnol.* 24:2577–2585.
12. Figurski, D.H., and D.R. Helinski. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in trans. *Proc. Natl. Acad. Sci. U.S.A.* 76:1648–1652.
13. Firestone, M.K., R.B. Firestone, and J.M. Tiedje. 1980. Nitrous oxide from soil denitrification: factors controlling its biological production. *Science* 208:749–751.
14. Forster, P., V. Ramaswamy, P. Artaxo, *et al.* 2007. Changes in atmospheric constituents and in radiative forcing, p. 129–234. In Solomon DQ (ed.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, New York.
15. Ghosh, S., D. Majumdar, and M.C. Jain. 2002. Nitrous oxide emissions from *kharif* and *rabi* legumes grown on an alluvial soil. *Biol. Fertil. Soils* 35:473–478.
16. Hartmann, A., M. Rothballer, and M. Schmid. 2008. Lorenz Hiltner, a pioneer in rhizosphere microbial ecology and soil bacteriology research. *Plant Soil* 312:7–14.
17. Hénault, C., and C. Revellin. 2011. Inoculants of leguminous crops for mitigating soil emissions of the greenhouse gas nitrous oxide. *Plant Soil* 346:289–296.
18. Hirayama, J., S. Eda, H. Mitsui, and K. Minamisawa. 2011. Nitrate-dependent N₂O emission from intact soybean nodules via denitrification by *Bradyrhizobium japonicum* bacteroids. *Appl. Environ. Microbiol.* 77:8787–8790.
19. Hoshino, T., T. Terahara, S. Tsuneda, A. Hirata, and Y. Inamori. 2005. Molecular analysis of microbial population transition associated with the start of denitrification in a wastewater treatment process. *J. Appl. Microbiol.* 99:1165–1175.
20. Inaba, S., K. Tanabe, S. Eda, S. Ikeda, A. Higashitani, H. Mitsui, and K. Minamisawa. 2009. Nitrous oxide emissions and microbial community in the rhizosphere of nodulated soybeans during the late growth period. *Microbes Environ.* 24:64–67.
21. Isobe, K., Y. Suwa, J. Ikutani, *et al.* 2011. Analytical techniques for quantifying ¹⁵N/¹⁴N of nitrate, nitrite, total dissolved nitrogen and ammonium in environmental samples using a gas chromatograph equipped with a quadrupole mass spectrometer. *Microbes Environ.* 26:46–53.
22. Isobe, K., K. Koba, S. Ueda, K. Senoo, S. Harayama, and Y. Suwa. 2011. A simple and rapid GC/MS method for the simultaneous determination of gaseous metabolites. *J. Microbiol. Methods* 84:46–51.
23. Itakura, M., K. Tabata, S. Eda, H. Mitsui, K. Murakami, J. Yasuda, and K. Minamisawa. 2008. Generation of *Bradyrhizobium japonicum* mutants with increased N₂O reductase activity by selection after introduction of a mutated *dnaQ* gene. *Appl. Environ. Microbiol.* 74:7258–7264.
24. Jumadi, O., Y. Hala, A. Muis, A. Ali, M. Palennari, K. Yagi, and K. Inubushi. 2008. Influences of chemical fertilizers and a nitrification inhibitor on greenhouse gas fluxes in a corn (*Zea mays* L.) field in Indonesia. *Microbes Environ.* 23:29–34.

25. Kaneko, T., Y. Nakamura, S. Sato, *et al.* 2002. Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res.* 9:189–197.
26. Katsuyama, C., N. Kondo, Y. Suwa, T. Yamagishi, M. Itoh, N. Ohte, H. Kimura, K. Nagaosa, and K. Kato. 2008. Denitrification activity and relevant bacteria revealed by nitrite reductase gene fragments in soil of temperate mixed forest. *Microbes Environ.* 23:337–345.
27. Kim, Y., N. Nakayama, T. Nakamura, M. Takahashi, S. Shimada, and J. Arihara. 2005. NO and N₂O emissions from fields in the different nodulated genotypes of soybean (in Japanese). *Jpn. J. Crop Sci.* 74:427–430.
28. Leonard, L.T. 1943. A simple assembly for use in the testing of cultures of rhizobia. *J. Bacteriol.* 45:523–525.
29. Lim, B.L. 2010. TonB-dependent receptors in nitrogen-fixing nodulating bacteria. *Microbes Environ.* 25:67–74.
30. Masuda, S., S. Eda, C. Sugawara, H. Mitsui and K. Minamisawa. 2010. The *cbbL* gene is required for thiosulfate-dependent autotrophic growth of *Bradyrhizobium japonicum*. *Microbes Environ.* 25:220–223.
31. Minamisawa, K., M. Itakura, M. Suzuki, K. Ichige, T. Isawa, K. Yuhashi, and H. Mitsui. 2002. Horizontal transfer of nodulation genes in soil and microcosmos from *Bradyrhizobium japonicum* to *B. elkanii*. *Microbes Environ.* 17:82–90.
32. Mori, A., M. Hojito, H. Kondo, H. Matsunami, and D. Scholefield. 2005. Effects of plant species on CH₄ and N₂O fluxes from a volcanic grassland soil in Nasu, Japan. *Soil Sci. Plant Nutr.* 51:19–27.
33. Okabe, S., M. Oshiki, Y. Kamagata, *et al.* 2010. A great leap forward in microbial ecology. *Microbes Environ.* 25:230–240.
34. Okazaki, S., M. Sugawara, and K. Minamisawa. 2004. *Bradyrhizobium elkanii* *rtxC* gene is required for expression of symbiotic phenotypes in the final step of rhizobitoxine biosynthesis. *Appl. Environ. Microbiol.* 70:535–541.
35. Okubo, T., S. Ikeda, A. Yamashita, K. Terasawa, and K. Minamisawa. 2012. Pyrosequence read length of 16S rRNA gene affects phylogenetic assignment of plant-associated bacteria. *Microbes Environ.* 27:204–208.
36. Paulsen, I.T., C.M. Press, J. Ravel, *et al.* 2006. Complete genome sequence of the plant commensal *Pseudomonas fluorescens* Pf-5. *Nat. Biotechnol.* 23:873–878.
37. Prentki, P., and H.M. Krisch. 1984. *In vitro* insertional mutagenesis with a selectable DNA fragment. *Gene* 29:303–313.
38. Ryuda, N., T. Hashimoto, D. Ueno, K. Inoue, and T. Someya. 2011. Visualization and direct counting of individual denitrifying bacterial cells in soil by *nirK*-targeted direct in situ PCR. *Microbes Environ.* 26:74–80.
39. Saeki, Y., S. Ozumi, A. Yamamoto, Y. Umehara, M. Hayashi, and G.C. Sigua. 2010. Changes in population occupancy of bradyrhizobia under different temperature regimes. *Microbes Environ.* 25:309–312.
40. Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
41. Sameshima-Saito, R., K. Chiba, and K. Minamisawa. 2004. New method of denitrification analysis of *Bradyrhizobium* field isolates by gas chromatographic determination of ¹⁵N-labeled N₂. *Appl. Environ. Microbiol.* 70:2886–2891.
42. Sameshima-Saito, R., K. Chiba, and K. Minamisawa. 2006. Correlation of denitrifying capability with the existence of *nap*, *nir*, *nor* and *nos* genes in diverse strains of soybean bradyrhizobia. *Microbes Environ.* 21:174–184.
43. Sameshima-Saito, R., K. Chiba, J. Hirayama, M. Itakura, H. Mitsui, S. Eda, and K. Minamisawa. 2006. Symbiotic *Bradyrhizobium japonicum* reduces N₂O surrounding the soybean root system via nitrous oxide reductase. *Appl. Environ. Microbiol.* 72:2526–2532.
44. Schäfer, A., A. Tauch, W. Jäger, J. Kalinowski, G. Thierbach, and A. Pühler. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145:69–73.
45. Shoun, H., D.H. Kim, H. Uchiyama, and J. Sugiyama. 1992. Denitrification by fungi. *FEMS Microbiol. Lett.* 73:277–281.
46. Smith, K.A., P.J. Crutzen, A.R. Mosier, and W. Winiwarter. 2010. The global nitrous oxide budget: A reassessment, p 63–84. *In* Smith KA (ed.), *Nitrous Oxide and Climate Change*. Earthscan, London, UK.
47. Smith, P., and D. Martino. 2007. Chapter 8. Agriculture, p. 506–507. *In* B. Mets, O. Davidson, P. Bosch, R. Dave, and L. Meyer (ed.), *Climate Change 2007 Mitigation, The Intergovernmental Panel on Climate Change*. Cambridge University Press, New York, NY.
48. Spector, M.P., F. Garcia del Portillo, S.M. Bearson, A. Mahmud, M. Magut, B.B. Finlay, G. Dougan, J.W. Foster, and M.J. Pallen. 1999. The *rpoS*-dependent starvation stress response locus *stiA* encodes a nitrate reductase (*narZYWV*) required for carbon-starvation-inducible thermotolerance and acid tolerance in *Salmonella typhimurium*. *Microbiology* 145:3035–3045.
49. Tago, K., S. Ishii, T. Nishizawa, S. Otsuka, and K. Senoo. 2011. Phylogenetic and functional diversity of denitrifying bacteria isolated from various rice paddy and rice-soybean rotation fields. *Microbes Environ.* 26:30–35.
50. Takahashi, M., T. Yamada, M. Tanno, H. Tsuji, and A. Hiraishi. 2011. Nitrate removal efficiency and bacterial community dynamics in denitrification processes using poly (L-lactic acid) as the solid substrate. *Microbes Environ.* 26:212–219.
51. Takaya, N., and H. Shoun. 2000. Nitric oxide reduction, the last step in denitrification by *Fusarium oxysporum*, is obligatorily mediated by cytochrome P450_{nor}. *Mol. Gen. Genet.* 263:342–348.
52. Thieme, F., R. Koebe, T. Bekel, *et al.* 2005. Insights into genome plasticity and pathogenicity of the plant pathogenic bacterium *Xanthomonas campestris* pv. *vesicatoria* revealed by the complete genome sequence. *J. Bacteriol.* 187:7254–7266.
53. Trung, B.C., and S. Yoshida. 1983. Improvement of Leonard jar assembly for screening of effective *Rhizobium*. *Soil Sci. Plant Nutr.* 29:97–100.
54. van Berkum, P., and H.H. Kayser. 1985. Anaerobic growth and denitrification among different serogroups of soybean rhizobia. *Appl. Environ. Microbiol.* 49:772–777.
55. Yang, L., and Z. Cai. 2005. The effect of growing soybean (*Glycine max. L.*) on N₂O emission from soil. *Soil Biol. Biochem.* 37:1205–1209.
56. Ye, B., A. Saito, and K. Minamisawa. 2005. Effect of inoculation with anaerobic nitrogen-fixing consortium on salt tolerance of *Miscanthus sinensis*. *Soil Sci. Plant Nutr.* 51:243–249.
57. Yoshida, M., S. Ishii, S. Otsuka, and K. Senoo. 2010. *nirK*-harboring denitrifiers are more responsive to denitrification-inducing conditions in rice paddy soil than *nirS*-harboring bacteria. *Microbes Environ.* 25:45–48.
58. Zhaorigetu, M. Komatsuzaki, Y. Sato, and H. Ohta. 2008. Relationships between fungal biomass and nitrous oxide emission in upland rice soils under no tillage and cover cropping systems. *Microbes Environ.* 23:201–208.