Short Communication

Identification of the Hydrogen Uptake Gene Cluster for Chemolithoautotrophic Growth and Symbiosis Hydrogen Uptake in *Bradyrhizobium diazoefficiens*

SACHIKO MASUDA^{1*†}, MASAKI SAITO¹, CHIAKI SUGAWARA¹, MANABU ITAKURA¹, SHIMA EDA¹, and KIWAMU MINAMISAWA¹

¹Graduate School of Life Sciences, Tohoku University, Katahira, Aoba-ku, Sendai, Miyagi 980–8577, Japan

(Received November 6, 2015—Accepted December 18, 2015—Published online February 25, 2016)

The hydrogen uptake (Hup) system of *Bradyrhizobium diazoefficiens* recycles the H₂ released by nitrogenase in soybean nodule symbiosis, and is responsible for H₂-dependent chemolithoautotrophic growth. The strain USDA110 has two *hup* gene clusters located outside (locus I) and inside (locus II) a symbiosis island. Bacterial growth under H₂-dependent chemolithoautotrophic conditions was markedly weaker and H₂ production by soybean nodules was markedly stronger for the mutant of *hup* locus I ($\Delta hupS_1L_1$) than for the mutant of *hup* locus II ($\Delta hupS_2L_2$). These results indicate that locus I is primarily responsible for Hup activity.

Key words: *Bradyrhizobium diazoefficiens*, chemolithoautotrophic growth, hydrogenase, symbiosis

Soybean bradyrhizobia have two lifestyles: as symbiotic bacteroids that fix atmospheric nitrogen in host plants or as free-living soil bacteria. As a symbiont, *B. diazoefficiens* synthesizes a hydrogen uptake (Hup) system that recycles the H₂ formed as a byproduct of nitrogenase activity (4). This symbiotic hydrogen oxidation increases nitrogen fixation efficiency, thereby enhancing the productivity of the legume host (3, 6). As free-living cells, *Bradyrhizobium diazoefficiens* Hup⁺ strains have the ability to grow chemolithoautotrophically by using H₂ as an electron donor (7).

Two sets of *hup* genes have been identified in *B. diazoefficiens*: a large cluster outside the symbiosis island (*hup* locus I, genome position 7,620,025–7,645,755) and a small cluster inside the symbiosis island (*hup* locus II, genome position 1,888,916–1,902,575) (6, 11). A transcriptome analysis previously showed that several *hup* genes located outside the symbiosis island were up-regulated during H₂-dependent chemolithoautotrophic growth, whereas several *hup* genes located inside the symbiosis island were up-regulated in symbiotic bacteroids (1, 6). These findings imply that *hup* locus I plays an important role in chemolithoautotrophic growth, while symbiotic *hup* locus II may contribute to Hup activity in the nodules. In the present study, we constructed *hupSL* deletion mutants in order to clarify the contribution of each *hup* gene cluster during the chemolithoautotrophic growth and nodulation of *B. diazoefficiens* USDA110.

Materials and Methods

The strains and plasmids used in this study are listed in Table 1. The HM salt medium for the preculture and Hup medium for chemolithoautotrophic growth were described previously (14). Antibiotics were added to the media for *B. diazoefficiens* USDA110 and *Escherichia coli* strains as described previously (13).

We generated *hupS₁L₁* and *hupS₂L₂* deletion mutants as follows. A DNA fragment containing *hupFDCL₁S₁V* was isolated from brp15657, a plasmid from the pUC18 clone library of *B. diazoefficiens* USDA110 sequences (11), and inserted into the *Hind*III site of pK18mob. The resultant plasmid, pK18mob-*hupI*, was digested with *Apa*I and ligated with the *Apa*I/*Eco*RI adaptor, yielding pK18mob-*hupI* Δ . The omega cassette isolated from pHP45 Ω was inserted at the *Eco*RI site of pK18mob-*hupI* Δ , yielding pK18mob-*hupI*omega. Triparental mating using pRK2013 was performed as described previously (13). A similar strategy was used to construct the *hupS₂L₂* deletion mutant. Briefly, *hupKHFDC₂S₂* was isolated from brp07423, and inserted into pK18mob, yielding pK18mob-*hupII*. pK18mob-*hupII* was ligated with the *Eco*O109I/*Bam*HI adaptor, resulting in pK18mob-*hupII* Δ . The Tc^r cassette was isolated from p34S-Tc and inserted into the *Bam*HI site of pK18mob-*hupII* Δ , yielding pK18mob-*hupII*Tc. The double crossover events of these deletion mutations were verified by a Southern blot analysis.

The inoculants were prepared as described previously (14). Aliquots (10 μ L) of the cells (OD₆₆₀ at 0.1) were streaked on Hup medium, and the agar plates were statically incubated at 25°C for 14 d in an atmosphere containing 1% O₂, 5% CO₂, 10% H₂, and 84% N₂.

Soybean seedlings (*Glycine max* cv. Enrei) grown in a plant box in a growth chamber were inoculated with *B. diazoefficiens* as described previously (8, 9). The nodulated roots were transferred into a 300-mL bottle 30 d later, and a 0.5-mL gas sample from the head space of the bottle was injected into a GC-2014 gas chromatograph (Shimadzu, Kyoto, Japan) as described previously (13). The flow rate of the carrier gas (N₂) was 30 mL min⁻¹.

Results

Wild-type USDA110 and the $\Delta hupS_2L_2$ mutant grew on Hup agar medium under chemolithoautotrophic growth conditions (Fig. 2A). However, the $\Delta hupS_1L_1$ mutant showed markedly weaker growth than that of the wild type on this medium (Fig. 2A). The height of plants inoculated with $\Delta hupS_1L_1$ appeared to be lower than those inoculated with the wild type and $\Delta hupS_2L_2$ mutant (Fig. 2B); however, no significant differences were observed in plant dry weights or fresh nodule weights (Table 2). H₂ was not produced from the nodulating roots of the wild type or $\Delta hupS_2L_2$ mutant (Fig. 2C), indicating that Hup activity compensated for the production of H₂ via

* Corresponding author. E-mail: smasuda@cc.tuat.ac.jp;
Tel/Fax: +81-42-367-5847.

† Present address: Graduate School of Agriculture, Tokyo University of Agriculture and Technology, Saiwaicho, Fuchu, Tokyo 183–8509, Japan

Table 1. Strains and plasmids used in the present study.

Strain or plasmid	Relevant characteristics ^a	Reference or source
<i>Bradyrhizobium diazoefficiens</i>		
USDA110	Soybean bradyrhizobia, wild type	11
$\Delta hupS_1L_1$	USDA110 <i>hupS₁L₁::aadA</i> ; Sm ^r Sp ^r	This study
$\Delta hupS_2L_2$	USDA110 <i>hupS₂L₂::tet</i> ; Tc ^r	This study
<i>Escherichia coli</i>		
DH5 α	cloning strain	Toyobo Inc. ^a
Plasmids		
p34S-Tc	Plasmid carrying a 2.1-kb Tc cassette; Tc ^r	2
pHP45 Ω	Plasmid carrying a 2.1-kb Ω cassette; Sp ^r Sm ^r Ap ^r	15
pK18mob- <i>hupI</i>	pK18mob carrying a 5.9-kb <i>hupFDCL₁S₁VU</i> HindIII fragment of brp15657; Km ^r	This study
pK18mob- <i>hupI</i> ad	pK18mob- <i>hupI</i> with an <i>ApaI/EcoRI</i> adaptor; Km ^r	This study
pK18mob- <i>hupI</i> omega	pK18mob carrying <i>hupS₁L₁::aadA</i> ; Sm ^r Sp ^r Km ^r	This study
pK18mob- <i>hupII</i>	pK18mob carrying a 7.6-kb <i>hupS₂L₂CDFHK</i> XhoI fragment of brp07428; Tc ^r	This study
pK18mob- <i>hupII</i> ad	pK18mob- <i>hupII</i> with an <i>EcoO109I/BamHI</i> adaptor; Km ^r	This study
pK18mob- <i>hupII</i> Tc	pK18mob carrying <i>hupS₂L₂::tet</i> ; Tc ^r Km ^r	This study
pK18mob	cloning vector; pMB1ori oriT; Km ^r	16
pRK2013	ColE1 replicon carrying RK2 transfer genes; Km ^r	5
brp15657	pUC18 carrying <i>hupUVS₁L₁CDFG</i>	11
brp07423	pUC18 carrying <i>hupKHFDC₂S₂</i>	11

Ap^r, ampicillin-resistant; Tc^r, tetracycline-resistant; Km^r, kanamycin-resistant; Sm^r, streptomycin-resistant; Sp^r, spectinomycin-resistant.

^a Osaka, Japan.

Fig. 1. Gene maps of *hup* gene clusters in *B. diazoefficiens*. (A) Gene map of *hup* locus I showing the genome position of *hup* locus I and brp15657. pK18mob-*hupI* carries a 5.9-kb DNA fragment containing *hupFDCL₁S₁V* genes. pK18mob-*hupI*ad was ligated with an *ApaI/EcoRI* adaptor. pK18mob-*hupI*omega had an omega cassette, which was inserted at the *EcoRI* site of pK18mob-*hupI*ad. (B) Gene map of *hup* locus II showing the genome position of *hup* locus II and brp07428. The strategy used to construct the *hupS₂L₂* deletion mutant is shown as the *hupS₁L₁* deletion mutant. H, *HindIII*; A, *ApaI*; E, *EcoRI*; X, *XhoI*; Ec, *EcoO109I*; B, *BamHI*; Tc^r, tetracycline-resistant.

nitrogenase. In contrast, H₂ was produced by the $\Delta hupS_1L_1$ nodules (6.4 $\mu\text{mol h}^{-1} \text{g}$ fresh nodule weight⁻¹) (Fig. 2C), indicating that Hup activity was lower than the production of H₂ via nitrogenase. These results suggest that *hup* genes outside the symbiosis island are the primary cluster involved in chemolithoautotrophic growth and Hup activity in the nodules.

Discussion

In the present study, the mutation of *hupS₂L₂* did not change nodule H₂ production from that by wild-type USDA110 (Fig. 2C), even though some genes on *hup* locus II were up-regulated in symbiotic bacteroids (1, 6). *Hup* locus I contains a complete set of the *hup-hyp-hox* cluster, and is missing from *hup* locus II (11). Thus, *hupS₂L₂* in locus II may not be fully induced without the *hup* gene assemblage, resulting in no or weak Hup activity by the *hupS₂L₂* genes. On the other hand, the *hup* gene cluster outside the symbiosis island, which we identified as the primary *hup* gene cluster contributing to Hup activity in free-living and symbiotic cells, is located on a typical genomic island (trnM element) of *B. diazoefficiens* USDA110. The trnM element is likely acquired in the USDA110 lineage after the divergence of strains USDA110 and USDA6^T because *B. japonicum* USDA6^T completely lacks this element (10, 11, 12).

Hup genes were also found in the symbiosis island of the USDA6^T genome even though USDA6^T was previously reported to exhibit no Hup activity (10, 11, 12). The *hup* genes in USDA6^T on the symbiosis island had 99–100% amino acid sequence identity to the corresponding genes in *hup* locus II of USDA110. In contrast, USDA6^T *hup* genes had only 43–83% amino acid sequence identity to genes in *hup* locus I of USDA110, which is similar to the homology (45–83%) between *hup* genes in loci I and II of USDA110. These results suggest that *hup* genes on locus II of USDA110 and *hup* genes in USDA6^T do not contribute to the Hup activities of these strains and appear to be derived from the acquisition of symbiosis islands. Therefore, our results imply the horizontal gene transfer of the primary *hup* cluster via the genomic

Fig. 2. Comparison of chemolithoautotrophic growth (A) and symbiotic phenotypes (BC) between *hup* mutants and the wild-type strain of *Bradyrhizobium diazoefficiens* USDA110. (A) Growth phenotype on Hup agar medium under an atmosphere of 84% N_2 , 10% H_2 , 5% CO_2 , and 1% O_2 at 25°C. (B) Plant growth 30 d after inoculation. (C) Concentration of H_2 produced by the root nodules. White squares, wild type; black squares, $\Delta hupS_1L_1$ mutant; grey squares, $\Delta hupS_2L_2$ mutant.

Table 2. Plant dry weights and fresh nodule weights of inoculated wild-type USDA110 and mutants.

Strains	Plant dry weight (g)	Fresh nodule weight (g)
USDA110	6.9 ± 0.9 ^a	1.36 ± 0.20 ^a
$\Delta hupS_1L_1$	6.8 ± 1.1 ^a	1.38 ± 0.20 ^a
$\Delta hupS_2L_2$	6.9 ± 0.6 ^a	1.29 ± 0.13 ^a

Plants were harvested 30 d after inoculation. Values are presented as an average and standard deviation ($n=5$). Tukey's multiple comparison test was used for statistical analyses ($p<0.05$).

island to the lineage of *B. diazoefficiens* rather than symbiosis island transfer.

Acknowledgements

This study was supported by a grant from the Ministry of Agriculture, Forestry, and Fisheries of Japan (BRAIN) and by Grants-in-Aid for Scientific Research (A) 26252065 from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Chang, W.S., W.L. Franck, E. Cytryn, S. Jeong, T. Joshi, D.W. Emerich, M.J. Sadowsky, D. Xu, and G. Stacey. 2007. An oligonucleotide microarray resource for transcriptional profiling of *Bradyrhizobium japonicum*. *Mol. Plant-Microbe Interact.* 20:1298–1307.
- Dennis, J.J., and G.J. Zylstra. 1998. Plasposons: modular self-cloning minitransposon derivatives for rapid genetic analysis of gram-negative bacterial genomes. *Appl. Environ. Microbiol.* 64:2710–2715.
- Drevon, J., V.C. Kalia, M. Heckmann, and I. Salsac. 1987. Influence of the *Bradyrhizobium* hydrogenase on the growth of *Glycine* and *Vigna* species. *Appl. Environ. Microbiol.* 53:610–612.
- Evans, H.J., A.R. Harker, H. Papen, S.A. Russell, F.J. Hanus, and M. Zuber. 1987. Physiology, biochemistry, and genetics of the uptake hydrogenase in rhizobia. *Annu. Rev. Microbiol.* 41:335–361.
- Figurski, D.H., and D.R. Helinski. 1979. Replication of an origin-containing derivative of plasmid RK2 dependent on a plasmid function provided in *trans*. *Proc. Natl. Acad. Sci. U.S.A.* 76:1648–1652.
- Franck, W.L., W.S. Chang, J. Qiu, M. Sugawara, M.J. Sadowsky, S.A. Smith, and G. Stacey. 2008. Whole-genome transcriptional profiling of *Bradyrhizobium japonicum* during chemoautotrophic growth. *J. Bacteriol.* 190:6697–6705.
- Hanus, F.J., R.J. Maier, and H.J. Evans. 1979. Autotrophic growth of H_2 -uptake-positive strains of *Rhizobium japonicum* in an atmosphere supplied with hydrogen gas. *Proc. Natl. Acad. Sci. U.S.A.* 76:1788–1792.
- Inaba, S., F. Ikenishi, M. Itakura, M. Kikuchi, S. Eda, N. Chiba, C. Katayama, Y. Suwa, H. Mitsui, and K. Minamisawa. 2012. N_2O emission from degraded soybean nodules depends on denitrification by *Bradyrhizobium japonicum* and other microbes in the rhizosphere. *Microbes Environ.* 27:470–476.
- Itakura, M., K. Tabata, S. Eda, H. Mitsui, K. Murakami, J. Yasuda, and K. Minamisawa. 2008. Generation of *Bradyrhizobium japonicum* mutants with increased N_2O reductase activity by selection after inoculation of a mutated *dnaQ* gene. *Appl. Environ. Microbiol.* 74:7258–7264.
- Itakura, M., K. Saeki, H. Omori, *et al.* 2008. Genomic comparison of *Bradyrhizobium japonicum* strains with different symbiotic nitrogen-fixing capabilities and other Bradyrhizobiaceae members. *ISME J.* 3:326–339.
- Kaneko, T., Y. Nakamura, S. Sato, *et al.* 2002. Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res.* 9:189–197.
- Kaneko, T., H. Maita, H. Hirakawa, N. Uchiike, K. Minamisawa, A. Watanabe, and S. Sato. 2011. Complete genome sequence of the soybean symbiont *Bradyrhizobium japonicum* strain USDA6^T. *Gene.* 2:763–787.
- Masuda, S., S. Eda, S. Ikeda, H. Mitsui, and K. Minamisawa. 2010. Thiosulfate-dependent chemolithoautotrophic growth in *Bradyrhizobium japonicum*. *Appl. Environ. Microbiol.* 76:2402–2409.
- Masuda, S., S. Eda, C. Sugawara, H. Mitsui, and K. Minamisawa. 2010. The *cbbL* gene is required for thiosulfate-dependent autotrophic growth of *Bradyrhizobium japonicum*. *Microbes Environ.* 25:220–223.
- Prentki, P., and H.M. Krisch. 1984. *In vitro* insertional mutagenesis with a selective DNA fragment. *Gene.* 29:303–313.
- Schäfer, A., A. Tauch, W. Jäger, J. Kalinowski, G. Thierbach, and A. Pühler. 1994. Small mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* 145:69–73.