The Honeybee Gut Microbiota Is Altered after Chronic Exposure to Different Families of Insecticides and Infection by *Nosema ceranae*

RÉGIS ROUZÉ¹, ANNE MONÉ¹, FRÉDÉRIC DELBAC¹, LUC BELZUNCES², and NICOLAS BLOT^{1*}

¹Université Clermont Auvergne, CNRS, Laboratoire “Microorganismes: Génome et Environnement”, F-63000 Clermont–Ferrand, France; and ²INRA, Laboratoire de Toxicologie Environnementale, UR 406 A&E, CS 40509, 84914 Avignon, France.

(Received December 21, 2018—Accepted April 1, 2019—Published online August 3, 2019)

The gut of the European honeybee *Apis mellifera* is the site of exposure to multiple stressors, such as pathogens and ingested chemicals. Therefore, the gut microbiota, which contributes to host homeostasis, may be altered by these stressors. The abundance of major bacterial taxa in the gut was evaluated in response to infection with the intestinal parasite *Nosema ceranae* or chronic exposure to low doses of the neurotoxic insecticides coumaphos, fipronil, thiamethoxam, and imidacloprid. Experiments were performed under laboratory conditions on adult workers collected from hives in February (winter bees) and July (summer bees) and revealed season-dependent changes in the bacterial community composition. *N. ceranae* and a lethal fipronil treatment increased the relative abundance of both *Gilliamella apicola* and *Snodgrassella alvi* in surviving winter honeybees. The parasite and a sublethal exposure to all insecticides decreased the abundance of *Bifidobacterium* spp. and *Lactobacillus* spp. regardless of the season. The similar effects induced by insecticides belonging to distinct molecular families suggested a shared and indirect mode of action on the gut microbiota, possibly through aspecific alterations in gut homeostasis. These results demonstrate that infection and chronic exposure to low concentrations of insecticides may affect the honeybee holobiont.

Key words: honeybee microbiota, insecticides, parasite, dysbiosis

Insects are exposed to many stressors in their environment, particularly their gut since it is both the site of development of many pathogens and the site of exposure to ingested xenobiotics. At the interface between the environment and the insect, the gut microbiota is sensitive to biotic and abiotic stressors. The consequences for the host of dysbiosis may be pleiotropic because the microbiota is connected to the physiologies of the gut and nervous and immune systems (56, 71).

The environmental stressors of the European honeybee (*Apis mellifera*) have attracted interest due to the ecological and economical importance of this pollinator as well as the colony losses that have been observed worldwide over the last few decades (19, 79). The origin of these losses is assumed to be multicausal, involving pathogens, such as the highly prevalent gut parasite *Nosema ceranae*, and agrochemical contaminants, particularly neurotoxic insecticides (34, 39).

N. ceranae is a fungal-related parasite belonging to *Microsporidia*. This obligate intracellular parasite proliferates within the epithelial cells of the adult honeybee midgut and affects metabolic, digestive, and immune functions as well as the gut epithelium renewal rate (5, 39, 64). Consequently, *N. ceranae* may reduce the honeybee lifespan and provoke colony depopulation.

Honeybees are also exposed to a number of pesticides through the foraging of contaminated pollen and nectar that are brought back to the colony (10). Some chemicals are deliberately introduced into hives by beekeepers, particularly the acaricides used against the ectoparasite *Varroa destructor*. These pesticides may be detected in hive matrices, including wax, pollen, and honey (8, 18, 51, 62). Therefore, all colony members are chronically exposed, particularly through their

diet. The most abundant and ubiquitous contaminants of matrices are varroacides. Systemic insecticides commonly used in agriculture, including neonicotinoids and phenylpyrazoles, are also frequently detected in hives. These compounds are neurotoxic insecticides (14). Even at low doses, they may alter honeybee behavior, learning, and memory and induce an immune response as well as increases in detoxifying activities (9, 17, 20, 32, 33). Some of these effects, including cell damage, have also been observed in the midgut (5, 13, 15, 63).

N. ceranae and ingested pesticides are in contact with the honeybee gut and may alter its physiology. Therefore, the composition and function of gut microbial communities may be affected by these stressors (35, 44, 53). The worker honeybee gut core microbiota, which is shared among individuals, is composed of five major bacterial taxa comprising the *Lactobacillus* Firm-4 and *Lactobacillus* Firm-5 clades, the *Gammaproteobacteria* *Gilliamella apicola* and *Snodgrassella alvi*, and species related to the *Bifidobacterium asteroides* cluster (49). Other taxa, although less abundant, are also present, particularly *Alphaproteobacteria*. Most of the 10⁸ to 10⁹ intestinal bacteria are localized within the hindgut. The crop and midgut are less colonized (36). The intestinal microbiota contributes to host homeostasis. It stimulates the immune system, constitutes a biological barrier against pathogens, and participates in digestion (29, 49, 50, 52, 80). Thus, the impact of stressors on the honeybee and its gut may be explained, at least partially, by an imbalanced microbiota, *i.e.* dysbiosis.

In the present study, we investigated how the composition of the honeybee gut microbiota may be affected under laboratory conditions after an infection with *N. ceranae* or after a chronic exposure to neurotoxic insecticides belonging to three families: the phenylpyrazole fipronil, the neonicotinoids thiamethoxam and imidacloprid, and the organophosphate acaricide coumaphos. Quantitative real-time PCR (qPCR) was used to assess changes

* Corresponding author. E-mail: nicolas.blot@uca.fr;
Tel: +33 4 73 40 74 57.

in the abundance of the major bacterial groups of the honeybee core microbiota, namely, *Lactobacillus* spp., *Bifidobacterium* spp., *Alphaproteobacteria* and *Gammaproteobacteria*, and the species *G. apicola* and *S. alvi*.

Materials and Methods

Honeybee artificial rearing, infection with N. ceranae, and chronic exposure to insecticides

Two experiments were performed using *A. mellifera* interior worker bees collected on the frames of five colonies of the same apiary at the Laboratoire Microorganismes: Génome et Environnement (Clermont-Ferrand, France) in February 2017 (winter bees) and July 2017 (summer bees). Each of the five colonies represented an independent replicate. Honeybees were anesthetized with CO₂ and placed into groups of 70 individuals in Pain-type cages containing a 5-mm piece of PseudoQueen (Inkto Supply, Vancouver, Canada). Honeybees were maintained at 33±1°C under a humidified atmosphere and fed 50% (w/v) sucrose syrup supplemented with 1% (w/v) Provita'Bee (ATZ Diététique, Mas Cabardès, France) *ad libitum*, as previously described (4). Dead bees were removed and sucrose consumption was quantified daily.

Seven experimental groups, including an untreated control, were tested for each colony replicate. Honeybees were collectively infected using a feeder containing approximately 150,000 spores per bee in 2.5 mL of sugar syrup that was maintained for 24 to 34 h until complete consumption. *N. ceranae* spores were obtained from previously infected honeybees and purified according to Vidau *et al.* (81). Chronic exposure to insecticides was performed by feeding non-infected honeybees with sugar syrup supplemented with either 0.25 or 1.0 µg kg⁻¹ fipronil, 650 µg kg⁻¹ coumaphos, 1.7 µg kg⁻¹ thiamethoxam, or 3.5 µg kg⁻¹ imidacloprid. Insecticide stock solutions were prepared in DMSO, leading to a final concentration of 0.05% DMSO (v/v) in the syrup. Non-intoxicated bees (*i.e.* untreated control and infected honeybees) were fed syrup supplemented with 0.05% DMSO. According to the acute lethal dose (LD50) (62), the daily mean consumption of insecticides was below 1:194th of the LD50 for coumaphos, thiamethoxam, and imidacloprid, and below 1:550th and 1:150th of the LD50 for low and high fipronil exposure, respectively (Fig. S1).

Gut sampling and DNA purification

Honeybees were sacrificed 18 d after the initiation of experiments, but 15 d for the highest dose of fipronil due to high mortality, along with the corresponding controls. Regarding each replicate, the hindguts of 7 honeybees were dissected on ice, pooled, flash frozen in liquid nitrogen, and stored at -80°C. The presence or absence of *N. ceranae* spores was checked by microscopy (×400) before pooling. Samples were homogenized in 800 µL of RLT buffer (Qiagen) and ground using a microtube pestle. The mixture was transferred to a tube containing 0.150 g of 0.1-mm silica beads and cells were mechanically disrupted by a set of 6 bead-beating pulses of 30 s at 30 Hz, with 30-s interruptions (28). After 3 series of thermal shocks at 65°C and in liquid nitrogen, samples were centrifuged at 8,000×g for 5 min and 200 to 300 µL of the clear supernatant was transferred to a new tube, carefully avoiding both pelleted and floating slimy materials. One volume of ethanol containing 1% (v/v) of 2-mercaptoethanol was added to the sample, which was then transferred to a DNeasy spin column for DNA purification (DNeasy Mini Kit, Qiagen). DNA concentrations were quantified using the bisBenzimide DNA Quantitation Kit, Fluorescence Assay (Sigma-Aldrich).

PCR quantification of bacterial taxa abundance

The theoretical specificity of primer pairs for the 16S rRNA gene was evaluated using the Testprime 1.0 tool (46) to estimate the coverage of bacterial taxa from the SILVA 132 database. Coverage was also estimated on honeybee microbiota 16S rRNA sequences from previous experiments (2, 21, 23, 59, 61, 83), requiring

sequence assignment and selection. Taxonomic assignment was performed through a Blast search against the SILVA 132 database, keeping only sequences with >98% identity. Data reported by Corby-Harris *et al.* (21) were used to assign *Lactobacillus* sp. sequences to the Firm-4 and Firm-5 clades. Only sequences covering the whole amplicon were considered for each primer set, while sequences with ambiguous positions in the priming regions were removed. Primer sets have been experimentally tested for their ability to fulfill qPCR quality requirements (12) on honeybee gut material by checking their specificity (by gel electrophoresis), linear dynamic range, efficiency, and amplification reproducibility using 1:4 serial dilutions of gut DNA extracts.

qPCR was performed on a CFX96 Real-Time System Thermocycler (BioRad) in 96-well plates (Eurogentec RT-PL96-MQ) in a final volume of 20 µL containing 10 µL of 2X Absolute Blue qPCR SYBR Green Mix (Thermo Scientific), 10 pmol of each primer (Table 1), and 2 to 24 ng of total genomic DNA. The PCR program consisted of an initial step at 94°C for 10 min, and 40 cycles comprising 94°C for 25 s, 25 s of annealing at 53°C, and 25 s of elongation at 72°C. Treated and untreated samples from an identical replicate were always deposited on the same plate and all plates were duplicated. The specificity of the reactions was assessed by analyzing the melting curves of the amplified products (Bio-Rad CFX Manager software). The Cycle Quantification (C_Q) values obtained were proportional to the log of input DNA amounts.

In contrast to previous studies, in which qPCR data were normalized to total DNA content or sample weights, we selected the total bacterial content as a reference. Therefore, we avoided biases due to non-bacterial DNA, *e.g.* DNA from *N. ceranae* or the honeybee, or those due to the loss of material following dissection or defecation. The total bacterial DNA content was estimated by the mean of C_Q values obtained with both the 341F/534R and BAC338F/BAC805R primer pairs (Table 1). It is important to note that the two pairs gave very similar C_Q values (correlation coefficient of 0.985; *n*=197). In each taxonomic group, the corresponding C_Q was subtracted from the mean reference C_Q of the same sample. Normalized C_Q were then compared pairwise. The log₂ of the ratio of a taxonomic group under the treated condition relative to the untreated condition was obtained by subtracting the normalized C_Q of the control from the normalized C_Q of the treated sample from the same colony and sampling day. Statistical analyses were performed using PAST software (37).

Results

Honeybees were exposed to the parasite *N. ceranae* or to low doses of the four neurotoxic insecticides. Daily sugar consumption was constant throughout the experiment and not significantly different between groups (Fig. S1). Survival analyses showed a significant decrease in honeybee survival for both winter and summer honeybees infected with *N. ceranae* or exposed to the highest dose of fipronil (Fig. S2). The four other insecticide treatments did not significantly reduce honeybee survival.

In order to assess the abundance of major bacterial taxa, published primer pairs targeting the 16S rRNA gene were checked for qPCR quality (Table 1). Ten primer pairs were eventually selected that target: total bacteria (two pairs), *Alphaproteobacteria*, *Bifidobacteriaceae* (almost exclusively represented by *Bifidobacterium* spp. in the honeybee gut), *Lactobacillus* spp. with one pair recognizing all *Lactobacillus* spp. (including Firm-4, Firm-5, and *L. kunkeei* clades) and another the *Lactobacillus* Firm-5 clade only, *Gammaproteobacteria*, *G. apicola* species, and *S. alvi* species (two pairs, including one recognizing *Neisseriaceae* that are almost exclusively represented by *S. alvi* in the honeybee microbiota). Principal component and ANOVA2 analyses suggested that season

Table 1. Primer pairs tested for the QPCR quantification of the honeybee microbiota major bacterial taxa.

Primer pairs and sequences (5'→3')	Targeted group and % coverage in SILVA database (in honeybee microbiota)	Amplicon size (bp)	Linearity (r ²) Efficiency (E)	validated DNA range ^a and C _Q range	Reference
Bacteria					
341F CCTACGGGAGGCAGCAG *	<i>Bacteria</i> 81.7	169–195	r ² = 0,999 E = 0,92	50 pg to 50 ng C _Q 10 to 22	11
534R ATTACCGCGGCTGCTGGCA *					
BAC338F ACTCCTACGGGAGGCAG *	<i>Bacteria</i> 77.4	443–469	r ² = 0,999 E = 0,85	50 pg to 10 ng ^b C _Q 13.5 to 23	86
BAC805R GACTACAGGGTATCTAATCC *					
BACT1369F CGGTGAATACGTTCYCGG	<i>Bacteria</i> 25.4	133–146	r ² = 0,999 E = 0,84	50 pg to 50 ng C _Q 11 to 23	77
PROK1492R GGWTACCTTGTTACGACTT					
926F AAACCTCAAAGAATTGACGG	<i>Bacteria</i> 78.4	172–179	r ² = 0,999 E = 0,87	50 pg to 50 ng C _Q 11 to 22	6
1062R CTCACRRACAGAGCTGAC					
Bifidobacterium spp.					
Bifi-F2 TCGCGTCYGGTGTGAAAG *	<i>Bifidobacteriaceae</i> 89.3	243	r ² = 0,999 E = 0,78	50 pg to 50 ng C _Q 14.5 to 25	70
Bifi-R2 CCACATCCAGCRTCCAC *	<i>Bifidobacterium</i> spp. 89.0 (96.1, n=51)				
Bifi-F1 CAAGCGAGAGTGAGTGATACC	<i>Bifidobacterium</i> spp. 2.5 (88.7, n=53)	165	r ² = 0,999 E = 0,90	50 pg to 50 ng C _Q 14.5 to 25.5	53
Bifi-R1 GCCGATCCACCGTTAAGC					
Act920F3 TACGGCCGCAAGGCTA	<i>Actinobacteria</i> 86.7	299–307	r ² = 0,999 E = 0,67	not validated	6
Act1200R TCRTCCCCACCTTCTCTCCG	<i>Bifidobacteriaceae</i> 84.8 <i>Bifidobacterium</i> spp. 86.5 (94.2, n=52)				
Lactobacillus spp.					
Lact-F1 TAACGCATTAAGCACTCC *	<i>Lactobacillus</i> 3 (28.6, n=1,705)	270	r ² = 0,998 E = 0,89	50 pg to 50 ng C _Q 13.5 to 24.5	53
Lact-R1 GCTGGCAACTAATAATAAGG *	<i>L. kunkei</i> (0, n=875) Firm-4 (0, n=88) Firm-5 (97.2, n=424)				
Lact-F2 AGCAGTAGGGAATCTTCCA *	<i>Lactobacillus</i> 87.2 (95.7, n=1,690)	341	r ² = 0,997 E = 1,08	50 pg to 10 ng ^b C _Q 14 to 23	70
Lact-R2 CACCGCTACACATGGAG *	<i>L. kunkei</i> (96.8, n=875) Firm-4 (97.7, n=88) Firm-5 (91.7, n=424)				
928F-Firm TGAAACTYAAAGGAATTGACG	<i>Firmicutes</i> 73.7	149–157	r ² = 0,998 E = 0,83	50 pg to 10 ng ^b C _Q 14.5 to 23.5	6
1040FirmR ACCATGCACCACCTGTC	<i>Lactobacillales</i> 73.3 <i>Lactobacillus</i> (43.6) (69.1, n=1,705) <i>L. kunkei</i> (97.7-8, n=875) Firm-4 (97.7, n=88) Firm-5 (0.5, n=424) <i>Mollicutes</i> 39.8				
Alphaproteobacteria					
a682F CDAGTGTAGAGGTGAAATT *	<i>Alphaproteobacteria</i> 63.6	245–250	r ² = 0,998 E = 0,91	50 pg to 50 ng C _Q 14.5 to 25	6
908aR CCCCCTCAATTCCTTTGAGTT *	<i>Bartonella</i> 25.4 (89.6, n=77) <i>Acetobacteraceae</i> 0.8 (10.0, n=201)				
Gammaproteobacteria					
1080yF TCGTCAGCTCGTGTGTGTA *	<i>Gammaproteobacteria</i> 61.2	146–160	r ² = 0,999 E = 0,90	50 pg to 50 ng C _Q 14.5 to 25	6
γ1202R CGTAAGGGCCATGATG *	<i>Gilliamella</i> 96.9 (97.1, n=516) <i>Snodgrassella</i> 0 (0, n=342) <i>Enterobacteriales</i> 94.0				
Snodgrassella alvi					
Neiss-F AAGCGGTGGATGATGTGG *	<i>Neisseriaceae</i> 79.4	194–199	r ² = 0,999 E = 0,90	50 pg to 50 ng C _Q 13.5 to 23.5	53
Neiss-R TGATGGCAACTAATGACAAGG *	<i>Snodgrassella</i> 86.4 (95.1, n=344) <i>Snodgrassella alvi</i> 100 (95.6, n=226)				
Beta-1009-qtF CTTAGAGATAGGAGAGTG *	<i>Snodgrassella</i> 77.3 (93.0, n=344)	127–128	r ² = 0,999 E = 0,87	50 pg to 50 ng C _Q 15 to 24	60
Beta-1115-qtR AATGATGGCAACTAATGACAA *	<i>Snodgrassella alvi</i> 76.9 (94.2, n=226)				
Snod-F GGAATTTCTTAGAGATAGGAAAGTG	<i>Snodgrassella</i> 11.4 (0, n=344)	136	r ² = 0,999 E = 0,86	50 pg to 50 ng C _Q 15 to 26	85
Snod-R TTAATGATGGCAACTAATGACAA	<i>Snodgrassella alvi</i> 23.1 (0, n=226)				
Gilliamella apicola					
G1-459-qtF GTATCTAATAGGTGCATCAATT *	<i>Gilliamella</i> 23.0 (71.6, n=232)	210	r ² = 0,999 E = 0,91	50 pg to 50 ng C _Q 14.5 to 25.5	60
G1-648-qtR TCCTCTACAATACTCTAGTT *	<i>Gilliamella apicola</i> 22.9 (80.7, n=109)				
Gill-F CCTTTGTTGCCATCGATTAGG	<i>Gilliamella</i> 28.0 (1.7, n=355)	249	r ² = 0,999 E = 0,83	50 pg to 50 ng C _Q 16.5 to 28	85
Gill-R GACATTTGATTACGATTACTAGC	<i>Gilliamella apicola</i> 37.5 (0.9, n=233)				
Past-F TTGTTGCCAGCGATTAGG	<i>Gilliamella apicola</i> 4.2 (2.1, n=233)	243	r ² = 0,998 E = 0,75	200 pg to 50 ng ^c C _Q 18 to 27.5	53
Past-R ATTCTGATTACGATTACTAGC	<i>Frischella perrara</i> 100 (94.9, n=59)				
Bacteroidetes					
798cfbF CRAACAGGATTAGATACCCT	<i>Bacteroidia</i> 86.2 (100, n=44)	202–208	r ² = 0,998 E = 0,90	50 pg to 50 ng C _Q 21.5 to 33 ^d	6
cfb967R GGTAAGGTTCTCGCGTAT	<i>Spirochaetaceae</i> 42.8				

^a total DNA from the honeybee gut^b not quantitative for higher DNA concentrations^c not quantitative for lower DNA concentrations^d high C_Q precluded any analyses

Selected pairs are indicated by an *.

was the main factor explaining differences in the abundance of these taxa (Fig. S3). More than 79% of variance was explained by one principal component that appeared to be mainly linked to season and partly to treatments, particularly to *N. ceranae* infection.

The abundance of the major bacterial taxa was initially compared between summer and winter in untreated honeybees. *Bifidobacterium* spp., *Lactobacillus* spp., *Lactobacillus* Firm-5, and *Alphaproteobacteria* were significantly more abundant in winter, while *G. apicola* and *Gammaproteobacteria* were both more abundant in summer (Fig. 1). The core microbiota of the gut was subjected to global seasonal variations.

The abundance of bacterial groups in honeybees chronically exposed to fipronil, imidacloprid, thiamethoxam, or coumaphos was then compared to the untreated control (Fig. 2). Sublethal exposure to all insecticides induced a significant decrease in the relative abundance of *Lactobacillus* spp. in winter and summer honeybees, except for the treatment with the lower dose of fipronil (0.25 $\mu\text{g kg}^{-1}$) for which the decrease was not significant in summer. The highest and lethal dose of fipronil (1 $\mu\text{g kg}^{-1}$) did not significantly change *Lactobacillus* spp. abundance. Similar results were obtained with all *Lactobacillus* spp. or the Firm-5 clade only. A significant decrease in the abundance of *Bifidobacterium* spp. was observed in response to imidacloprid and thiamethoxam in summer honeybees and to coumaphos and the low dose of fipronil (0.25 $\mu\text{g kg}^{-1}$) in winter honeybees. The abundance of *Bifidobacterium* spp. was not affected by the lethal dose of fipronil. Chronic exposure to insecticides induced other significant variations, but in winter honeybees only, with a decrease in the relative abundance

Fig. 1. Comparison of the abundance of major bacterial taxa in the gut microbiota between summer and winter honeybees: *Alphaproteobacteria* (α -P), *Bifidobacterium* spp. (Bifi), *Lactobacillus* spp. (Lact), *Lactobacillus* Firm-5 clade only (Firm5), *Gammaproteobacteria* (γ -P), *Gilliamella apicola* (G.api), and *Snodgrassella alvi* (S.alvi1 and S.alvi2 using the primer pairs Neiss-F/Neiss-R and Beta-1009-qtF/Beta-1115-qtR, respectively). Interior worker honeybees were maintained for 18 d in cages (control conditions) before 16S rRNA gene quantification by qPCR in the hindgut. The y-axis represents the mean log₂ of fold changes (FC) in taxon abundance between summer and winter workers. Positive and negative values denote higher and lower abundance, respectively, in summer bees than in winter bees. Data were gathered from five independent colony replicates. Bars represent 95% confidence intervals. Stars indicate significant differences ($P < 0.05$) by a paired Wilcoxon signed-rank test.

of *Alphaproteobacteria* in response to thiamethoxam (Fig. 2A), an increase in *G. apicola* in response to thiamethoxam and 1 $\mu\text{g kg}^{-1}$ fipronil (Fig. 2F), and an increase in *S. alvi* in response to 1 $\mu\text{g kg}^{-1}$ fipronil using *S. alvi*-, but not *Neisseriaceae*-specific primers (Fig. 2G and H). Collectively, these results showed the similar effects of sublethal doses of insecticides that were more significant in winter honeybees.

N. ceranae infection induced variations in the proportion of all bacterial groups tested (Fig. 2). The abundance of *Alphaproteobacteria*, *Bifidobacterium* spp., and *Lactobacillus* spp. was significantly decreased by the parasite regardless of the season. In contrast, *Gammaproteobacteria*, *G. apicola*, and *S. alvi* significantly increased in winter honeybees only. However, one phylotype identified as *S. alvi* was less abundant in infected bees, as shown by Denaturing Gradient Gel Electrophoresis (Fig. S4), suggesting the complex rebalancing of *S. alvi* strains in infected honeybees.

Discussion

Seasonality of the honeybee microbiota and of stressor-induced variations

The abundance of the main bacterial groups of the gut microbiome differed in summer and winter honeybees (Fig. 1). The higher abundance of *G. apicola* during the beekeeping season was already suggested by Ludvigsen *et al.* (57). In contrast to previous studies (21, 57) seasonal dynamics were also observed for *Alphaproteobacteria*, *Bifidobacterium* spp., and *Lactobacillus* spp., which were less abundant in summer. Since the honeybee gut microbiota is influenced by the environmental landscape (42), it is logically influenced by season in temperate areas. Genomic data suggested that *G. apicola* participates in digestion through the breakdown of complex carbohydrates, such as the pectin contained in pollen (29, 52). The increased abundance of *G. apicola* in summer may be related to increases in these resources.

Responses to stressors also differed according to season (Fig. 2), with more significant results being observed in winter. The mortality of honeybees exposed to acute neonicotinoid treatments was previously reported to be higher in winter than in summer (7). We herein demonstrated that the microbiota, and, thus, the holobiont were more sensitive in winter. The lack of significance in summer bees may be related to a higher heterogeneity in microbial communities, reflected by the greater variance of data (Fig. 2 and S3) and higher α -diversity (57). These findings may be explained by the greater diversity of resources in summer (25, 42). Moreover, short-living summer bees have a more diverse physiology, engaging in the age-dependent division of labor with successive changes in the endocrine system (73), and microbiota richness and abundance have been suggested to differ with ontogenetic stages and behavioral tasks (40, 43). In contrast, in February, aged overwintering bees are more homogeneous, clustering inside the hive and feeding on less diverse reserves.

Alterations in the honeybee microbiota by the intestinal parasite *N. ceranae*

Pathogens may alter the composition of the honeybee microbiota (30, 35), and the present results revealed that all

Fig. 2. Abundance of major taxa in the gut microbiota of pesticide-exposed or *N. ceranae*-infected honeybees and untreated honeybees. Interior worker honeybees were submitted to a chronic exposure to 650 $\mu\text{g kg}^{-1}$ coumaphos (COM), 1.7 $\mu\text{g kg}^{-1}$ thiamethoxam (TMX), 3.5 $\mu\text{g kg}^{-1}$ imidacloprid (IMI), 0.25 $\mu\text{g kg}^{-1}$ (FIP¼), or 1 $\mu\text{g kg}^{-1}$ fipronil (FIP1) or to an infection with *N. ceranae* (Ncer). The abundance of bacterial taxa was assessed by qPCR after 15 or 18 d in winter (white bars) and summer (hatched bars) honeybees. The y-axes depict the mean $\log_2$ of fold changes (FC) in abundance between treated and untreated samples. Negative and positive values denote lower and higher abundance, respectively, in response to stressors. Data were gathered from five independent colony replicates. Bars represent 95% confidence intervals. Stars indicate significant differences ($P < 0.05$) by a paired Wilcoxon signed-rank test.

major bacterial taxa were affected by *N. ceranae* (Fig. 2), with a decrease in lactic acid bacteria (LABs), including *Lactobacillus* spp. and *Bifidobacterium* spp., and an increase in *S. alvi* and *G. apicola*. The increase in *S. alvi* and *G. apicola* may have been due to a rebalance in the microbiota by occupying niches left vacant by LABs. The results of the DGGE analysis suggested that while all *S. alvi* species experienced an increase in abundance, the content of a specific strain decreased. This opposite variation in strain content was previously reported for *G. apicola* in response to an antibiotic treatment, suggesting variations in the ability of strains to cope with stress (66). This important dysbiosis was not observed in *A. mellifera* in previous studies (41, 54, 58). In the Asian honeybee *A. cerana*, Li *et al.* (53) observed a similar decrease in the abundance of *Bifidobacterium* spp., but no significant change in *Lactobacillus* spp..

It is unlikely that *N. ceranae* directly affects gut microbial communities because its development entirely occurs within midgut epithelial cells. However, it may alter honeybee midgut homeostasis. *N. ceranae* induces a degeneration in the midgut epithelium and peritrophic matrix (26, 39, 48) that may alter the activation of the immune system by luminal bacteria (38, 47). The infection also up-regulates the antioxidant system, reduces the production of reactive oxygen species (ROS) (26, 65, 81, 82), and may provoke immunosuppression by inhibiting the production of antimicrobial peptides (AMP) (3, 5, 16, 55). No data are available on the honeybee hindgut. Interestingly, the microsporidian parasite *Paranosema locustae* alters the gut microbiota of *Locusta migratoria*, but also induces acidification and an increase in ROS in the hindgut (74, 78). By affecting the gut ecosystem, possibly through changes in AMPs, ROS, or pH, *N. ceranae* may favor or impair bacterial communities.

Alterations in the honeybee gut microbiota by insecticides

Honeybees were chronically exposed to concentrations of insecticides that correspond to residue levels detected in hive matrices (18, 45, 51, 62), denoting realistic exposure. These treatments led to variations in the honeybee core microbiota taxa, with a general decrease in LABs (Fig. 2). Kakumanu *et al.* (44) also observed a significant decrease in *Lactobacillus* spp. in response to the fungicide chlorothalonil, but a slight increase in *Bifidobacterium* spp. in response to coumaphos. This opposite variation in *Bifidobacterium* spp. suggests the strong influence of the mode of exposure (oral exposure in cages vs an in-hive CheckMite+® treatment) on microbial responses. In contrast to our results, Raymann *et al.* (68) reported that imidacloprid did not significantly affect the microbiota. The treatment with 1.0 µg kg⁻¹ fipronil did not have a stronger effect on LABs than the 0.25 µg kg⁻¹ treatment, but induced an increase in *G. apicola* and *S. alvi* in winter (Fig. 2). This higher insecticide concentration was lethal (Fig. S2) and may have had a strong effect on morbidity in the microbiota because dying honeybees were sampled.

Insecticides may have a direct impact on bacterial growth that strongly varies upon insecticides and bacterial strains (75). Thus, insecticides from different families may not have the same direct impact on gut bacteria. The similar effects observed in response to different insecticides (Fig. 2) may have been due to aspecific changes in the gut homeostasis that

may secondarily affect the microbiota. In the midgut, chronic and oral exposure to insecticides altered the immune system, activated antioxidant defenses, and induced damage in epithelial cells (5, 9, 15, 20, 31, 68). Thus, gut bacterial communities may have to cope with changes in their environment.

Ingested neurotoxic insecticides are absorbed in the gut and diffuse throughout the whole insect body (76), eventually affecting the nervous system (14). Neonicotinoids, such as thiamethoxam and imidacloprid, induce morphological, cytological, and transcriptomic alterations in the brain and disrupt the hormonal balance (20). Thus, besides local effects on the gut, insecticides exert pleiotropic effects that may, in turn, affect the gut and its microbiota, which is considered to be connected to the nervous system in mammals (69). This indirect impact on the gut has been demonstrated based on changes in midgut antioxidant activities following a single topical exposure to fipronil (13).

Dysbiosis may sensitize the honeybee to other stressors

The microbiota protects the host against pathogens by stimulating the immune system and providing competition for niche occupation (36). The honeybee gut microbiota is considered to induce AMP production and the LABs themselves exhibit antibacterial properties (50, 67, 80). The suppression of honeybee gut bacteria by antibiotics increased honeybee vulnerability to *N. ceranae* (54), and some bacteria, including LABs, may conversely reduce parasite proliferation (22, 27). In contrast, feeding supplemented with *S. alvi* increased honeybee sensitivity to the trypanosomal parasite *Lotmaria passim* (72).

The insect microbiota may also comprise strains with the ability to enhance host resistance to insecticides (84). In the moth *Plutella xylostella*, a positive correlation was observed between *Lactobacillales* abundance and fipronil resistance. Moreover, some gut bacteria isolated from insects, including *Lactobacillus* spp., are considered to metabolize insecticides (1, 24). Thus, dysbiosis and a reduction in LABs may sensitize the honeybee to pathogens and pesticides.

The season-dependent gut microbiota may be regarded as a modulator of interactions between the honeybee and its stressors, emphasizing the need to consider the holobiont, *i.e.* the insect and its symbionts, both in summer and winter, when assessing the effects of stressors.

Acknowledgements

The project was funded by the ANR MIXTRESS program (ANR-15-CE34-0004). The authors are thankful to Jérôme Lesobre, Johan Panek, and Laurianne Paris for their support in beekeeping activities and to Clermont Auvergne Métropole for lending the beehive site. The authors declare no conflict of interest.

References

- Almeida, L.G. de, L.A.B. de Moraes, J.R. Trigo, C. Omoto, and F.L. Cônsoli. 2017. The gut microbiota of insecticide-resistant insects houses insecticide-degrading bacteria: a potential source for biotechnological exploitation. *PLoS One* 12:e0174754.
- Anderson, K.E., T.H. Sheehan, B.M. Mott, P. Maes, L. Snyder, M.R. Schwan, A. Walton, B.M. Jones, and V. Corby-Harris. 2013. Microbial ecology of the hive and pollination landscape: Bacterial associates from floral nectar, the alimentary tract and stored food of honey bees (*Apis mellifera*). *PLoS One* 8:e83125.

3. Antúnez, K., R. Martín-Hernández, L. Prieto, A. Meana, P. Zunino, and M. Higes. 2009. Immune suppression in the honey bee (*Apis mellifera*) following infection by *Nosema ceranae* (Microsporidia). *Environ. Microbiol.* 11:2284–2290.
4. Aufauvre, J., D.G. Biron, C. Vidau, R. Fontbonne, M. Roudel, M. Diogon, B. Viguès, L.P. Belzunces, F. Delbac, and N. Blot. 2012. Parasite-insecticide interactions: a case study of *Nosema ceranae* and fipronil synergy on honeybee. *Sci. Rep.* 2:326.
5. Aufauvre, J., B. Misme-Aucouturier, B. Viguès, C. Texier, F. Delbac, and N. Blot. 2014. Transcriptome analyses of the honeybee response to *Nosema ceranae* and insecticides. *PLoS One* 9:e91686.
6. Bacchetti De Gregoris, T., N. Aldred, A.S. Clare, and J.G. Burgess. 2011. Improvement of phylum- and class-specific primers for real-time PCR quantification of bacterial taxa. *J. Microbiol. Methods.* 86:351–356.
7. Baines, D., E. Wilton, A. Pawluk, M. de Gorter, and N. Chomistek. 2017. Neonicotinoids act like endocrine disrupting chemicals in newly-emerged bees and winter bees. *Sci. Rep.* 7:10979.
8. Beyer, M., A. Lenouvel, C. Guignard, M. Eickermann, A. Clermont, F. Kraus, and L. Hoffmann. 2018. Pesticide residue profiles in bee bread and pollen samples and the survival of honeybee colonies—a case study from Luxembourg. *Environ. Sci. Pollut. Res.* <https://doi.org/10.1007/s11356-018-3187-4>.
9. Boncristiani, H., R. Underwood, R. Schwarz, J.D. Evans, J. Pettis, and D. vanEngelsdorp. 2012. Direct effect of acaricides on pathogen loads and gene expression levels in honey bees *Apis mellifera*. *J. Insect Physiol.* 58:613–620.
10. Bonmatin, J.-M., C. Giorio, V. Girolami, *et al.* 2015. Environmental fate and exposure; neonicotinoids and fipronil. *Environ. Sci. Pollut. Res.* 22:35–67.
11. Bru, D., F. Martin-Laurent, and L. Philippot. 2008. Quantification of the detrimental effect of a single primer-template mismatch by real-time PCR using the 16S rRNA gene as an example. *Appl. Environ. Microbiol.* 74:1660–1663.
12. Bustin, S.A., V. Benes, J.A. Garson, *et al.* 2009. The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* 55:611–622.
13. Carvalho, S.M., L.P. Belzunces, G.A. Carvalho, J.-L. Brunet, and A. Badiou-Beneteau. 2013. Enzymatic biomarkers as tools to assess environmental quality: A case study of exposure of the honeybee *Apis mellifera* to insecticides. *Environ. Toxicol. Chem.* 32:2117–2124.
14. Casida, J.E., and K.A. Durkin. 2013. Neuroactive insecticides: targets, selectivity, resistance, and secondary effects. *Annu. Rev. Entomol.* 58:99–117.
15. Catae, A.F., T.C. Roat, M. Pratavieira, A.R. da Silva Menegasso, M.S. Palma, and O. Malaspina. 2018. Exposure to a sublethal concentration of imidacloprid and the side effects on target and nontarget organs of *Apis mellifera* (Hymenoptera, Apidae). *Ecotoxicology* 27:109–121.
16. Chaimanee, V., P. Chantawannakul, Y. Chen, J.D. Evans, and J.S. Pettis. 2012. Differential expression of immune genes of adult honey bee (*Apis mellifera*) after inoculated by *Nosema ceranae*. *J. Insect Physiol.* 58:1090–1095.
17. Charreton, M., A. Decourtye, M. Henry, G. Rodet, J.-C. Sandoz, P. Charvet, and C. Collet. 2015. A locomotor deficit induced by sublethal doses of pyrethroid and neonicotinoid insecticides in the honeybee *Apis mellifera*. *PLoS One* 10:e0144879.
18. Chauzat, M.-P., J.-P. Faucon, A.-C. Martel, J. Lachaize, N. Cougoule, and M. Aubert. 2006. A survey of pesticide residues in pollen loads collected by honey bees in France. *J. Econ. Entomol.* 99:253–262.
19. Chauzat, M.-P., A. Jacques, EPILOBEE Consortium, M. Laurent, S. Bougeard, P. Hendrikx, and M. Ribière-Chabert. 2016. Risk indicators affecting honeybee colony survival in Europe: one year of surveillance. *Apidologie* 47:348–378.
20. Christen, V., F. Mittner, and K. Fent. 2016. Molecular effects of neonicotinoids in honey bees (*Apis mellifera*). *Environ. Sci. Technol.* 50:4071–4081.
21. Corby-Harris, V., P. Maes, and K.E. Anderson. 2014. The bacterial communities associated with honey bee (*Apis mellifera*) foragers. *PLoS One* 9:e95056.
22. Corby-Harris, V., L. Snyder, C.A. Meador, R. Naldo, B. Mott, and K.E. Anderson. 2016. *Parasaccharibacter apium*, gen. nov., sp. nov., improves honey bee (Hymenoptera: Apidae) resistance to *Nosema*. *J. Econ. Entomol.* 109:537–543.
23. Cox-Foster, D.L., S. Conlan, E.C. Holmes, *et al.* 2007. A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* 318:283–287.
24. Daisley, B.A., M. Trinder, T.W. McDowell, S.L. Collins, M.W. Sumarah, and G. Reid. 2018. Microbiota-mediated modulation of organophosphate insecticide toxicity by species-dependent lactobacilli interactions in a *Drosophila melanogaster* insect model. *Appl. Environ. Microbiol.* 84:e02820–17.
25. Danner, N., A. Keller, S. Härtel, and I. Steffan-Dewenter. 2017. Honey bee foraging ecology: Season but not landscape diversity shapes the amount and diversity of collected pollen. *PLoS One* 12:e0183716.
26. Dussaubat, C., J.-L. Brunet, M. Higes, *et al.* 2012. Gut pathology and responses to the microsporidium *Nosema ceranae* in the honey bee *Apis mellifera*. *PLoS One* 7:e37017.
27. El Khoury, S., A. Rousseau, A. Lecoeur, B. Cheaib, S. Bouslama, P.-L. Mercier, V. Demey, M. Castex, P. Giovenazzo, and N. Derome. 2018. Deleterious interaction between honeybees (*Apis mellifera*) and its microsporidian intracellular parasite *Nosema ceranae* was mitigated by administrating either endogenous or allochthonous gut microbiota strains. *Front. Ecol. Evol.* 6:58.
28. Engel, P., R.R. James, R. Koga, W.K. Kwong, Q.S. McFrederick, and N.A. Moran. 2013. Standard methods for research on *Apis mellifera* gut symbionts. *J. Apic. Res.* 52:1–24.
29. Engel, P., and N.A. Moran. 2013. Functional and evolutionary insights into the simple yet specific gut microbiota of the honey bee from metagenomic analysis. *Gut Microbes.* 4:60–65.
30. Erban, T., O. Ledvinka, M. Kamler, B. Hortova, M. Nesvorna, J. Tyl, D. Titera, M. Markovic, and J. Hubert. 2017. Bacterial community associated with worker honeybees (*Apis mellifera*) affected by European foulbrood. *PeerJ* 5:e3816.
31. Garrido, P.M., M.P. Porrini, K. Antúnez, B. Branchiccela, G.M.A. Martínez-Noël, P. Zunino, G. Salerno, M.J. Eguaras, and E. Ieno. 2016. Sublethal effects of acaricides and *Nosema ceranae* infection on immune related gene expression in honeybees. *Vet. Res.* 47:51.
32. Godfray, H.C.J., T. Blacquiere, L.M. Field, R.S. Hails, G. Petrokofsky, S.G. Potts, N.E. Raine, A.J. Vanbergen, and A.R. McLean. 2014. A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proc. R. Soc. B.* 281:20140558.
33. Godfray, H.C.J., T. Blacquiere, L.M. Field, R.S. Hails, S.G. Potts, N.E. Raine, A.J. Vanbergen, and A.R. McLean. 2015. A restatement of recent advances in the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proc. R. Soc. B.* 282:20151821.
34. Goulson, D., E. Nicholls, C. Botias, and E.L. Rotheray. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 347:1255957–1255957.
35. Guo, J., J. Wu, Y. Chen, J.D. Evans, R. Dai, W. Luo, and J. Li. 2015. Characterization of gut bacteria at different developmental stages of Asian honey bees, *Apis cerana*. *J. Invertebr. Pathol.* 127:110–114.
36. Hamdi, C., A. Balloi, J. Essanaa, *et al.* 2011. Gut microbiome dysbiosis and honeybee health. *J. Appl. Entomol.* 135:524–533.
37. Hammer, O., D.A.T. Harper, and P.D. Ryan. 2001. PAST: Paleontological Statistics software package for education and data analysis. *Palaeontol. Electronica* 4:9.
38. Hegedus, D., M. Erlandson, C. Gillott, and U. Toprak. 2009. New insights into peritrophic matrix synthesis, architecture, and function. *Annu. Rev. Entomol.* 54:285–302.
39. Higes, M., A. Meana, C. Bartolomé, C. Botías, and R. Martín-Hernández. 2013. *Nosema ceranae* (Microsporidia), a controversial 21st century honey bee pathogen: *N. ceranae* an emergent pathogen for beekeeping. *Environ. Microbiol. Rep.* 5:17–29.
40. Hroncova, Z., J. Havlik, J. Killer, *et al.* 2015. Variation in honey bee gut microbial diversity affected by ontogenetic stage, age and geographic location. *PLoS One* 10:e0118707.
41. Hubert, J., M. Bicianova, O. Ledvinka, M. Kamler, P.J. Lester, M. Nesvorna, J. Kopecky, and T. Erban. 2017. Changes in the bacteriome of honey bees associated with the parasite *Varroa destructor*, and pathogens *Nosema* and *Lotmaria passim*. *Microb. Ecol.* 73:685–698.
42. Jones, J.C., C. Fruciano, F. Hildebrand, H.A. Toufalilia, N.J. Balfour, P. Bork, P. Engel, F.L. Ratnieks, and W.O. Hughes. 2018. Gut microbiota composition is associated with environmental landscape in honey bees. *Ecol. Evol.* 8:441–451.
43. Jones, J.C., C. Fruciano, J. Marchant, F. Hildebrand, S. Forslund, P. Bork, P. Engel, and W.O.H. Hughes. 2018. The gut microbiome is associated with behavioural task in honey bees. *Insectes Soc.* 65:419–429.

44. Kakumanu, M.L., A.M. Reeves, T.D. Anderson, R.R. Rodrigues, and M.A. Williams. 2016. Honey bee gut microbiome is altered by in-hive pesticide exposures. *Front. Microbiol.* 7:1255.
45. Kasiotis, K.M., C. Anagnostopoulos, P. Anastasiadou, and K. Machera. 2014. Pesticide residues in honeybees, honey and bee pollen by LC–MS/MS screening: Reported death incidents in honeybees. *Sci. Total Environ.* 485–486:633–642.
46. Klindworth, A., E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, and F.O. Glöckner. 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* 41:e1.
47. Kuraishi, T., O. Binggeli, O. Oputa, N. Buchon, and B. Lemaitre. 2011. Genetic evidence for a protective role of the peritrophic matrix against intestinal bacterial infection in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. U.S.A.* 108:15966–15971.
48. Kurze, C., Y. Le Conte, P. Kryger, O. Lewkowski, T. Müller, and R.F.A. Moritz. 2018. Infection dynamics of *Nosema ceranae* in honey bee midgut and host cell apoptosis. *J. Invertebr. Pathol.* 154:1–4.
49. Kwong, W.K., and N.A. Moran. 2016. Gut microbial communities of social bees. *Nat. Rev. Microbiol.* 14:374–384.
50. Kwong, W.K., A.L. Mancenido, and N.A. Moran. 2017. Immune system stimulation by the native gut microbiota of honey bees. *R. Soc. Open Sci.* 4:170003.
51. Lambert, O., M. Piroux, S. Puyo, C. Thorin, M. L'Hostis, L. Wiest, A. Buleté, F. Delbac, and H. Pouliquen. 2013. Widespread occurrence of chemical residues in beehive matrices from apiaries located in different landscapes of western France. *PLoS One* 8:e67007.
52. Lee, F.J., D.B. Rusch, F.J. Stewart, H.R. Mattila, and I.L.G. Newton. 2015. Saccharide breakdown and fermentation by the honey bee gut microbiome: Fermentation by honey bee gut microbes. *Environ. Microbiol.* 17:796–815.
53. Li, J., H. Qin, J. Wu, B.M. Sadd, X. Wang, J.D. Evans, W. Peng, and Y. Chen. 2012. The prevalence of parasites and pathogens in Asian honeybees *Apis cerana* in China. *PLoS One* 7:e47955.
54. Li, J.H., J.D. Evans, W.F. Li, Y.Z. Zhao, G. DeGrandi-Hoffman, S.K. Huang, Z.G. Li, M. Hamilton, and Y.P. Chen. 2017. New evidence showing that the destruction of gut bacteria by antibiotic treatment could increase the honey bee's vulnerability to *Nosema* infection. *PLoS One* 12:e0187505.
55. Li, W., J.D. Evans, J. Li, S. Su, M. Hamilton, and Y. Chen. 2017. Spore load and immune response of honey bees naturally infected by *Nosema ceranae*. *Parasitol. Res.* 116:3265–3274.
56. Lu, K., R. Mahbub, and J.G. Fox. 2015. Xenobiotics: Interaction with the intestinal microflora. *ILAR J.* 56:218–227.
57. Ludvigsen, J., A. Rangberg, E. Avershina, M. Sekelja, C. Kreibich, G. Amdam, and K. Rudi. 2015. Shifts in the midgut/pyloric microbiota composition within a honey bee apiary throughout a season. *Microbes Environ.* 30:235–244.
58. Maes, P.W., P.A.P. Rodrigues, R. Oliver, B.M. Mott, and K.E. Anderson. 2016. Diet-related gut bacterial dysbiosis correlates with impaired development, increased mortality and *Nosema disease* in the honeybee (*Apis mellifera*). *Mol. Ecol.* 25:5439–5450.
59. Martinson, V.G., B.N. Danforth, R.L. Minckley, O. Rueppell, S. Tingek, and N.A. Moran. 2011. A simple and distinctive microbiota associated with honey bees and bumble bees: the microbiota of honey bees and bumble bees. *Mol. Ecol.* 20:619–628.
60. Martinson, V.G., J. Moy, and N.A. Moran. 2012. Establishment of characteristic gut bacteria during development of the honeybee worker. *Appl. Environ. Microbiol.* 78:2830–2840.
61. Moran, N.A., A.K. Hansen, J.E. Powell, and Z.L. Sabree. 2012. Distinctive gut microbiota of honey bees assessed using deep sampling from individual worker bees. *PLoS One* 7:e36393.
62. Mullin, C.A., M. Frazier, J.L. Frazier, S. Ashcraft, R. Simonds, D. vanEngelsdorp, and J.S. Pettis. 2010. High levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *PLoS One* 5:e9754.
63. Oliveira, R.A., T.C. Roat, S.M. Carvalho, and O. Malaspina. 2014. Side-effects of thiamethoxam on the brain and midgut of the Africanized honeybee *Apis mellifera* (Hymenoptera: Apidae). *Environ. Toxicol.* 29:1122–1133.
64. Panek, J., L. Paris, D. Roriz, A. Mone, A. Dubuffet, F. Delbac, M. Diogon, and H. El Alaoui. 2018. Impact of the microsporidian *Nosema ceranae* on the gut epithelium renewal of the honeybee, *Apis mellifera*. *J. Invertebr. Pathol.* <https://doi.org/10.1016/j.jip.2018.09.007>.
65. Paris, L., M. Roussel, B. Pereira, F. Delbac, and M. Diogon. 2017. Disruption of oxidative balance in the gut of the western honeybee *Apis mellifera* exposed to the intracellular parasite *Nosema ceranae* and to the insecticide fipronil. *Microb. Biotechnol.* 10:1702–1717.
66. Raymann, K., L.-M. Bobay, and N.A. Moran. 2018. Antibiotics reduce genetic diversity of core species in the honeybee gut microbiome. *Mol. Ecol.* 27: 2057–2066.
67. Raymann, K., and N.A. Moran. 2018. The role of the gut microbiome in health and disease of adult honey bee workers. *Curr. Opin. Chem. Biol.* 26:97–104.
68. Raymann, K., E.V.S. Motta, C. Girard, I.M. Riddington, J.A. Dinser, and N.A. Moran. 2018. Imidacloprid decreases honey bee survival but does not affect the gut microbiome. *Appl. Environ. Microbiol.* 84:e00545-18.
69. Rhee, S.H., C. Pothoulakis, and E.A. Mayer. 2009. Principles and clinical implications of the brain–gut–enteric microbiota axis. *Nat. Rev. Gastroenterol. Hepatol.* 6:306–314.
70. Rinttilä, T., A. Kassinen, E. Malinen, L. Krogus, and A. Palva. 2004. Development of an extensive set of 16S rDNA-targeted primers for quantification of pathogenic and indigenous bacteria in faecal samples by real-time PCR. *J. Appl. Microbiol.* 97:1166–1177.
71. Ryu, J.-H., S.-H. Kim, H.-Y. Lee, J.Y. Bai, Y.-D. Nam, J.-W. Bae, D.G. Lee, S.C. Shin, E.-M. Ha, and W.-J. Lee. 2008. Innate immune homeostasis by the homeobox gene caudal and commensal-gut mutualism in *Drosophila*. *Science* 319:777–782.
72. Schwarz, R.S., N.A. Moran, and J.D. Evans. 2016. Early gut colonizers shape parasite susceptibility and microbiota composition in honey bee workers. *Proc. Natl. Acad. Sci. U.S.A.* 113:9345–9350.
73. Seeley, T.D. 1985. *Honeybee Ecology: A Study of Adaptation in Social Life*. Princeton University Press, Princeton, New Jersey.
74. Shi, W., Y. Guo, C. Xu, S. Tan, J. Miao, Y. Feng, H. Zhao, R.J. St. Leger, and W. Fang. 2014. Unveiling the mechanism by which microsporidian parasites prevent locust swarm behavior. *Proc. Natl. Acad. Sci. U.S.A.* 111:1343–1348.
75. Staley, Z.R., V.J. Harwood, and J.R. Rohr. 2015. A synthesis of the effects of pesticides on microbial persistence in aquatic ecosystems. *Crit. Rev. Toxicol.* 45:813–836.
76. Suchail, S., G.D. Sousa, R. Rahmani, and L.P. Belzunces. 2004. *In vivo* distribution and metabolism of 14C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Manage. Sci.* 60:1056–1062.
77. Suzuki, M.T., L.T. Taylor, and E.F. Delong. 2000. Quantitative analysis of small-subunit rRNA genes in mixed microbial populations via 5'-nuclease assays. *Appl. Environ. Microbiol.* 66:4605–4614.
78. Tan, S., K. Zhang, H. Chen, Y. Ge, R. Ji, and W. Shi. 2015. The mechanism for microsporidian parasite suppression of the hindgut bacteria of the migratory locust *Locusta migratoria manilensis*. *Sci. Rep.* 5:17365.
79. vanEngelsdorp, D., and M.D. Meixner. 2010. A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *J. Invertebr. Pathol.* 103:S80–S95.
80. Vázquez, A., E. Forsgren, I. Fries, R.J. Paxton, E. Flaberg, L. Szekely, and T.C. Olofsson. 2012. Symbionts as major modulators of insect health: lactic acid bacteria and honeybees. *PLoS One* 7:e33188.
81. Vidau, C., M. Diogon, J. Aufaivre, et al. 2011. Exposure to sublethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*. *PLoS One* 6:e21550.
82. Vidau, C., J. Panek, C. Texier, D.G. Biron, L.P. Belzunces, M. Le Gall, C. Broussard, F. Delbac, and H. El Alaoui. 2014. Differential proteomic analysis of midguts from *Nosema ceranae*-infected honeybees reveals manipulation of key host functions. *J. Invertebr. Pathol.* 121:89–96.
83. Vojvodic, S., S.M. Rehan, and K.E. Anderson. 2013. Microbial gut diversity of Africanized and European honey bee larval instars. *PLoS One* 8:e72106.
84. Xia, X., B. Sun, G.M. Gurr, L. Vasseur, M. Xue, and M. You. 2018. Gut microbiota mediate insecticide resistance in the Diamondback Moth, *Plutella xylostella* (L.). *Front. Microbiol.* 9:25.
85. Xu, L.-L., J. Wu, J. Guo, and J.-L. Li. 2014. Dynamic variation of symbionts in bumblebees during hosts growth and development. *Sci. Agric. Sin.* 47:2030–2037.
86. Yu, Y., C. Lee, J. Kim, and S. Hwang. 2005. Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. *Biotechnol. Bioeng.* 89:670–679.