THREE DIMENSIONAL OBSERVATION OF CELLULAR MICROSTRUCTURES WITH A NOVEL FLUORESCENT OPTICAL TOMOGRAPHY

T. Matsumoto*, M. Ogawa*, F. Wake*, K. Nagayama*, and Y. Ishino**

*Biomechanics Laboratory and **Energy Conversion Engineering Laboratory,
Department of Mechanical Engineering, Nagoya Institute of Technology, Nagoya, Japan

takeo@nitech.ac.jp

Introduction

Detailed knowledge on the three dimensional (3D) microstructure of the cells is crucial in cell biomechanics. The 3D morphology of intracellular structure has been reconstructed with a stack of images obtained from a confocal laser scanning microscope (CLSM). However, the resolution of the images in the direction of the optical axis, *i.e.*, the direction of cell thickness, is much lower than that in the direction perpendicular to it. To overcome this problem, we have been developing a cell rotation system (CRS) in which specimen cell is rotated under a microscope around the axis perpendicular to the optical axis of the microscope to observe the intracellular microstructure precisely from their images obtained at various rotation angles. In this study, we observed fluorescent images of cell nuclei and actin filaments (AF) from various angles and reconstruct their 3D images with a computed tomography algorithm.

Materials and Methods

Cell rotation system: The outline of the CRS is shown in Figure 1. A specimen cell was held with a glass micropipette by aspirating its surface gently under a microscope and the pipette was rotated around the axis perpendicular to the optical axis with a stepping motor via a belt and pulley mechanism. The pipette tip position was coarsely adjusted on the rotation axis with two micrometer heads that was aligned to bend the axis of pipette holder in two orthogonal directions (Figure 2). Displacement of the pipette tip position from the rotation axis during rotation was compensated with two sets of piezo mechanism.

Observation of cell microstructures: Cultured porcine aortic smooth muscle cells (SMCs) were detached from the substrate by

Figure 1 Schematic diagram of the cell rotation device.

Figure 2 Coarse centering mechanism for pipette tip.

trypsinization, stained with Hoechst 33342 for their nuclei and observed with the CRS under a conventional fluorescent microscope at every 4.5° from 0° to 180°. One image was taken at each angle. Their 3D images were reconstructed from these images by applying a distributing back projection method [1]. To observe the nuclear shape of the cells attached to the substrate, cells cultured on a narrow glass strip were also stained with Hoechst 33342. The strip was connected to the CRS and the nuclei were observed similarly. For the observation of AFs, SMCs freshly isolated from rat aortic tissue were stained with Alexa Fluor 546 Phalloidin and observed with the CRS under a CLSM at every 9° from 0° to 180°. Pan-focus fluorescent images of the actin filaments were obtained in a maximum intensity projection process from ~100 confocal images taken at 0.25 μ m intervals at each angle. Their 3D images were reconstructed from the pan-focus images similarly to those of the cell nuclei.

Results and Discussion

Clear images were obtained both for cell nuclei (Figure 3) and AFs (Figure 4). The nucleus of the detached cell had irregular shape and had dents on its surface, while that of the attached cell had flat and smooth surface. This may indicate that cell nuclei are buckled in the cells detached from the substrate. The AFs aligned helically along the long axis of the cell body as suggested in TEM observation [2]. These morphological characteristics would not be clearly observed in 3D images obtained in the conventional method. The present method is useful for quantitative 3D analysis of intracellular structures.

References

- [1] Ishino Y and Oiwa N: Scanless 3D-CT Reconstruction of Turbulent Flame Shape, *J Combustion Soc Jpn*, **46**: 142-152, 2004.
- [2] Small JV: Studies on isolated smooth muscle cells: The contractile apparatus, *J Cell Sci*, **24**: 327-49, 1977.

Figure 3 Reconstructed images of cell nuclei.

Figure 4 Reconstructed images of actin filaments.