【国際セッション】

The influence of cardiac output variability on cerebral blood flow velocity variability during exercise in humans. Shigehiko Ogoh^{1,2}, Mads K. Dalsgaard², Chie Yoshiga², Ellen A. Dawson², Niels H. Secher² and Peter B. Raven¹¹Department of Integrative Physiology, University of North Texas Health Science Center, Fort Worth, TX, USA²Department of Anaesthesia, The Copenhagen Muscle Research Centre, University of Copenhagen, Denmark**Introduction**

Cerebral autoregulation (CA) is important in the regulation of cerebral blood flow (Ogoh *et al.* 2005a, b). In addition, the middle cerebral artery mean blood flow velocity (MCA V_{mean}) is influenced by systemic blood pressure regulation (Ide & Secher, 2000). The increase in MCA V_{mean} reported with exercise appears to depend on the ability to increase cardiac output (Q), as demonstrated in response to β -1 blockade and in patients with cardiac insufficiency or atrial fibrillation (Ide & Secher, 2000). Arterial baroreflex (ABR) control of arterial blood pressure (ABP) operates via two separate reflex arcs, which for the carotid baroreflex (CBR) are defined as the carotid-cardiac or carotid-vasomotor arm. During exercise, the carotid-vasomotor arm of the reflex predominates in the regulation of ABP (Ogoh *et al.* 2003). However, the cardiac ABR responses at the operating point (OP) decrease from rest to maximal exercise (Ogoh *et al.*, 2005c). Therefore, if the contribution of the variability in Q to the beat-to-beat MCA V_{mean} remains stable during exercise, the reduction of cardiac ABR sensitivity at the OP would result in less control of the beat-to-beat MCA V_{mean} . This study evaluated the influence of Q variability on beat-to-beat MCA V_{mean} during exercise with and without cardiac β -1 adrenergic blockade.

Methods

Eight men (22 $\pm$ 1 yrs; mean $\pm$ SE) performed 15 min bouts of moderate (EX130; 105 $\pm$ 11W) and heavy (EX160; 162 $\pm$ 8W) intensity cycling before and after cardio-selective β -1 adrenergic blockade (0.15 mg/kg metoprolol). The beat-to-beat changes in Q (%), determined by Modelflow technology, heart rate (HR), ABP and MCA V_{mean} were also obtained. Transfer function gain, phase shift and coherence between changes in Q and MCA V_{mean} and between mean arterial pressure (MAP) and MCA V_{mean} were calculated. In addition, transfer function gain between MAP and HR was calculated to evaluate dynamic cardiac ABR gain.

Results

At rest, metoprolol did not significantly affect blood gas variables or HR, stroke volume (SV) and Q. However, the MAP and MCA V_{mean} decreased by 7.6% and 13.4%, respectively ($P<0.05$). Without metoprolol, HR, MAP, Q and blood lactate concentration (La) gradually increased from rest to EX160 ($P<0.05$). The MCA V_{mean} and PaCO₂ increased by 25% and 9.6%, respectively ($P<0.05$) from rest to EX130, but the MCA V_{mean} decreased by 11% and PaCO₂ by 5% from EX130 to EX160. After the administration of metoprolol during exercise, all cardiovascular variables increased during exercise, however, the increases in HR, MAP, Q, MCA V_{mean} and La were lower than control values ($P<0.05$). The SV and PaCO₂ were not influenced significantly by cardio-selective β -1 blockade. Both exercise and metoprolol decreased the Low frequency (LF) transfer function gain between %Q and MCA V_{mean} ($P<0.05$). However, during EX130 and EX160 there was no significant difference in the LF transfer function gain with and without after β -1 blockade. During exercise, the transfer function gain between MAP and MCA V_{mean} remained stable and were not different from the resting values with and without metoprolol ($P>0.05$). The transfer function gain between HR and MAP, an index of cardiac ABR responses at the OP, decreased from rest to EX160 ($P<0.05$). In addition, the exercise-induced reduction in cardiac ABR gain was significantly related to a decrease in the transfer function gain between Q and MCA V_{mean} from rest to EX160 ($R=0.942$, $P<0.05$).

Discussion/Conclusion

An important finding of the present investigation was that the contribution of Q variability (cardiac-arterial baroreflex) to MCA V_{mean} variability decreased from rest to heavy exercise. Furthermore, a reduced steady-state Q or MCA V_{mean} induced by β -1 blockade did not alter the influence of Q variability on MCA V_{mean} variability during exercise. However, exercise and the administration of β -1 blockade did not significantly affect the transfer function gain between beat-to-beat MAP and MCA V_{mean} . These data suggest that during exercise the control of systemic blood pressure (i.e., arterial baroreflex control of vasomotion) becomes more important than the cardiac ABR in the beat-to-beat regulation of MCA V_{mean} , even though steady-state Q determined MCA V_{mean} .

References

- Ide K, Secher NH (2000). *Prog Neurobiol* 61, 397-414.
- Ogoh S *et al* (2005a) *Am J Physiol*, 288(3):H1461-1467.
- Ogoh S *et al* (2005b) *Am J Physiol*, 288(4):H1526-1531.
- Ogoh S *et al* (2003) *J Physiol* 550, 317-324.
- Ogoh S *et al* (2005c) *J Physiol* (in press).