<招待講演 1>

Cerebral Blood Vessels and Stroke: Some New (and Forgotten) Principles

Donald D. Heistad

Department of Internal Medicine, University of Iowa Carver College of Medicine, U.S.A.

(脳卒中 29 : 680, 2007)

Some unique aspects of regulation of cerebral circulation

Resistance of large arteries is greater in the brain than in any other organ. Thus, blood pressure in the cerebral microcirculation is remarkably low. Cerebral microvessels have thinner walls than microvessels in other organs, but they are protected against rupture by the low level of intravascular pressure.

"Breakthrough" of autoregulation refers to large increases in cerebral blood flow at high levels of arterial pressure. This phenomenon generally is thought to be a passive, perhaps irreversible response. But the increase in blood flow often is reversible, and may be an active vasodilator response that is mediated by activation of K⁺ channels.

Cerebral ischemia and stroke

Hypertrophy and inward "remodeling" of cerebral arterioles reduce wall stress and protect against hypertension. Sympathetic nerves, angiotensin II, and superoxide contribute to these structural changes that protect against stroke.

TIAs with repetitive similar symptoms are intriguing. It seems unlikely that platelets or thrombi would

embolize repeatedly from carotid arteries to the same location in the brain. An alternative possibility is that aggregation of platelets in a large extracranial artery might release vasoconstrictor substances that produce vasospasm, and ischemia in a downstream region.

It is amazing that the primary mediator of vasospasm after subarachnoid hemorrhage (SAH) is not well defined. The most logical therapy to prevent vasospasm after SAH might be to block vasoconstrictor responses. An alternative approach, which works very well in prevention of vasospasm in experimental models of SAH, is gene transfer of calcitonin gene-related peptide, an extremely potent cerebral vasodilator.

Hemorrhagic stroke

In contrast to the large number of studies of mechanisms of ischemic stroke, relatively few studies focus on intracranial hemorrhage (ICH). We have developed a model of spontaneous ICH in hypertensive mice. It appears that oxidative stress is increased in these mice. The finding is of special interest because matrix metalloproteinases, which appear to play an important role in ICH, are activated by oxidative stress.