イブニングセミナー 1 特別講演

Emotions, mood and behavior after stroke

Prof. Dr. Julien Bogousslavsky

Department de Neurologie Center Hospitalier Universitaire Vaudois CHUV
CH-1011 Lausanne, Switzerland

While emotional outcome may be a critical factor influencing early evolution and late prognosis after stroke, few relevant studies have been performed on that subject. However, mood changes, modified judgement, and emotional reactions may also dramatically alter recruitment into clinical trials; for instance, up to one third of patients with acute stroke may have altered time perception, inappropriate self-evaluation of their condition, and attentional or memory dysfunction, with subsequent increase in referral-to-hospital delays. Also, the value of the "informed" consenting process may be questionable, in the setting of urgent randomization into an acute stroke clinical trial. Data from ongoing studies suggest that behavior and emotional reactions in acute stroke patients may be classified into a few broad categories, with considerable overlap. Correlations with the type, location and severity of stroke may give useful findings for improving management of patients, including prediction of functional evolution and late prognosis. While depressive reactions have been rather widely studied in the recovery-rehabilitation phase after stroke, consistent significant depression is uncommon shortly after stroke. On the other hand, related though different emotional behavioral changes may be more frequent, and have often been confused with depression, including catastrophic reaction, emotionalism or athymhormia. Late depression is the most common mood alteration during the first year after stroke. It bears specific characteristics, which makes it different from classical endogenous and reactive depression, emphasizing a critical role of brain damage in the pathogenesis of poststroke depression. Early recognition and management of mood disorders after stroke seem critical for the functional improvement of individual patients. However, little remains known on the specific indications of different antidepressant drugs in poststroke depression and related disorders. Ongoing research has identified a "new" emotional-behavioral disorder called poststroke fatigue, which is clearly separate from depression in most instances. It is especially disabling and frustrating, in that it typically involves patients with total or nearly-total neurological recovery, who should have been able to go back to their previous activities, but who become severely disabled because of early and persisting exhaustion. Preliminary neuropsychological, MR and PET imaging suggest that disruption of subtle mechanisms underlying attention may be responsible, in the absence of significant cognitive and mood alterations. Research projects are now being launched to better delineate poststroke fatigue and its management.

Acknowledgement: I would like to thank all the collaborators of the Lausanne Emotion in Stroke Study, who made this work possible: F. Guenther-Genta, P. Guex, F. Ghika-Schmid, A. Berney, A. Nicola, S. Aybek, A. Carota, F. Staub, L. Urben.