食肉類（ビーグルを中心として）の眼底検査とその問題点

福井正信、古川敏紀

国立予防衛生研究所 獣疫部

わが国における実験動物の眼底検査は、10数年ほど前の福井らのビーグルを主たる対象とする活動に端を発する。以来、有蹄類、霊長類、ネコ、ウサギ、その他マウス・ラットなどの小型齧歯類にいたるまで、検眼鏡や眼底カメラを使用しての調査・研究・検査が行われるようになった。

今回、ここに採り上げる食肉類は、その中でも歴史は古く、経験と蓄積の多さでは他の実験動物にはるかにまさる。比較的大型の実験動物として、イヌは取扱いが容易であり、目的によっては無麻酔下実験も可能なことから、多くの薬理・薬物実験に用いられてきた。

しかしながら、実験動物の中で、食肉類に属するイヌの眼底は、ヒトなどのそれと大きく異なり、タペタム(Tapetum lucidum)と呼ばれる特殊な光輝板を、眼底の網膜と脈絡膜の間に保有している。これは、組織学的にはPremelanomaterialとされており、事実、Melanocyte中のメラニン産生に必須とされる亜鉛のキレート剤投与や、成長過程における亜鉛欠乏食投与によって、形成に障害がみられる。このタペタムは、同じく食肉類のネコにもみられ、両者とも、そのタペタム組織は細胞性であり、注意深く観察すれば、小さな粒の集まりであることが理解される。これは、同じくタペタムを有する有蹄類（ウシ・ウマ・ヤギなど）のそれとは異なり、さらにタペタムを有しない高等なサル類（カニクイザルやアカゲザル）、齧歯類動物（マウス、ラット、モルモットなど）、ウサギなどとも異なる。この点が、本種動物の検査をやや困難にさせている理由でもある。

撮影条件として、準備すべき機器、その他を以下に列挙する。

通常撮影時：眼底カメラ（携帯用）。

電源。

フィルター（タペタムからの反射を抑え、露光過度の像を防ぐため）。

散瞳剤、縮瞳剤。

必要に応じて麻酔・鎮静剤など。

単色光撮影時：上記機材に加えての各種フィルター類。

蛍光撮影時：蛍光眼底撮影用フィルター、蛍光剤（静注用10%フルオレスセイン液）。

通常撮影は、多数の動物をスクリーニング的に行う場合に用いる。これに対し、単色光撮影や蛍光撮影は、やや特殊な目的に使用される。前者は、食肉類のように、タペタムを有する動物の眼底検査には特に有効で、微少

な障害部位を明確に特定するために必要である。これに対し、蛍光眼底撮影は、眼底血管系の観察に有用であり、タペタムを有する食肉類においても、フィルター類（励起用フィルター及び濾過用フィルター）の好選択と、適当な濃度・量の蛍光剤投与を行い、かつ、撮影手技に習熟すれば、ヒト蛍光眼底撮影像と同程度の像を得ることが出来る（Fig. 1-4）。我々の収集し得た像の中から、幾つかの異常眼底例も併せて示す（Fig. 5-8）。

食肉類動物にみられる薬物中毒について、幾つか報告のみられるものを、ここに掲げる。Phenothiazines¹⁾, Quinolines²⁾, Piperidylchlorophenothiazine³⁾, Vasodilates⁴⁾, Ethambutol⁵⁾, Dithizone⁶⁾, Rafoxanide⁷⁾, Naphthalene⁸⁾, Hexachlorophene⁹⁾, Chloroquin¹⁰⁾, Quinine derivatives¹¹⁾, Trenimon¹²⁾, Iodate¹³⁾, Iodocetate¹⁴⁾, Glutamate¹⁵⁾, Taurin¹⁶⁾, Aspartate¹⁷⁾, Lead¹⁸⁾, Methanol¹⁹⁾, Arsenicals²⁰⁾, Cobalt²¹⁾, Oxygen²²⁾などがそれである（Table 1）。

食肉類中イヌの眼底検査を行う上で、忘れてならないものに、各犬種の有する先天性眼疾患がある²³⁾。我々は近年、ネコ各品種にみられるネコ中心性網膜変性（FCRD）を対象を絞り、調査を進めているが^{24, 25)}、先天性疾患は、たくみに利用することで、ヒト先天性疾患モデル動物への、食肉類利用の道を開くものと考ええる。

引用文献（抄）

- 1) Meier-Ruge, W. & Cerletti, A. (1966): Zur experimentelle Pathologie der Phenothiazine-Retinopathie. *Ophthalmologica*, 151, 512-533.
- 2) Jess, A. (1921): Die Gefahren der Chemotherapie für das Auge, insbesondere über eine das Schrägen schwer schädigende Komponente des Chinins und seiner Derivate. *Albrecht von Graefes Arch. Ophthalmol.*, 104, 48-74.
- 3) Rintelen, F.G., & Wagner, P. (1957): Zur Klinik und Experimentellen pathologie der Pigmentepithelerkrankung nach Medikation mit einem Piperidinphenothiadine. *Ophthalmologica*, 133, 277-283.
- 4) Rubin, L.F. (1974): *Atlas of Veterinary Ophthalmoscopy*. Lea & Febiger, Philadelphia, 428.
- 5) Vogel, A.W. & Kaiser, J.A. (1963): Ethambutol induced transient change and reconstitution (in vivo) of the tapetum lucidum color in the dog. *Exp. Mol. Pathol. suppl.* 2, 136-149.
- 6) Scholler, J., Brown, D.E., & Timmens, E.K. (1961): Toxicological and Pathological studies with diethyldithiocarbamate. *Pharmacologist*. 3, 62.
- 7) Brown, W.R., Rubin, L., Hite, M., & Zwickey, R.E. (1971): Experimental papilledema induced by a salicylanilide. *Toxicol. Appl.*

Pharmacol. 27, 532-541.

8) Pirie, A.(1968): Pathology in the eye of the naphthalene-fed rabbit. Exp. Eye Res. 7, 354-357.

9) Kimbrough, R.D. (1971): Review of the toxicity of hexachlorophene. Arch. Environ. Health. 23, 119-122.

10) Berson, E.L. (1970): Acute toxic effects of chloroquine on the cat retina. Invest. Ophthalmol. 9, 618-628.

11) Holden, W.A. (1898): The Pathology of experimental quinine amblyopia. Arch. Ophthalmol. 27, 583-592.

12) Apponi, G.O., & Rinaldi, E. (1964): The retinotoxic action of 2:3:5:-trichylenimino 1:4-benzoquinone (Trenimon) Acta Ophthalmol., 42, 64-67.

13) Grignolo, A.N. & Calabria, G.A. (1966): Studies on the fine structure and the rhodopsin cycle of the rabbit retina in experimental degeneration induced by sodium iodate. Exp. Eye Res., 5, 86-97.

14) Noell, W.K. (1951): The effects of iodoacetate in the vertebrate retina. J. Cell Comp. Physiol., 37, 283-307.

15) Lucas, D.R. & Newhouse J.P. (1957): The toxic effect of sodium L-glutamate on the inner layers of the retina. Arch. Ophthalmol., 58, 193-201.

16) Schmidt, S.V., Berson, E.L., & Hayes, K.C. (1976): Retinal degeneration in cats fed casein. I. Taurine deficiency. Invest. Ophthalmol., 15, 47.

17) Rubin, L.F. (1974): Atlas of Veterinary Ophthalmoscopy. Lea & Febiger, Philadelphia, 447.

18) Hass, G.M., Brown, D.V.L., & Eisenstein, R., & Hemmens, A.(1964): Relations between lead poisoning in rabbit and man. Am. J. Pathol. 45, 691-727.

19) Gilger, A. & Potts, A. (1955): Studies on the visual toxicity of methanol. Am. J. Ophthalm., 39, 63-86.

20) Longley, B.J., Clausen, N.M., & Tatum, A.L.(1942): The experimental production of primary optic atrophy in monkeys by administration of organic arsenical compounds. J. Pharmacol. Exp. Ther., 76, 202-206.

21) Alagna, G., & d'Aquino, S.(1956): Alterazioni oculari da cloruro di cobalto. Arch. Ottal., 60, 5-29.

22) Beehler, C.C., Newton, N.L., Culver, J.F., & Tredici, T.J. (1964): Retinal detachment in dogs resulting from oxygen toxicity.

Arch. Ophthalmol., 71, 665-670.

23) Blogg, J.R. (1980): Breed Predisposition to Eye Disease. W.B. Saunders Comp., Philadelphia, 501-510.

24) 福井正信、林昭一、稲月一高(1984): ネコの眼底異常、とくに非感染性疾患について 比較眼科3: 27-32.

25) Fukui, M., Furukawa, T., Kurashima, Y., Hayashi, S. & Sugiyama, F. (1985): A Case of Asymmetric Feline Central Retinal Degeneration. Anim. Eye Res. 4: 23-26.

参考図書

1) 福井正信(1976): 実験用ビーグルの眼底撮影について。実験用ビーグルの研究(福井、友田、上田編)、287-297, ソフトサイエンス社、東京。

Table 1 List of Chemicals showing Toxicity on Ocular Fundus
of Carnivores

Chemicals	Conditions
Phenothiazines	Retinopathy
Quinolines	Retinopathy
Piperidylchlorophenothiazine	Retinopathy
Vasodilates	Retinal color change
Ethambutol	Retinal decoloration
Dithizone	Retinal edema, Degeneration
Rafoxanide	Optic nerve edema
Naphthalene	Retinal edema, Retinal atrophy
Hexachlorophene	Edema of optic disc
Chloroquine	Atrophy, Degeneration
Quinine derivatives	Optic nerve atrophy
Trenimon	Retinal edema
Iodate	Retinal atrophy
Iodocetate	Degeneration
Glutamate	Retinopathy
Taurin	Retinopathy
Aspartate	Retinopathy
Lead	Retinopathy
Methanol	Retinal edema
Arsenicals	Optic nerve atrophy
Cobalt	Retinal edema
Oxygen	Retinal detachment

Explanation of figures

FLUOROSCEIN ANGIOGRAPHS OF BEAGLE

- Figure 1) Fundus photograph 3.2 sec after injection with 10% solution of sodium fluorescein. (Early anterior phase)
Figure 2) " 5.4 sec after injection with 10% solution of sodium fluorescein. (Late anterior ph.)
Figure 3) " 7.6 sec after injection with 10% solution of sodium fluorescein. (Early venous ph.)
Figure 4) " 9.8 sec after injection with 10% solution of sodium fluorescein. (Late venous ph.)

* The dog is adult male.

SOME ABNORMAL CASES RECORDED IN CARNIVORES

- Figure 5) Injurious retinal detachment of adult male beagle by a traffic accident.
Figure 6) Retinal atrophy, adult male Japanese native dog.
Figure 7) Retinal atrophy, adult female Japanese domestic shorthaired cat.
Figure 8) Collie ectasia syndrome observed in an adult male.

Ophthalmoscopic examination of the canine ocular fundus,
in particular in the beagle and some other breeds

Masanobu FUKUI, Toshinori FURUKAWA

Department of Veterinary Science
National Institute of Health of Japan
35-10, Kamiosaki 2 chome, Shinagawa-ku, Tokyo, 141

At academic meetings in this country since 1971, Fukui and his colleagues have presented standard and abnormal retinal findings as seen in dogs and other mammals used for biomedical research.

Among mammals used for research, study of the ocular fundus of the dog has the longest history, for the dog is easy to handle, and its ocular fundus can be easily observed, just as in human patients, without the use of anesthetics.

The ocular fundus of the dog, which belongs to the order Carnivora, has a zone which reflects light, the tapetum lucidum. This tissue can be observed between the retina and the choroid by low-power light microscopic examination.

The cat, another domestic animal belonging to the Carnivora, has an even more brilliantly reflecting zone in the posterior part of the eye. Therefore, investigators must devise some kind of filter or light reducing apparatus to avoid trouble from reflected light when examining the ocular fundus.

The portable ophthalmoscopic camera, although developed for human eye research, can be used in animal research as well, particularly for non-human primates, rabbits, small ruminants, and carnivores.

For drug-safety assessment, the beagle is one of the most useful mammals presently used. The ocular fundus should be checked in dogs at the time of introduction to the research facility, and before the application of target chemicals. Animals with abnormal findings in the ocular region should be excluded from groups used for testing.

It is preferable that a team for investigating and recording findings should consist of 3 persons of equal academic achievement and technical skill.

The area centralis, the optic disk, and all corners of the retina in the posterior part of the eye should be observed.

Generally, in the course of an ocular toxicologic check, regular photography is used as a first step in the examination. Later, monocular photography with color film and fluorescein angiography may be used for special purposes.

Regular photography with a mono peaked gelatin filter at the light source is an effective method for examining the tapetal fundus in animals that have brilliant reflection from this structure, such as carnivores, some species of ruminants, and prosimians. Monocular photography with some kind of barrier or excitation gelatin filter, on the other hand, is a useful technique for checking the retinal angiosystem. If effective filters are used, the photographs will yield just as much information as when the techniques are used on human subjects. Both of these techniques are simple and useful for testing. As a general rule, positive color film should be used in all cases.

As a final check, results obtained by light or electron microscopy of the examined eye should be compared with the recorded ophthalmoscopic data.

The chemicals which produce various types of ocular abnormalities are listed in Table I.

Congenital malformations or genetical abnormalities in the ocular region of carnivores, such as feline central retinal degeneration as seen in Japanese domestic shorthaired cats (Fukui et al., 1984, 1985) can be useful as animal models of human diseases.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8