Simultaneous Inactivation of the *p16*, *p15* and *p14* Genes Encoding Cyclin-Dependent Kinase Inhibitors in Canine T-Lymphoid Tumor Cells

Aki FUJIWARA-IGARASHI¹⁾, Yuko GOTO-KOSHINO¹⁾, Hiroyuki MOCHIZUKI¹⁾, Shingo MAEDA¹⁾, Yasuhito FUJINO¹⁾, Koichi OHNO¹⁾ and Hajime TSUJIMOTO^{1)*}

¹⁾Department of Veterinary Internal Medicine, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

(Received 7 August 2012/Accepted 10 January 2013/Published online in J-STAGE 24 January 2013)

ABSTRACT. The *p16*, *p15* and *p14* genes are widely known as tumor suppressor genes in human medicine. Although a large number of genetic and epigenetic aberrations in these genes have been reported in human malignancies, canine malignancies have not been well analyzed on the aberrations of these genes. In this study, the full-length complementary DNA (cDNA) of the canine *p16* gene was cloned using the 5' and 3' rapid amplification of cDNA ends methods. Based on the sequence data, primers specific for *p16*, *p15* and *p14* were designed. Using these primers, the expression of *p16*, *p15* and *p14* mRNAs could be individually evaluated by reverse transcriptase polymerase chain reaction. Genomic aberrations were also examined using genomic polymerase chain reaction. Two of the 6 canine lymphoid tumor cell lines did not express detectable levels of *p16*, *p15* and *p14* mRNAs, and wide-ranging deletions in the *p15-p14-p16* genomic locus were suspected. Wide-ranging deletions were also speculated in 2 of 14 dogs with T-cell lymphoid tumors. On the other hand, similar failure of amplification suggesting wide-ranging deletions were not observed in any of the 14 dogs with B-cell lymphoma. Deletion of the *p15-p14-p16* genomic locus could be one of the molecular aberrations in canine lymphoid tumor cells.

KEY WORDS: canine, lymphoid tumor, *p16*, tumor suppressor gene.

doi: 10.1292/jvms.12-0351; *J. Vet. Med. Sci.* 75(6): 733–742, 2013

The *p16*, *p15* and *p14* genes are widely known as tumor suppressor genes in human medicine [8, 11]. They are clustered in a region on human chromosome 9 corresponding to mouse chromosome 4, and canine chromosome 11 [9, 40]. The P16 and P15 proteins belong to the inhibitor of cyclin-dependent kinase 4 (INK4) family, and are known as INK4a (cyclin-dependent kinase inhibitor 2A [CDKN2A]) and INK4b (CDKN2b), respectively. By binding directly to CDK4 and CDK6, P16 and P15 block the formation of cyclin D-CDK complexes, resulting in the phosphorylation of retinoblastoma (RB) proteins and G1-phase cell cycle arrest. The *p14* gene, which is also known as *alternative reading frame* (*ARF*), regulates the cell cycle via an alternative pathway. By binding to the murine double minute 2 (MDM2) protein, P14 inhibits the degradation of P53 by MDM2. Because P53 induces G1-phase cell cycle arrest in conjunction with P21, the outcome of this pathway is similar to that of the P16 and P15 pathways. The human *p16* and *p14* genes are composed of 3 exons each, and are known to share common exons 2 and 3.

A large number of genetic and epigenetic aberrations of the *p16*, *p15* and *p14* genes have been reported in human malignancies. Of these 3 genes, inactivation of *p16* has been most frequently found in hematological malignancies, such

as lymphoma [2, 3, 5, 6, 10, 16, 17, 20, 24, 36, 44], leukemia [13, 18, 19, 22, 30, 38, 42] and multiple myeloma [25, 32] as well as other non-hematological malignancies of various types [25, 32]. Inactivation of *p16* results from 4 types of alterations, namely, homozygous deletion, promoter hypermethylation, loss of heterozygosity and point mutations. Inactivation of *p16*, occurring mainly due to deletions and methylations [8], was observed in 44% of hematological tumor patients [39]. Furthermore, deletions of the 3 genes occur more frequently in patients with T-cell malignancies than in those with B-cell malignancies [8]. By contrast, patients with B-cell malignancies exhibit a higher incidence of inhibition of expression due to hypermethylations [8].

Partial or whole sequence data of the *p16*, *p15* and *p14* open reading frames have been reported. Four reports on the canine *p16* gene have been published [7, 9, 21, 27] since the identification of the partial sequence of its exon 2 (GenBank accession no. FJ542309) [21]. In another study [1], the entire coding sequence of the canine *p15* gene (GenBank accession no. NM_00114629) and partial sequence of the canine *p14* gene (GenBank accession no. FM883643) were reported. However, the genomic locus harboring *p16*, *p15* and *p14* has not been thoroughly examined in dogs. The predicted canine *p16*, *p15* and *p14* gene locus, deduced from the corresponding loci in other species, is shown in Fig. 1. Canine *p16* and *p14* genes are thought to share common exons 2 and 3, with a unique alternatively spliced first exon. Further, the previously reported partial sequence of canine *p16* gene exon 2 exhibited high homology to the sequence of *p15* gene exon 2. Therefore, the primers designed to amplify exon 2 of the canine *p16* gene in previous studies [7, 9, 21, 27] would also amplify the *p15* and *p14* genes. To examine the expression

*CORRESPONDENCE TO: TSUJIMOTO, H., Department of Veterinary Internal Medicine, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan.

e-mail: atsui@mail.ecc.u-tokyo.ac.jp

©2013 The Japanese Society of Veterinary Science

Fig. 1. Predicted gene locus of the *p16*, *p15* and *p14* genes on canine chromosome 11 deduced from the gene locus on human chromosome 9. The identified and unidentified exons of *p16*, *p15* and *p14* are indicated by white and black squares, respectively. Arrow with asterisk (*) indicates a gap in canine genome database. Locations of the primers in exon 2 of *p16* used in previous studies [21] are indicated by daggers (†).

of *p16*, *p15* and *p14* independently, it would be necessary to determine the whole sequence of exon 1 of the canine *p16* gene.

Using primers that would amplify not only *p16* but also *p15* and *p14*, 2 previous studies reported low or undetectable expression of these 3 genes in canine lymphoma [9, 27]. Partial loss of chromosome 11, on which these genes are located, has also been reported in canine lymphoid malignancies using fluorescent *in situ* hybridization analysis and comparative genomic hybridization [9, 12, 40]. Based on these reports, the *p16*, *p15* and *p14* genes are speculated to be partially or completely deleted in canine lymphoid malignancies, similar to human lymphoid tumors.

The first purpose of this study was to determine the complete nucleotide sequence of the canine *p16* cDNA, in order to design primers specific to *p16*, *p15* and *p14*. The second purpose was to examine the aberration of the *p16*, *p15* and *p14* genes in the canine lymphoid tumor cell lines and primary tumor cells from dogs with various lymphoid tumors.

MATERIALS AND METHODS

Cells and patient samples: Six lymphoid tumor cell lines derived from dogs with naturally occurring lymphoid malignancies were examined. These included CLBL-1 (multicentric B-cell lymphoma) [34], GL-1 (B-cell acute lymphoblastic leukemia [ALL]) [29], UL-1 (renal T-cell lymphoma) [43], CL-1 (mediastinal T-cell lymphoma) [28], Nody-1 (alimentary T-cell lymphoma) [15] and Ema (mediastinal T-cell lymphoma) [15]. Four of these cell lines (CLBL-1, GL-1, UL-1 and CL-1) were cultivated in RPMI-1640 (Invitrogen, Grand Island, NY, U.S.A.) supplemented with 10% fetal bovine serum (Biowest, Nuaille, France), penicillin (100 units/ml) and streptomycin (0.1 mg/ml) (Sigma-Aldrich, St. Louis, MO, U.S.A.). The other 2 cell lines (Nody-1, Ema) were cultivated in RPMI-1640 supplemented with 20% fetal bovine serum (Nichirei Biosciences Inc., Tokyo, Japan), penicillin (100 units/ml), and streptomycin (0.1 mg/ml). All cell lines were maintained in an atmosphere of 5% CO₂.

Lymph node (LN) cells were obtained as a control from a

healthy dog kept for blood transfusion in our teaching hospital via fine-needle aspiration (FNA). The dog was kept in accordance with the guidelines of Animal Care Committee of the Graduate School of Agricultural and Life Sciences, the University of Tokyo.

Dogs with lymphoid tumors were cytologically or pathologically diagnosed. Furthermore, clonal rearrangement of the *immunoglobulin heavy chain (IgH)* gene or the *T-cell receptor gamma chain (TCRγ)* gene in tumor cells confirmed using previously described primers [4, 41] was examined to indicate their clonal origin and B- or T-cell lineage. Fourteen dogs (dogs nos. 1–14) with *IgH* rearrangement were included, and all were diagnosed with multicentric high-grade lymphoma. Primary tumor cell samples were obtained from the LN of these dogs by FNA. Of the other 14 dogs (dogs nos. 15–28) showing *TCRγ* rearrangement, 11 dogs were diagnosed as having alimentary lymphoma, 1 with ALL and 2 with chronic lymphocytic leukemia (CLL). Primary tumor cell samples were obtained by endoscopy from dogs with the alimentary lymphoma, and by peripheral blood mononuclear cell (PBMC) separation by gradient centrifugation [43] from the dogs with ALL and CLL. All primary tumor samples were obtained with owners' informed consent.

Full-length cloning of canine *p16* complementary DNA: Total RNA was isolated using Illustra RNA Spin Mini (GE Healthcare, Little Chalfont, Buckinghamshire, U.K.), followed by DNase I (Invitrogen) treatment. Complimentary DNA (cDNA) was synthesized using PrimeScript® RT reagent Kit (Takara Bio, Otsu, Japan). All protocols were performed in accordance with the manufacturer's instructions. To amplify a part of exon 1 of canine *p16*, we designed degenerative primers, p16 EXON-F94 and p16 EXON-R293 (Table 1), based on the sequence of *p16* exon 1 from other species (human, chimpanzee, cattle, mouse and rat). PCR amplification was performed with AmpliTaq gold® 360 (Applied Biosystems, Grand Island, NY, U.S.A.) using normal LN cDNA as template. Cycle conditions included an initial denaturation step at 95°C for 10 min, followed by 35 cycles of denaturation at 95°C for 3 sec, annealing at 68°C for 30 sec and extension at 72°C for 60 sec, with a final extension step at 72°C for 7 min. The PCR amplification products were electrophoresed on 2% agarose gel (BMBio, Tokyo, Japan), purified using Wizard® SV Gel and PCR Clean-Up System (Promega, Madison, WI, U.S.A.) and inserted into the pGEM®-T Easy Vector System (Promega). After transformation, plasmid DNA was extracted using NucleoSpin® Plasmid QuickPure (Macherey-Nagel, Düren, Germany) and subjected to sequence analysis using BigDye® Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) and 3130xl Genetic Analyzer (Applied Biosystems).

The 5' Rapid amplification of cDNA ends (5' RACE) method was used to determine the full-length sequence of exon 1 of the *p16* gene using 5'-full RACE Core Set (TakaraBio). CL-1 cDNA template was generated using the p16 EXON2-R1481 primer phosphorylated at the 5' end. After circularization using T4 ligase, the cDNA was amplified by nested-PCR using 4 primers: p16 EXON2-F1374 and p16EXON2-F1411, and p16 EXON 1-R169 and p16 EXON

Table 1. Primer sequences used for full-length complementary DNA cloning and expression analysis of the *p16* gene in this study

Primer name	Primer sequence (5'-3')	Genbank no., nucleotide number	Amplicon size (bp)
p16 EXON-F94	GASGTGCGGGCRCTGCTSG	AB675384, nt. 94-112	200
p16 EXON-R293	GTGTCCAGGAAGCCCTCCCG	AB675384, nt. 293-274	
p16 EXON2-R1481	P-CAACTCAGATGTGGC	FM883643, nt. 1481-1466	NA
p16 EXON1-R169	ATGACCTGAATCGGGCTCCGA	AB675384, nt. 169-150	
p16 EXON1-R146	AGGCGGTTTCGGCGCGTCGGGA	AB675384, nt. 146-126	
p16 EXON2-F1374	GCTGTCGCTGCGTACCTGCG	FM883643, nt. 1374-1385	
p16 EXON2-F1411	CCGAAAGTGGTAGCCACGCC	FM883643, nt. 1411-1430	
p16 EXON2-F153	GAGCCCGATTACAGTTCAT	AB675384, nt. 153-170	
p16 EXON1-F148	GGTCGGAGCCCGATTCA	AB675384, nt. 148-169	95
p16 EXON1-R242	ACGGGGTCGGCACAGTT	AB675384, nt. 242-226	
RPL13A-F	GCCGGAAGGTTGTAGTCGT	AJ388525, nt. 87-105	87
RPL13A-R	GGAGGAAGGCCAGGTAATTC	AJ388525, nt. 173-154	
p15 EXON1-F90	GCGGCAGCTCCTGGAAG	NM_00114629, nt. 90-106	140
p15 EXON1-R230	GGGTCGGCACAGTTGG	NM_00114629, nt. 230-215	
p14 EXON1-F330	GCCCTGGTGCTAAAGCTAGT	FM883643, nt. 330-349	154
p14 EXON1-R1230	TGACGGGGTCGGCACAGTT	FM883643, nt. 1230-1212	
GAPDH-F	TCACCAGGGCTGCTTTTAAC	AB038240, nt. 47-67	403
GAPDH-R	AGGAGGCATTGCTGACAATC	AB038240, nt. 448-429	

GAPDH, Glyceraldehyde-3-phosphate dehydrogenase; NA, not applicable (primers used for 5' and 3' RACE); P, Phosphate group; R represents A or G, S represents C or G.

1-R146 (Table 1). Three primers, p16 EXON2-R1481, p16 EXON2-F1374 and p16 EXON-2F1411, were designed on the basis of the exon 2 sequences of the *p16/p14* genes in GenBank (GenBank accession no. FM883643); the other 2 primers, p16 EXON1-R169 and p16 EXON1-R146, were designed on the basis of the partial sequence of *p16* exon 1 determined in this study (fragment amplified with p16 EXON-F94 and p16 EXON-R293 primers) (GenBank accession no. AB675384). Both the first and second PCR amplifications were performed using Takara LA Taq with GC Buffer (Takara Bio). Cycle conditions included an initial denaturation step at 94°C for 1 min, followed by 25 cycles of denaturation at 94°C for 30 sec, annealing at 62°C for 30 sec, and extension at 72°C for 60 sec, with a final extension step at 72°C for 5 min. The PCR product was cloned and sequenced as described above.

The 3' RACE method was used to determine the universal exon 3 of the *p16/p14* genes using 3'-full RACE Core Set (Takara Bio). A primer, p16 EXON2-F153, was designed on exon 2 of the *p16/p14* genes (Table 1). The first strand of cDNA was synthesized from CL-1 RNA using oligo-dT primer containing an adaptor sequence. Subsequent PCR amplification was performed with the p16 EXON2-F153 primer and a supplied primer complementary to the adaptor sequence using AmpliTaq gold® 360. The protocol consisted of an initial denaturation step at 95°C for 10 min, followed by 35 cycles of denaturation at 95°C for 3 sec, annealing at 60°C for 35 sec and extension at 72°C for 60 sec, with a final extension step at 72°C for 7 min. The PCR amplification product was cloned and sequenced as described above.

Real-time PCR analysis of *p16* expression in canine lymphoid tumor cell lines: The primers p16 EXON1-F148 and p16 EXON1-R242 were designed to amplify 95 base pairs

(bp) of *p16* exon 1 based on the *p16* sequence determined in this study (Table 1). The mRNA expression in each cell line was measured by real-time PCR using SYBR® Premix Ex Taq (TakaraBio). *Ribosomal protein L13A (RPL13A)* was used as the endogenous control (RPL13A F and RPL13A R; amplicon length 87 bp, Table 1). The cycle protocol was as follows: denaturation step at 95°C for 10 sec; 40 cycles of denaturation at 95°C for 5 sec and annealing/extension at 60°C for 30 sec; followed by the dissociation step at 95°C for 5 sec, 60°C for 30 sec and 95°C for 15 sec. The PCR amplicons were electrophoresed on 12.5% polyacrylamide gel (ATTO, Tokyo, Japan).

The comparative cycle threshold (Ct) method was used to quantify *p16* transcript levels. ΔC_t was determined by subtracting the Ct value of *RPL13A* from the Ct value of the *p16* gene. $\Delta\Delta C_t$ was calculated by subtracting the ΔC_t of the control sample from the ΔC_t of the target sample. Normal LN was used as the control sample in this study. The levels of the *p16* transcripts relative to that in the normal LN were calculated as $2^{-\Delta\Delta C_t}$. To confirm the amplification efficacy of the primers, a standard curve was generated using serial dilutions of the CL-1 cDNA as template. All samples were evaluated in triplicate.

Expression analysis of the *p15* and *p14* genes in canine lymphoid tumor cell lines: Primers specific for *p15*, p15 EXON1-F90 and p15 EXON1-R230, and *p14*, p14 EXON1-F330 and p14 EXON1-R1230, were designed based on the sequences of *p15* and *p14* registered in GenBank (accession nos. NM_00114629 and FM883643), respectively (Table 1). *GAPDH* was used as the endogenous control gene (primers GAPDH-F and GAPDH-R; Table 1). Reverse transcriptase-polymerase chain reaction (RT-PCR) amplification was performed for the *p15* and *p14* genes using AmpliTaq

Fig. 2. Locations for the genomic PCR amplification on the *p15-p14-p16* gene locus. Arrow with asterisk (*) indicates a gap in the canine genome database. Black bars at the bottom represent the regions amplified by 8 primer pairs.

Table 2. Primer sequences used to amplify the *p15-p14-p16* genomic locus in this study

Primer name	Primer sequence (5'–3')	Genbank no., nucleotide number	Amplicon size (bp)
p15 usEXON1-F1	TGCCCTAACATTAATTCTTCTCAG	AAEX02016331.1, nt. 54947–54971	399
p15 usEXON1-R1	GACCTTTGACCAGAATGAAGAGATA	AAEX02016331.1, nt. 54777–54753	
p15 usEXON1-F2	GGACATGTAATTTGAACCCAGATAG	AAEX02016331.1, nt. 44716–44740	303
p15 usEXON1-R2	TCCTAATCTCCCTTCCTAGACACTT	AAEX02016331.1, nt. 44462–44438	
p16 dsEXON3-F	CACCTTGTGGGAACCTCTAGAAAAGTA	AAEX02016329.1, nt. 1182–1206	529
p16 dsEXON3-R	TATTCAAAAGGGGTATTGGTCTGTGA	AAEX02016329.1, nt. 802–778	
p15 usEXON1-F	GAAGAGGGCTGAAAGCTGACCT	AAEX02016331.1, nt. 13506–13522	561
p15 dsEXON1-R	CCAAACTTAAAGTAGCGTTTGTGTGT	AAEX02016331.1, nt. 12984–12962	
p15 usEXON2-F	ATCAACAAAGTATAACGGACTCCTG	AAEX02016331.1, nt. 10872–10896	500
p15 dsEXON2-R	ACATTTGTAGCCTCCCTTCAACT	AAEX02016331.1, nt. 10169–10146	
p16 dsEXON1-F	CTCTCAAAATCCCTTCACACTTAAA	AAEX02016329.1, nt. 9417–9441	529
p16 dsEXON1-R	CTACCATTTTGAGGGTACCTCCTAT	AAEX02016329.1, nt. 8937–8913	
p16 dsEXON2-F	GTCATGATGATGGGCAGCACCCGCGT	AAEX02016329.1, nt. 7139–7164	1,093
p16 dsEXON2-R	AATGGTTCTCCCTTCAGAAAAAG	AAEX02016329.1, nt. 6095–6072	
p16 usEXON3-F	TGAGGTGCCACATCTGAGTT	AAEX02016329.1, nt. 6831–6856	2,600
p16 usEXON3-R	CAAGCATGCAGGGAAGAGTT	AAEX02016329.1, nt. 4555–4534	
GAPDHgen-F	TCACCAGGGCTGCTTTTAAC	AB038240, nt. 47–67	818
GAPDHgen-R	AGGAGGCATTGCTGACAATC	AB038240, nt. 448–429	

GAPDH, Glyceraldehyde-3-phosphate dehydrogenase.

gold® 360 and the cDNAs of the cell lines as templates in the following protocol: initial denaturation at 95°C for 10 min, followed by 35 cycles of denaturation at 95°C for 3 sec, annealing at 60°C for 30 sec and extension at 72°C for 60 sec, with a final extension step at 72°C for 7 min. The PCR amplification products were electrophoresed on 12.5% polyacrylamide gel.

Amplification of the *p15-p14-p16* gene locus on chromosome 11 by genomic PCR: Genomic DNA was extracted using QIAmp DNA Mini kit (Qiagen, Hilden, Germany). Eight pairs of primers (Table 2) covering a 100-kb region of canine chromosome 11, including the introns and exons of the *p16*, *p15* and *p14* genes, were designed (Fig. 2). The primer sequences are listed in Table 2. The *GAPDH* gene was used as the endogenous control. PCR amplification was performed

using AmpliTaq gold® 360 and the DNAs of cell lines and patient samples as templates. The PCR protocol was as follows: initial denaturation at 95°C for 10 min, followed by 35 cycles with denaturation at 95°C for 3 sec, annealing at 60°C for 30 sec and extension at 72°C for 60 sec, with a final extension step at 72°C for 7 min. Electrophoresis was performed on 2% agarose gel. Amplification using distilled water was regarded as negative control.

RESULTS

Full-length cloning of canine *p16* cDNA: Using 5' RACE and 3' RACE methods, the nucleotide sequence of full-length canine *p16* cDNA was determined. A 705-bp entire open reading frame encoding 151 amino acids could be iden-

		20		40	
Human	MEPAAGSSME	PSADWLATAA	ARGRVEEVRA	LLEAGALPNA	PNSYGRRPIQ 50
Chimpanzee	MEPAAGSSME	PSADWLATAA	ARGRVEEVRA	LLEAGALPNA	PNSYGRRPIQ 50
Cattle	MQPSLGSSME	TADLLAAAA	ALGWAEVRA	LLEAGASANA	PNRYGRSAIQ 50
Dog	-----ME	PFADWLASAA	ARGRADEVRA	LLAAGAPPDA	PNRLGRSPIQ 42
Mouse	-----ME	SAADRLARAA	AQGRVHDVRA	LLEAGVSPNA	PNSFGRTPIQ 42
Rat	-----ME	SSADRLARAA	ALGREHEVRA	LLEAGASPNA	PNTFGRTPIQ 42
		60	Exon 1	80	100
Human	VMMMGSARVA	ELLLLHGAEP	NCADPATLTR	PVHDAAREGF	LDTLVVLHRA 100
Chimpanzee	VMMMGSARVA	ELLLLHGAEP	NCADPATLTR	PVHDAAREGF	LDTLVVLHRA 100
Cattle	VMMMGSARVA	ELLLLHGADP	NCADPATLTR	PVHDAAREGF	LDTLVALHRA 100
Dog	VMMMGSTRVA	QLLLLHGANP	NCADPVTLTR	PVHDAAREGF	LDTLVVLHRA 92
Mouse	VMMMGNVHVA	ALLLNYGADS	NCEDPTTFSR	PVHDAAREGF	LDTLVVLHGS 92
Rat	VMMMGNVKVA	ALLLSYGADS	NCEDPTTFSR	PVHDAAREGF	LDTLVVLHQA 92
		120	Exon 2	140	
Human	GARLDVRDAW	GRLPVDLAE	LGHDRVARYL	RAAAGGTRGS	NHARI DAAEG 150
Chimpanzee	GARLDVRDAW	GRLPVDLAE	LGHDRVARYL	RAAAGGTRGS	NHARI DAAEG 150
Cattle	GARLDVRDAW	GRLPVDLAE	RGHDRVARYL	RAAAEDTEGG	SHASADSAEG 150
Dog	GARLDVRDAW	GRLPVDLAE	RGHGAVAYL	RAAAGGTESG	SHARTEGAEG 142
Mouse	GARLDVRDAW	GRLPLDLAQE	RGHQDIVRYL	RSAGCSLCSA	GWSLCTAG - - 140
Rat	GARLDVRDAW	GRLPLDLALE	RGHHDVVRYL	R - - - - -	-YLLSSAG - - 130
		160	Exon 2	180	200
Human	PS - - - - -	- - - - -	- - - - -	-DI PD - - - -	156
Chimpanzee	PL - - - - -	- - - - -	- - - - -	-DI PD - - - -	156
Cattle	PAGQGDTFA	SSRLELCVVN	NLALVIDPLC	SLNPTGCSTA	VSLPRRAPRE 200
Dog	HA - - - - -	- - - - -	- - - - -	-DSPDFKN - -	151
Mouse	- - - - -	- - - - -	- - - - -	-NVAQ - TDGH	148
Rat	- - - - -	- - - - -	- - - - -	-NVSRTDRH	139
		220	Exon 3		
Human	- - - - -	- - - - -	- - - - -	- - - - -	156
Chimpanzee	- - - - -	- - - - -	- - - - -	- - - - -	156
Cattle	RLGSPEQPRL	AADALRSSGG	- - - - -	- - - - -	220
Dog	- - - - -	- - - - -	- - - - -	- - - - -	151
Mouse	SFSSSTPRAL	ELRGQSSEQS	- - - - -	- - - - -	168
Rat	NFCSSTPRCL	GLRGQPPKQR	- - - - -	- - - - -	159

Fig. 3. Amino acid sequence comparison of the deduced canine P16 with P16 of human (GenBank accession no. NM_000077.3), chimpanzee (GenBank accession no. NM_001146290.1), cattle (GenBank accession no. XM_868375.4), mouse (GenBank accession no. NM_009877.2) and rat (GenBank accession no. NM_031550). The corresponding exons are indicated below the sequence. Dash denotes gap.

tified (GenBank accession no. AB675384, nucleotide [nt] 40–492). In the dog genome database, nt no. 166–472 and nt no. 473–705 of canine *p16* cDNA were found on canine chromosome 11 separately using BLAST (National Center for Biotechnology Information, Bethesda, MD, U.S.A.) (nt no.166–472, 32195005–32195311; nt no. 473–705, 32192558–32192790). However, no sequence corresponding to the 5' fragment of canine *p16* (nt no. 1–165) was found in the dog genome database. Because there is a gap region between exon 1β of the *p14* gene and exon 2 of the *p16/p14* genes where its exon 1α is expected to be located (Figs. 1 and 2) in the dog genome database, the 5' fragment of canine *p16* (nt no. 1–165) was expected to be located in this region. Nt no. 1–165 of canine *p16* was considered to be its exon 1α, and nt no. 166–472 and nt no. 473–705 were regarded as its exons 2 and 3 from the result of the alignment of the sequence (nt no.166–705) of canine *p16* cDNA using BLAST. Therefore, canine *p16* gene was assumed to be composed of 3 exons.

The determined full-length cDNA sequence and the de-

duced amino acid sequence of canine *p16* were compared with those of other species. The cDNA sequence of the coding region shared 85, 86, 79 and 76% identities with human, chimpanzee, cattle, mouse and rat sequences, respectively. The deduced amino acid sequence of canine *p16* was 81, 81, 82, 61 and 76% homologous to that of human, chimpanzee, cattle, mouse and rat, respectively (Fig. 3). Exons 1 and 2 of canine *p16* were shown to contain 4 ankyrin repeats highly conserved among various species [33].

Expression analysis of *p16*, *p15* and *p14* genes using real-time RT-PCR or RT-PCR: The levels of *p16* mRNA were determined using real-time RT-PCR. The levels of *p16* mRNA in CLBL-1 and GL-1 cells were lower than those in normal LN cells, and below the limit of detection in UL-1, Nody-1 and Ema cells. The amount of *p16* mRNA in CL-1 was approximately 100 fold higher than that in normal LN cells (Fig. 4). Further, the levels of *p15* and *p14* expression were examined using RT-PCR. The levels of *p15* and *p14* mRNAs were uniformly high in 4 cell lines (CLBL-1, GL-1, UL-1 and CL-1) compared to those in normal LN cells.

Fig. 4. Expression analysis of the *p16* gene in canine lymphoid tumor cell lines and a lymph node tissue using real-time polymerase chain reaction (RT-PCR). Levels of *p16* mRNA were measured relative to that in normal LN tissue as a control sample. The expression of *p16* mRNA is below the detection limit in UL-1, Nody-1 and Ema. Data are expressed as the mean value of triplicate assay.

On the other hand, expression of *p15* and *p14* mRNAs was below the limit of detection in Nody-1 and Ema cells (Fig. 5). Amplification was not observed in these 2 cell lines even when the number of PCR cycles was increased from 35 to 40 (data not shown).

Amplification of the *p15-p14-p16* gene locus by genomic PCR: To examine deletion in the genomic locus of *p16*, *p15* and *p14* genes, we designed 8 primer pairs to amplify each region of the locus (Fig. 2). All the 8 primer pairs designed in the locus amplified the fragments of expected sizes in 4 cell lines (CLBL-1, GL-1, UL-1 and CL-1) and LN cells, whereas no amplification was observed with any of the 8 primer pairs in Nody-1 and Ema cells (Fig. 6).

Primary lymphoid tumor samples were also examined by genomic PCR. Due to limited DNA samples from lymphoid tumor patients, 5 primer pairs covering the exon regions of *p16*, *p15* and *p14* were used to examine each gene locus. Faint bands were presumed to arise from non-tumor cells derived from the surrounding normal tissues.

The 14 dogs diagnosed with multicentric high-grade lymphoma and harboring clonal *IgH* gene rearrangements all exhibited amplifications products of the expected sizes in the regions p15 EXON1, p16 dsEXON2 and p16 usEXON3 (Fig. 7). Amplified bands corresponding to p15 EXON2 and p16 dsEXON1 were faint to absent in 2 (dog nos. 4 and 9) and 1 (dog no. 7) of the 14 dogs, respectively. Eleven dogs showed successful amplifications with all the 5 primer pairs.

Among the 14 dogs with clonal *TCR γ* gene rearrangements (alimentary lymphoma, ALL and CLL), amplifica-

Fig. 5. Expression analysis of the *p15* and *p14* genes in canine lymphoid tumor cell lines and a lymph node tissue using RT-PCR. Thirty-five cycles of PCR reaction was performed for each gene. *GAPDH* (glyceraldehyde-3-phosphate dehydrogenase) was used as a control.

Fig. 6. Genomic PCR amplification of regions corresponding to the *p15-p14-p16* gene locus on chromosome 11. The location of each region is shown in Fig. 2. *GAPDH* gene was used as a control.

tions of region p15 EXON1, p15 EXON2, p16 dsEXON1, p16 dsEXON2 and p16 usEXON3 were faint to absent in 2 (dog nos. 15 and 23), 3 (dog nos. 15, 18, and 23), 5 (dog nos. 15, 18, 19, 20, and 23), 2 (dog nos. 15, and 23) and 2 (dog nos. 15, and 23) dogs, respectively (Fig. 7). In 2 dogs (dog nos. 15 and 23), no or faint bands were detected with any of the 5 primer pairs. The results of genomic PCR in primary lymphoid tumor samples are summarized in Table 3.

DISCUSSION

In this study, we identified the full-length nucleotide sequence of the canine *p16* gene. As expected, canine *p16* was composed of 3 exons, sharing exons 2 and 3 with the *p14* gene. The deduced canine *p16* amino acid sequence was highly homologous to that of human, chimpanzee and cattle throughout the coding region. On the other hand, the exon 3 sequence exhibited limited identity with that of mouse and rat, although exons 1 and 2 were relatively conserved. Exons

Fig. 7. Genomic PCR amplification for the *p15-p14-p16* gene locus in 28 canine primary tumor samples. The location of each region is shown in Fig. 2. Band intensities fainter than that of normal LN were considered as absence of amplification. Dogs 1–14 were diagnosed as having multicentric high-grade lymphoma with clonal *immunoglobulin heavy chain* gene rearrangement. Dogs 15–25 were diagnosed with alimentary lymphoma, dog 26 with acute lymphoblastic leukemia (ALL) and dogs 27 and 28 with chronic lymphocytic leukemia (CLL). Dogs 15–28 exhibited clonal *T-cell receptor gamma chain* gene rearrangements. Genomic DNA purified from normal LN tissue and peripheral blood mononuclear cell (PBMC) was used as controls. The result of PBMC was similar to that of LN (data not shown). Amplification using distilled water was regarded as a negative control.

Table 3. Results of genomic PCR to amplify the *p15-p14-p16* locus in primary lymphoid tumor samples

Dogs No.	Multicentric high-grade lymphoma														Alimentary lymphoma										ALL	CLL			
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
p15 EXON 1	+	+	+	+	+	+	+	+	+	+	+	+	+	+	−	+	+	+	+	+	+	+	±	+	+	+	+	+	+
p15 EXON 2	+	+	+	±	+	+	+	+	±	+	+	+	+	+	−	+	+	−	+	+	+	+	±	+	+	+	+	+	+
p16 ds EXON 1	+	+	+	+	+	+	−	+	+	+	+	+	+	+	−	+	+	−	−	±	+	+	−	+	+	+	+	+	+
p16 ds EXON 2	+	+	+	+	+	+	+	+	+	+	+	+	+	+	−	+	+	+	+	+	+	+	±	+	+	+	+	+	+
p16 us EXON 3	+	+	+	+	+	+	+	+	+	+	+	+	+	+	−	+	+	+	+	+	+	+	−	+	+	+	+	+	+

ALL, acute lymphoblastic leukemia; CLL, chronic lymphocytic leukemia; +, distinct band; ±, faint band; –, no band.

1 and 2 contained 4 ankyrin repeats, which are known to be highly conserved among various species [33]. The ankyrin repeat motif may be important for maintaining the functional conformation of P16.

The cloning of *p16* exon 1 enabled us to design primers that distinguished *p16* from *p15* and *p14*. Although exon 2 of *p15* and *p16/p14* was highly homologous, exon 1 of each gene shared only limited identity, and therefore it was possible to design primers specific to each gene. However, the primers specific for *p15* and *p14* generated several non-specific products, and could not be used for quantification. It was difficult to design another primer for *p15*, because of the high homology of its exon 2 with that of the *p16/p14* genes (98%). Further, polymorphism in exon 1 of the *p15* gene was reported in normal dogs [1]. Indeed, the reported polymorphism in *p15* exon 1 was also observed in CLBL-1, GL-1, UL-1, CL-1 and LN tissues by sequence analysis (data not shown). Therefore, we did not quantify *p15* expression by real-time RT-PCR.

In this study, a part of the *p16* nucleotide sequence (GenBank accession no. AB675384, nt 1–91) was obtained from CL-1 cells, but not confirmed in normal cells. Because *p16* expression was low in normal cells [38, 45], it was difficult to obtain a clear band from normal tissue by RT-PCR and 5 RACE. Instead, CL-1 cDNA was used as a template for PCR, because this cell line exhibited remarkably high expression of the *p16* gene. Further, it was not possible to verify the sequence from the genome database, because *p16* exon 1a is located in the gap region. The remaining part of the *p16* gene (nt 92–705) was identical between normal LN and CL-1 cells. Although the sequence of 5' end of canine *p16* cDNA should be verified in normal LN cells, the data gathered in this study conceivably reflected the sequence of canine normal *p16* cDNA.

The amount of *p16* mRNA expression was smaller in 5 (B-cell lines; CLBL-1 and GL-1, and 3 T-cell lines; UL-1, Nody-1 and Ema) of the 6 canine lymphoid tumor cell lines, compared to that in normal LN cells. *p15* and *p14*

mRNAs were also undetectable in 2 lymphoid tumor cell lines (Nody-1 and Ema). On the other hand, *p16* mRNA was highly expressed in CL-1 (T-cell line). It has been reported that RB protein becomes inactivated when it is excessively phosphorylated, which elicits high expression of *p16* mRNA due to a feedback mechanism [14, 26, 31]. Examination of the phosphorylation status of RB in CL-1 cells will be needed. Furthermore, when some of the cell cycle regulators such as *p21*, *p27*, *p57* and *p53* are inactivated and the progression of cell cycle occurs, over expression of *p16* mRNA could be induced as a feedback mechanism. Since mediastinal lymphoma is very rare in dogs, we did not have an opportunity to examine the *p16* mRNA expression level in primary tumor samples from dogs with mediastinal lymphoma. In previous studies on the prognosis of canine lymphoma [37], the presence of mediastinal mass was shown as a poor prognostic factor. A further study is needed to examine the relationship between the augmented *p16* expression and the disease type or the biologic behavior of lymphoid tumors in dogs.

In Nody-1 and Ema, *p16*, *p15* and *p14* mRNAs were not detected. Furthermore, genomic PCR amplification for the *p15-p14-p16* locus of the DNA samples with the 8 primer pairs did not generate any product in locus in Nody-1 and Ema. These results suggest the presence of wide-ranging deletions in chromosome 11 in Nody-1 and Ema. To confirm the exact status of the genetic deletions in these 2 cell lines, Southern blot analysis would be necessary. In the cells with such wide-ranging deletion, there should be a simultaneous loss of *p16*, *p15* and *p14* resulting in the failure of cell cycle control. Inactivation of these genes, especially in their combinations, was implicated in the tumorigenesis and malignancy from the studies of the *in vitro* and *in vivo* mouse model systems with targeted disruption of the corresponding genes [23]. Induction of wild-type *p16* into *p16*-deficient human lymphoma cell lines leads to growth retardation and partial differentiation, indicating that *p16* deficiency might contribute to the malignant phenotype [23]. Mice with targeted disruption of both *p16/p19* loci developed lymphomas and lymphoid leukemias with a low penetrance, as well as other tumors [35]. From these findings, the simultaneous inactivation of *p16*, *p15* and *p14* genes would be also associated with the tumorigenesis and malignancy in canine T-lymphoid tumor cells.

The amounts of *p16* mRNA in CLBL-1, GL-1 and UL-1 were shown to be lower than that in normal LN, whereas those of *p15* and *p14* mRNAs in these cell lines were similar to that in normal LN. If only the amount of *p16* mRNA is decreased despite the presence of *p15* and *p14* mRNAs as shown in normal LN, there is a possibility that deletion around *p16* gene exon 1 exists without wide-ranging deletion over *p15-p14-p16* locus. However, PCR amplification for the *p16* gene exon 1 generated a distinct band in these 3 cell lines using genomic PCR. It is possible that genetic aberrations exist in *p16* gene although not detected in this study, or epigenetic mechanism to suppress its expression might be related in these 3 cell lines.

We further examined whether genetic aberrations similar to that in Nody-1 and Ema exist in naturally occurring canine

lymphoid tumor cases. Similar failure of amplification of the *p15-p14-p16* locus (*p15EXON1*, *p15EXON2*, *p16dsEXON1*, *p16dsEXON2* and *p16usEXON3*) was observed in 2 dogs (dog nos. 15 and 23) with T-cell alimentary lymphoma, although several faint bands conceivably derived from concomitant normal cells were observed in dog no. 23. Such failure of amplification was not observed in any of the 14 multicentric B-cell lymphoma dogs.

Thomas *et al.* [40] indicated the frequent DNA copy number losses in *p15-p14-p16* locus of chromosome 11 in canine T-cell lymphomas but not in B-cell malignancies, using array-based comparative genomic hybridization. Although further studies are needed, deletion of *p15-p14-p16* locus would be an important candidate to understand the tumorigenesis in a subtype (possibly high-grade T-cell lymphoma) of canine lymphoid tumors.

Failure of amplification suggesting wide-ranging deletion of *p15-p14-p16* locus was observed in 2 canine lymphoid tumor cell lines, Nody-1 and Ema. Nody-1 was established from tumor cells obtained from a patient with alimentary lymphoma [15]. Furthermore, similar failure of amplification was observed in 2 dogs (dog nos. 15 and 23) with alimentary lymphoma. Thus, such aberration of *p15-p14-p16* locus might be a type of genetic changes observed in canine alimentary lymphoma.

On the other hand, 6 of 28 primary lymphoid tumor samples exhibited faint amplification failure with only 1 or 2 of the 8 primer pairs (B-cell lymphoid tumor nos. 4, 7 and 9; T-cell lymphoid tumor nos. 18, 19 and 20) in this study, suggesting a different genetic aberration in the tumor nos. 15 and 23. Failures of amplification could occur, if mutation, single nucleotide polymorphisms at the primer-binding region, or heterozygous genetic deletion is present. Southern blotting or array comparative genomic hybridization would be required to reveal the cause of the failure of amplification.

In conclusion, the full-length canine *p16* cDNA was cloned, and the levels of *p16*, *p15* and *p14* expression in canine lymphoid tumor cell lines were determined. Failure of amplification suggesting wide-ranging deletions were observed in 2 canine T-cell lymphoid tumor cell lines lacking *p16*, *p15* and *p14* expression. Similar failure of amplification was also found in 2 primary tumor samples from alimentary T-cell lymphoma. Deletion of the *p15-p14-p16* genomic locus could be one of the molecular aberrations in canine lymphoid tumor cells.

ACKNOWLEDGMENTS. The authors would like to acknowledge Dr. Barbara C. Rütgen, Institute of Immunology, Department of Pathobiology, University of Veterinary Medicine Vienna, for providing a canine lymphoma cell line (CLBL-1) and Dr. Takuya Mizuno, Laboratory of Veterinary Internal Medicine, Faculty of Agriculture, Yamaguchi University, for providing 2 canine lymphoma cell lines (Nody-1 and Ema). This study was supported by the Japan Society for the Promotion of Science, KAKENHI 24658265.

REFERENCES

1. Aguirre-Hernandez, J., Milne, B. S., Queen, C., O'Brien, P. C., Hoather, T., Haugland, S., Ferguson-Smith, M. A., Dobson, J. M. and Sargan, D. R. 2009. Disruption of chromosome 11 in canine fibrosarcomas highlights an unusual variability of CDKN2B in dogs. *BMC Vet. Res.* **5**: 27. [Medline] [CrossRef]
2. Amara, K., Trimeche, M., Ziadi, S., Laatiri, A., Hachana, M. and Korbi, S. 2008. Prognostic significance of aberrant promoter hypermethylation of CpG islands in patients with diffuse large B-cell lymphomas. *Ann. Oncol.* **19**: 1774–1786. [Medline] [CrossRef]
3. Baur, A. S., Shaw, P., Burri, N., Delacretaz, F., Bosman, F. T. and Chaubert, P. 1999. Frequent methylation silencing of p15(INK4b) (MTS2) and p16(INK4a) (MTS1) in B-cell and T-cell lymphomas. *Blood* **94**: 1773–1781. [Medline]
4. Burnett, R. C., Vernau, W., Modiano, J. F., Olver, C. S., Moore, P. F. and Avery, A. C. 2003. Diagnosis of canine lymphoid neoplasia using clonal rearrangements of antigen receptor genes. *Vet. Pathol.* **40**: 32–41. [Medline] [CrossRef]
5. Cheung, H. H., Lee, T. L., Rennert, O. M. and Chan, W. Y. 2009. DNA methylation of cancer genome. *Birth Defects Res. C Embryo Today* **87**: 335–350. [Medline] [CrossRef]
6. Chu, L. C., Eberhart, C. G., Grossman, S. A. and Herman, J. G. 2006. Epigenetic silencing of multiple genes in primary CNS lymphoma. *Int. J. Cancer* **119**: 2487–2491. [Medline] [CrossRef]
7. DeInnocentes, P., Agarwal, P. and Bird, R. C. 2009. Phenotype-rescue of cyclin-dependent kinase inhibitor p16/INK4A defects in a spontaneous canine cell model of breast cancer. *J. Cell. Biochem.* **106**: 491–505. [Medline] [CrossRef]
8. Drexler, H. G. 1998. Review of alterations of the cyclin-dependent kinase inhibitor INK4 family genes p15, p16, p18 and p19 in human leukemia-lymphoma cells. *Leukemia* **12**: 845–859. [Medline] [CrossRef]
9. Fosmire, S. P., Thomas, R., Jubala, C. M., Wojcieszyn, J. W., Valli, V. E., Getzy, D. M., Smith, T. L., Gardner, L. A., Ritt, M. G., Bell, J. S., Freeman, K. P., Greenfield, B. E., Lana, S. E., Kisseberth, W. C., Helfand, S. C., Cutter, G. R., Breen, M. and Modiano, J. F. 2007. Inactivation of the p16 cyclin-dependent kinase inhibitor in high-grade canine non-Hodgkin's T-cell lymphoma. *Vet. Pathol.* **44**: 467–478. [Medline] [CrossRef]
10. Garcia, M. J., Castrillo, J. M., Granizo, J. J., Cazorla, A. and Rivas, C. 2002. Clinicopathological correlation, p16-p15 methylation status and outcome predictors in anaplastic large cell lymphoma. *Br. J. Haematol.* **119**: 877–878. [Medline] [CrossRef]
11. Gil, J. and Peters, G. 2006. Regulation of the INK4b-ARF-INK4a tumour suppressor locus: all for one or one for all. *Nat. Rev. Mol. Cell Biol.* **7**: 667–677. [Medline] [CrossRef]
12. Hahn, K. A., Richardson, R. C., Hahn, E. A. and Chrisman, C. L. 1994. Diagnostic and prognostic importance of chromosomal aberrations identified in 61 dogs with lymphosarcoma. *Vet. Pathol.* **31**: 528–540. [Medline] [CrossRef]
13. Hangaishi, A., Ogawa, S., Imamura, N., Miyawaki, S., Miura, Y., Uike, N., Shimazaki, C., Emi, N., Takeyama, K., Hirose, S., Kamada, N., Kobayashi, Y., Takemoto, Y., Kitani, T., Toyama, K., Ohtake, S., Yazaki, Y., Ueda, R. and Hirai, H. 1996. Inactivation of multiple tumor-suppressor genes involved in negative regulation of the cell cycle, MTS1/p16INK4A/CDKN2, MTS2/p15INK4B, p53, and Rb genes in primary lymphoid malignancies. *Blood* **87**: 4949–4958. [Medline]
14. Hara, E., Smith, R., Parry, D., Tahara, H., Stone, S. and Peters, G. 1996. Regulation of p16CDKN2 expression and its implications for cell immortalization and senescence. *Mol. Cell. Biol.* **16**: 859–867. [Medline]
15. Hiraoka, H., Minami, K., Kaneko, N., Shimokawa Miyama, T., Okamura, Y., Mizuno, T. and Okuda, M. 2009. Aberrations of the FHIT gene and Fhit protein in canine lymphoma cell lines. *J. Vet. Med. Sci.* **71**: 769–777. [Medline] [CrossRef]
16. Hutter, G., Scheubner, M., Zimmermann, Y., Kalla, J., Katzenberger, T., Hubler, K., Roth, S., Hiddemann, W., Ott, G. and Dreyling, M. 2006. Differential effect of epigenetic alterations and genomic deletions of CDK inhibitors [p16(INK4a), p15(INK4b), p14(ARF)] in mantle cell lymphoma. *Genes Chromosomes Cancer* **45**: 203–210. [Medline] [CrossRef]
17. Jardin, F., Jais, J. P., Molina, T. J., Parmentier, F., Picquenot, J. M., Ruminy, P., Tilly, H., Bastard, C., Salles, G. A., Feugier, P., Thieblemont, C., Gisselbrecht, C., De Reynies, A., Coiffier, B., Haioun, C. and Leroy, K. 2010. Diffuse large B-cell lymphomas with CDKN2A deletion have a distinct gene expression signature and a poor prognosis under R-CHOP treatment: a GELA study. *Blood* **116**: 1092–1104. [Medline] [CrossRef]
18. Kees, U. R., Terry, P. A., Ford, J., Everett, J., Murch, A., De Klerk, N. and Baker, D. L. 2005. Detection of hemizygous deletions in genomic DNA from leukaemia specimens for the diagnosis of patients. *Leuk. Res.* **29**: 165–171. [Medline] [CrossRef]
19. Kim, M., Yim, S. H., Cho, N. S., Kang, S. H., Ko, D. H., Oh, B., Kim, T. Y., Min, H. J., She, C. J., Kang, H. J., Shin, H. Y., Ahn, H. S., Yoon, S. S., Kim, B. K., Shin, H. R., Han, K. S., Cho, H. I. and Lee, D. S. 2009. Homozygous deletion of CDKN2A (p16, p14) and CDKN2B (p15) genes is a poor prognostic factor in adult but not in childhood B-lineage acute lymphoblastic leukemia: a comparative deletion and hypermethylation study. *Cancer Genet. Cytogenet.* **195**: 59–65. [Medline] [CrossRef]
20. Klangby, U., Okan, I., Magnusson, K. P., Wendland, M., Lind, P. and Wiman, K. G. 1998. p16/INK4a and p15/INK4b gene methylation and absence of p16/INK4a mRNA and protein expression in Burkitt's lymphoma. *Blood* **91**: 1680–1687. [Medline]
21. Koenig, A., Bianco, S. R., Fosmire, S., Wojcieszyn, J. and Modiano, J. F. 2002. Expression and significance of p53, rb, p21/waf-1, p16/ink-4a, and PTEN tumor suppressors in canine melanoma. *Vet. Pathol.* **39**: 458–472. [Medline] [CrossRef]
22. Kohno, T., Yamada, Y., Tawara, M., Takasaki, Y., Kamihira, S., Tomonaga, M. and Matsuyama, T. 2007. Inactivation of p14ARF as a key event for the progression of adult T cell leukemia/lymphoma. *Leuk. Res.* **31**: 1625–1632. [Medline] [CrossRef]
23. Krug, U., Ganser, A. and Koefler, H. P. 2002. Tumor suppressor genes in normal and malignant hematopoiesis. *Oncogene* **21**: 3475–3495. [Medline] [CrossRef]
24. Lee, S. M., Lee, E. J., Ko, Y. H., Lee, S. H., Maeng, L. and Kim, K. M. 2009. Prognostic significance of O6-methylguanine DNA methyltransferase and p57 methylation in patients with diffuse large B-cell lymphomas. *APMIS* **117**: 87–94. [Medline] [CrossRef]
25. Li, J., Poi, M. J. and Tsai, M. D. 2011. Regulatory mechanisms of tumor suppressor P16(INK4A) and their relevance to cancer. *Biochemistry* **50**: 5566–5582. [Medline] [CrossRef]
26. Masumoto, N., Fujii, T., Ishikawa, M., Saito, M., Iwata, T., Fukuchi, T., Susumu, N., Mukai, M., Kubushiro, K., Tsukazaki, K. and Nozawa, S. 2003. P16 overexpression and human papillomavirus infection in small cell carcinoma of the uterine cervix. *Hum. Pathol.* **34**: 778–783. [Medline] [CrossRef]
27. Modiano, J. F., Breen, M., Valli, V. E., Wojcieszyn, J. W. and Cutter, G. R. 2007. Predictive value of p16 or Rb inactivation in a model of naturally occurring canine non-Hodgkin's lym-

- phoma. *Leukemia* **21**: 184–187. [Medline] [CrossRef]
28. Momoi, Y., Okai, Y., Watari, T., Goitsuka, R., Tsujimoto, H. and Hasegawa, A. 1997. Establishment and characterization of a canine T-lymphoblastoid cell line derived from malignant lymphoma. *Vet. Immunol. Immunopathol.* **59**: 11–20. [Medline] [CrossRef]
29. Nakaichi, M., Taura, Y., Kanki, M., Mamba, K., Momoi, Y., Tsujimoto, H. and Nakama, S. 1996. Establishment and characterization of a new canine B-cell leukemia cell line. *J. Vet. Med. Sci.* **58**: 469–471. [Medline] [CrossRef]
30. Omura-Minamisawa, M., Diccianni, M. B., Batova, A., Chang, R. C., Bridgeman, L. J., Yu, J., Pullen, J., Bowman, W. P. and Yu, A. L. 2000. Universal inactivation of both p16 and p15 but not downstream components is an essential event in the pathogenesis of T-cell acute lymphoblastic leukemia. *Clin. Cancer Res.* **6**: 1219–1228. [Medline]
31. Parry, D., Bates, S., Mann, D. J. and Peters, G. 1995. Lack of cyclin D-Cdk complexes in Rb-negative cells correlates with high levels of p16INK4/MTS1 tumour suppressor gene product. *EMBO J.* **14**: 503–511. [Medline]
32. Popov, N. and Gil, J. 2010. Epigenetic regulation of the INK4b-ARF-INK4a locus: in sickness and in health. *Epigenetics* **5**: 685–690. [Medline] [CrossRef]
33. Russo, A. A., Tong, L., Lee, J. O., Jeffrey, P. D. and Pavletich, N. P. 1998. Structural basis for inhibition of the cyclin-dependent kinase Cdk6 by the tumour suppressor p16INK4a. *Nature* **395**: 237–243. [Medline]
34. Rutgen, B. C., Hammer, S. E., Gerner, W., Christian, M., De Arezacochaga, A. G., Willmann, M., Kleiter, M., Schwendenwein, I. and Saalmuller, A. 2010. Establishment and characterization of a novel canine B-cell line derived from a spontaneously occurring diffuse large cell lymphoma. *Leuk. Res.* **34**: 932–938. [Medline] [CrossRef]
35. Serrano, M., Lee, H., Chin, L., Cordon-Cardo, C., Beach, D. and Depinho, R. A. 1996. Role of the INK4a locus in tumor suppression and cell mortality. *Cell* **85**: 27–37. [Medline] [CrossRef]
36. Shiozawa, E., Takimoto, M., Makino, R., Adachi, D., Saito, B., Yamochi-Onizuka, T., Yamochi, T., Shimosuma, J., Maeda, T., Kohno, Y., Kawakami, K., Nakamaki, T., Tomoyasu, S., Shio-kawa, A. and Ota, H. 2006. Hypermethylation of CpG islands in p16 as a prognostic factor for diffuse large B-cell lymphoma in a high-risk group. *Leuk. Res.* **30**: 859–867. [Medline] [CrossRef]
37. Starrak, G. S., Berry, C. R., Page, R. L., Johnson, J. L. and Thrall, D. E. 1997. Correlation between thoracic radiographic changes and remission/survival duration in 270 dogs with lymphosarcoma. *Vet. Radiol. Ultrasound* **38**: 411–418. [Medline] [CrossRef]
38. Takasaki, Y., Yamada, Y., Sugahara, K., Hayashi, T., Dateki, N., Harasawa, H., Kawabata, S., Soda, H., Ikeda, S., Tomonaga, M. and Kamihira, S. 2003. Interruption of p16 gene expression in adult T-cell leukaemia/lymphoma: clinical correlation. *Br. J. Haematol.* **122**: 253–259. [Medline] [CrossRef]
39. Taniguchi, T., Chikatsu, N., Takahashi, S., Fujita, A., Uchimaru, K., Asano, S., Fujita, T. and Motokura, T. 1999. Expression of p16INK4A and p14ARF in hematological malignancies. *Leukemia* **13**: 1760–1769. [Medline] [CrossRef]
40. Thomas, R., Seiser, E. L., Motsinger-Reif, A., Borst, L., Valli, V. E., Kelley, K., Suter, S. E., Argyle, D., Burgess, K., Bell, J., Lindblad-Toh, K., Modiano, J. F. and Breen, M. 2011. Refining tumor-associated aneuploidy through ‘genomic recoding’ of recurrent DNA copy number aberrations in 150 canine non-Hodgkin lymphomas. *Leuk. Lymphoma* **52**: 1321–1335. [Medline] [CrossRef]
41. Valli, V. E., Vernau, W., De Lorimier, L. P., Graham, P. S. and Moore, P. F. 2006. Canine indolent nodular lymphoma. *Vet. Pathol.* **43**: 241–256. [Medline] [CrossRef]
42. Wong, I. H., Ng, M. H., Huang, D. P. and Lee, J. C. 2000. Aberrant p15 promoter methylation in adult and childhood acute leukemias of nearly all morphologic subtypes: potential prognostic implications. *Blood* **95**: 1942–1949. [Medline]
43. Yamazaki, J., Baba, K., Goto-Koshino, Y., Setoguchi-Mukai, A., Fujino, Y., Ohno, K. and Tsujimoto, H. 2008. Quantitative assessment of minimal residual disease (MRD) in canine lymphoma by using real-time polymerase chain reaction. *Vet. Immunol. Immunopathol.* **126**: 321–331. [Medline] [CrossRef]
44. Zainuddin, N., Kanduri, M., Berglund, M., Lindell, M., Amini, R. M., Roos, G., Sundstrom, C., Enblad, G. and Rosenquist, R. 2011. Quantitative evaluation of p16(INK4a) promoter methylation using pyrosequencing in de novo diffuse large B-cell lymphoma. *Leuk. Res.* **35**: 438–443. [Medline] [CrossRef]
45. Zindy, F., Quelle, D. E., Roussel, M. F. and Sherr, C. J. 1997. Expression of the p16INK4a tumor suppressor versus other INK4 family members during mouse development and aging. *Oncogene* **15**: 203–211. [Medline] [CrossRef]