NOTE

Virology

Detection of *felis catus* papillomavirus type 3 and 4 DNA from squamous cell carcinoma cases of cats in Japan

Nanako YAMASHITA-KAWANISHI¹⁾, Ryoma SAWANOBORI¹⁾,
Kosuke MATSUMIYA¹⁾, Akiko UEMA¹⁾, James K. CHAMBERS²⁾,
Kazuyuki UCHIDA²⁾, Hidekatsu SHIMAKURA¹⁾, Masano TSUZUKI¹⁾,
Chia-Yu CHANG³⁾, Hui-Wen CHANG³⁾ and Takeshi HAGA^{1)*}

¹⁾Division of Infection Control and Disease Prevention, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-Ku, Tokyo 113-8657, Japan

²⁾Department of Veterinary Pathology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-Ku, Tokyo 113-8657, Japan

³⁾Graduate Institute of Molecular and Comparative Pathobiology, School of Veterinary Medicine, National Taiwan University, Taipei 10617, Taiwan

J. Vet. Med. Sci.

80(8): 1236–1240, 2018

doi: 10.1292/jvms.18-0089

Received: 23 February 2018

Accepted: 27 May 2018

Published online in J-STAGE:
5 June 2018

ABSTRACT. Association of *felis catus* papillomaviruses (FcaPVs) with feline squamous cell carcinoma (SCC) has been reported worldwide, while there is limited information about FcaPVs in Asia. In this study, 21 feline SCC biopsy samples from cats in Japan were analyzed by PCR with PV consensus primers and type-specific primers for FcaPV type 2 (FcaPV-2), FcaPV-3 and FcaPV-4 designed in this study. Sequence analysis revealed that one sample was FcaPV-3, and two were FcaPV-4. In both FcaPV-4 positive samples, 334th tryptophan in L1 ORF was deleted compared with the reference sequence. Moreover, immunohistochemistry showed that p16 protein was positive in both FcaPV-4 detected samples. This study would contribute to the molecular epidemiological and pathological understanding of FcaPV in Japan.

KEY WORDS: FcaPV type 3, FcaPV type 4, Feline squamous cell carcinoma, Japan, type specific primer

In domestic cats, five types of papillomaviruses (PVs), designated as FcaPV (*Felis catus* papillomavirus), have been reported in association with cutaneous neoplastic lesions. Each genotype is classified by the sequence of L1 ORF, the major capsid protein of PV. FcaPV type 1 (FcaPV-1) DNA was detected from flat sessile cutaneous hyperkeratotic lesions [21], oral papillomas [12] and oral SCCs [17]. FcaPV-2 have been detected most frequently from feline cutaneous neoplastic diseases [10], including viral plaques [13], Bowenoid *in situ* carcinomas (BISCs) [8], and cutaneous SCCs [18]. The partial genome of FcaPV-3, previously described as FdPV-MY2, was detected in viral plaque [13], and also in SCCs [14]. Later on, the full genome of FcaPV-3 was identified from a multiple BISCs case of a domestic cat [15]. FcaPV-4 full genome DNA was identified from feline oral cavity with ulcerative gingivitis [4], and its partial genome DNA, named as FdPV-MY3 was reported from SCC cases in earlier study [14]. FcaPV-5 was recently identified from a feline viral plaque case [16]. These studies show that FcaPVs may associate with the cutaneous and mucosal neoplastic diseases in domestic cats, although the pathogenesis of FcaPVs are still uncertain.

SCC accounts for 15% of cutaneous neoplastic diseases in cats [11], and share the most in feline oral neoplastic diseases [20]. However, the definitive cause of feline SCC is not well understood. The exposure to ultraviolet (UV) is considered as one of the environmental risk factors in developing cutaneous SCCs [3] while a recent study revealed that FcaPV DNA was detected more in UV-protected SCCs than in UV-exposed cases [18], suggesting the involvement of PV in the pathogenesis of feline SCCs. FcaPVs have been detected from feline SCC lesions in Europe [5, 10], Oceania [18], and North America [19]. From those studies, FcaPV-2 have been detected the most in feline SCCs, and it is also suggested to infect the skin of cats asymptotically [22]. In Asian countries including Japan, there is limited information about FcaPVs. Based on these backgrounds described above, we sought to detect FcaPV DNA from feline SCC lesions. Although PCR methods are commonly used, primers may affect the specificity and sensitivity for the amplification of PV DNA [18]. Therefore, in addition to the previously established consensus PV primers, the type specific primers designed in this study were used for detecting PVs from feline SCCs.

Twenty-one feline SCC biopsy samples, collected in Japan between 2013 and 2015, were fixed in 10% formalin and routinely

*Correspondence to: Haga, T.: ahaga@mail.ecc.u-tokyo.ac.jp

©2018 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Table 1. Descriptions of the feline SCC samples, detection of FcaPV and the results of p16 immunohistochemical analysis

Sample ID	Breed	Age, in years	Sex	Anatomical site	PV genotype (GenBank accession number)	p16 Immunohistochemistry
13-060	Mixed-breed	13	FX ^{a)}	Skin	ND ^{d)}	–
13-0107	Mixed-breed	12	FX	Oral (submandible)	ND	NA ^{g)}
13-136	Mixed-breed	14	FX	Skin	FcaPV-4 (LC333412) ^{e)}	+
13-0153	Mixed-breed	12	MX ^{b)}	Skin (eyelid, left)	FcaPV-3 (LC333418) ^{f)}	NA
13-0297	Mixed-breed	12	MX	Oral (lower gingia, right)	ND	–
13-0317	American shorthair	15	MX	Oral (tongue)	ND	–
13-848	Mixed-breed	10	FX	Auditory canal (left)	ND	–
13-882	Mixed-breed	16	FX	Oral (tongue)	ND	NA
13-944	Scottish fold	10	MX	Oral (tongue)	ND	–
14-778	Mixed-breed	13	M ^{c)}	Skin (upper jaw, right)	ND	–
14-1018	Mixed-breed	16	FX	Auditory canal (left)	ND	–
14-1110	American shorthair	10	MX	Skin (forelimb, digit, right)	FcaPV-4 (LC333413)	+
15-291	Mixed-breed	10	FX	Oral (upper gingiva)	ND	–
15-358	Mixed-breed	13	FX	Skin (auditory canal, left)	ND	–
15-498	Mixed-breed	9	MX	Skin (external ear, left)	ND	–
15-528	Mixed-breed	16	MX	Skin (sublingual, right)	ND	–
15-577	Mixed-breed	15	FX	Oral (upper gingiva, right)	ND	–
15-615	Mixed-breed	10	MX	Esophagus	ND	–
15-637	Mixed-breed	11	FX	Skin (submandible)	ND	–
5741-2013	Mixed-breed	12	MX	Anus	ND	+
6997-2015	Mixed-breed	9	M	Oral (buccal mucosa)	ND	NA

FcaPV genotype confirmed by sequence analysis are also shown for PV positive samples. a) Female (spayed); b) Male (castrated); c) Male; d) Not detected; e) *Felis catus* papillomavirus type 4; f) *Felis catus* papillomavirus type 3; g) Not available.

Fig. 1. Representative histopathological findings of feline squamous cell carcinoma. Atypical squamous cells invade the fibrous tissue. (A) Case number 14-1110, FcaPV4-positive. (B) Case number 13-0153, FcaPV3-positive. (C) Case number 14-0778, FcaPV negative. Bars, 100 μ m.

embedded in paraffin and stained with hematoxylin and eosin for histopathological examination. Histopathological diagnoses of SCC were made on the consensus of two veterinary pathologists (certified by the Japanese College of Veterinary Pathologists) at the Department of Veterinary Pathology, the University of Tokyo. Clinical and histopathological findings of feline SCC cases are summarized in Table 1. The mean age at the diagnosis was 12.3 ± 2.3 (mean $\pm$ standard deviation) years with no sex predilection. In the present study, seven out of 21 cases were oral, nine were cutaneous, and the rest of the SCC case lesions were observed in other locations. Histopathological examination revealed the invasive growth of squamous epithelial tissue of the skin or the mucosa (Fig. 1). Abundant mitotic figures and severe nuclear atypia were observed. Nuclear inclusion bodies, indicative of papilloma virus infection, were not observed in any of the cases.

Immunohistochemistry (IHC) was conducted to detect the PV antigen and p16 protein, with mouse anti-PV (clone BPV-1/1H8 +CAMVIR, Abcam, Cambridge, U.K.) and mouse anti-CDKN2A/p16INK4a (Becton Dickinson Co., Franklin Lakes, NJ, U.S.A.) monoclonal antibodies, respectively. Horseradish peroxidase labeled detection kit (Envision + system, Dako Japan, Kyoto, Japan) was applied, and antigen-antibody complex was visualized by the chromogen treatment with 3,3'-diaminobenzidine (DAB). Sections were counterstained with Mayer's hematoxylin. A section of cutaneous bovine papilloma, positive for bovine papillomavirus type-2, was used as the positive control for anti-PV antibody. For the negative control, the primary antibodies were substituted with tris-buffered saline. Positive staining of more than 10% of the neoplastic tissue was considered to be p16-positive. Immunohistochemical analysis for p16 showed diffuse positive staining in the neoplastic tissue of sample 13-136, 14-1110 (Fig. 2) and 5741-2013 (Table 1).

Fig. 2. Immunohistochemical analysis against p16 protein in feline squamous cell carcinoma. (A) Intense positive p16 staining in the basal cell membrane (FcaPV4-positive, case number 14-1110). (B) Negative for p16 (FcaPV-negative, case number 14-0778). Bars, 100 μ m.

Table 2. The primer information used in this study

Primer pair		Alignment 5'→3'	Target gene (location)	Amplicon length (bp)	T _m (°C)	Reference
ACTB	Forward	CAA CCG TGA GAA GAT GAC TCA GA	Feline beta actin	410	54	Kessler <i>et al.</i> , 2009
	Reverse	CCC AGA GTC CAT GAC AAT AAC A				
MY09/11	Forward	GCM CAG GGW CAT AAY AAT GG	PV L1 (Consensus)	approximately 450	55	Manos <i>et al.</i> , 1989
	Reverse	CGT CCM ARR GGA WAC TGA TC				
CP4/5	Forward	ATG GTA CAR TGG GCA TWT GA	PV E1 (Consensus)	approximately 450	49	Tieben <i>et al.</i> , 1994
	Reverse	GAG GYT GCA ACC AAA AMT GRC T				
FcaPV-2 L1	Forward	CGC AAG GAC AGA ATA ATG GTA TTT GCT	L1 (7130–7156 nt) (7548–7525 nt)	419	58	This study
	Reverse	AAG ACG ATC CGA GAT GTC AAC AT				
FcaPV-3 L1	Forward	TCT GGT AAT CAG TAT AGG GTG TTC AGA GT	L1 (5868–5876 nt) (6340–6312 nt)	473	59	This study
	Reverse	ATT TCT AAA GGC ACC CCT GAT TTG TCT T				
FcaPV-4 L1	Forward	CTT TGG TAA CCA GCG ATT CC	L1 (6571–6590 nt) (7046–7022 nt)	476	52	This study
	Reverse	CAA TCT ATC CTT CAA GTC CAC TAC				

The name of the primer pairs, the target region of PV ORF, expected size of the PCR product, annealing temperature (T_m), and the reference information are shown respectively.

For the immunohistochemical analysis for PV antigen, positive staining in the nucleus of epithelial cells in the section of bovine fibropapilloma was observed, while no nuclear-positive staining was detected in the feline SCC samples (data not shown). Some of the paraffin blocks, including FcaPV-3 positive sample (13-0153) were not available for IHC due to insufficient amount of paraffin blocks.

DNA from 21 formalin-fixed paraffin-embedded tissue samples were extracted with Qiagen DNA FFPE Tissue kit (QIAGEN, Hilden, Germany) and the success in DNA extraction of all samples were confirmed by PCR using feline beta-actin primer pair, ACTB [7] (Table 2). To detect the PV genome, two consensus primer pairs, MY09/11 [9] and CP4/5 [23] (Table 2) were used. From two samples (Sample ID: 13-0153 and 14-1110), papillomaviral DNA was detected with the CP4/5 consensus primer pair, while no positive results were obtained with MY09/11 primer (data not shown). According to the criteria of International Committee on Taxonomy of Viruses (ICTV), PVs are genotyped based on the sequence similarity of L1 ORF. Primer pair, CP4/5 is to detect the sequence of E1 ORF, so we have additionally designed type-specific primers for FcaPV-2, FcaPV-3 and FcaPV-4 L1 gene (Table 2) based on the reference sequence (GenBank accession number EU796884.1, NC_021472.1 and NC_022373.1 respectively) with GENETYX Ver.11 software (GENETYX Corp., Tokyo, Japan). Water in place of DNA template was added to the PCR mixture for the negative control. For the positive control of FcaPV-2 L1 primer, extracted DNA from a feline BISC case was applied. For FcaPV-3 and FcaPV-4 L1 primers, positive results for each FcaPV genotype was confirmed in this study (Table 1). PCR reactions were carried with the respective annealing temperature listed in Table 2, and each result was confirmed by electrophoresis. For the PCR positive samples, products were cloned and sequenced. Sequence analysis was performed using GENETYX Ver.11 software, and BLAST tool of the National Center for Biotechnology Information (NCBI) was used for the sequence similarity analysis. With FcaPV-3 type-specific primer, sample 13-0153 showed 99% sequence similarity with the reference FcaPV-3 (NC_021472.1). Sample 14-1110 also showed 99% sequence similarity with the reference FcaPV-4 (NC_022373.1) with FcaPV-4 type specific primer. Additionally, one sample (13-136) was detected with FcaPV-4 type-specific primer (Table 1), showing 95% sequence similarity with the reference FcaPV-4 (NC_022373.1). None of the samples were positive with FcaPV-2 L1 type specific primer (data not shown). The nucleotide sequences of the FcaPV-3 and FcaPV-4 identified in this

study were deposited in GenBank (Table 1).

Sequence analysis revealed that two FcaPV-4 samples detected in this study showed a single deletion of tryptophan (W) in the position of 334th amino acid in the reference FcaPV-4 L1 sequence. To investigate whether the deletion of 334th tryptophan is limited to FcaPV-4, the amino acid sequences of the reference FcaPVs from genotype 1 to 5 (available on GenBank) were aligned. Interestingly, the 334th tryptophan was found only in the reference FcaPV-4 (NC_022373.1). In order to examine whether the deletion affects the structure of L1 protein, the translated amino acid sequence of FcaPV-4 samples, 13-136 and 14-110 was submitted to SWISS-MODEL (<https://swissmodel.expasy.org/interactive>). As results, no significant change of the structure of L1 was found. Further studies are required to figure out the significance of the 334th tryptophan.

In Table 1, the anatomical locations of the SCC lesion are summarized. Nine cases out of 21 were cutaneous, and eight cases were oral SCC. All three FcaPV-positive samples were detected from cutaneous SCCs (Table 1), which was consistent with the previous report that PV-associated SCCs in cats are common in cutaneous cases than oral SCCs [17].

In the present study, IHC for detecting p16 protein was conducted, as positive staining for p16 is one of the observations indicative of PV-infection in feline SCC [18]. Since PVs are recognized to asymptotically infect the epidermis of cats [22], immunostaining to detect p16 within the lesions are preferable to support the involvement of PV to the lesion development. As expected, both of the FcaPV-4 positive samples, 13-136 and 14-110 (Fig. 2) were immunopositive for p16 (Table 1). One sample, 5742-2013 was also positive for p16 while neither FcaPV-2, -3 and -4 was detected by PCR. This result suggests that non-PV factors or infection of other PV genotypes may be present. For the immunostaining of PV antigen, none of the feline SCC samples showed nuclear-positive staining. This result was not unexpected, because capsid antigen of PV is frequently disappeared during the malignant progression [1], and the antibodies used in this study (monoclonal BPV-1/1H8) react to a major capsid protein.

This study confirmed the detection of FcaPV-3 and FcaPV-4 by conventional PCR. As shown in Table 1, one additional sample, 13-136 was detected with the FcaPV-4 type-specific primer, but not by the consensus primer. Several mutations in sample 13-136 were observed within the sequence amplified with FcaPV-4 L1 type specific primer, assuming that additional mutations may be present in other genomic regions. As mutations in the primer attachment site can affect the primer specificity, this may be one of the reasons why PV-genome could not be detected with the consensus primers. Designing the PV-type specific primers are considered to be advantageous for detecting the target PV-genotype, while it may have difficulties in identifying the unknown PV genotypes [6]. In addition to FcaPV type-specific primers, designing the FcaPV consensus primers may be preferable for detecting the unknown FcaPV genotype.

From the previous studies in other geographical regions, FcaPV-2 has been reported to be found in feline SCC most frequently [10, 18, 19]. On the other hand, FcaPV-2 was not detected in this study. From the study on human papillomavirus (HPV), the distribution of high risk HPV type in cervical cancer differed depending on the geographical region [2]. Although the sample size of this study is small, the geographical difference may explain the discrepancies of dominated type of FcaPV found in SCCs.

To further understand the pathogenicity of FcaPVs and the aetiology of feline SCCs, concert effort is required including larger scale of PV molecular-epidemiological studies to determine the geographical FcaPV genotype distributions, pathological studies to determine the pathogenicity of FcaPVs for the SCC development, and clinical observation studies to investigate the malignance between FcaPV-positive and negative feline SCCs.

REFERENCES

1. Azzimonti, B., Hertel, L., Aluffi, P., Pia, F., Monga, G., Zocchi, M., Landolfo, S. and Gariglio, M. 1999. Demonstration of multiple HPV types in laryngeal premalignant lesions using polymerase chain reaction and immunohistochemistry. *J. Med. Virol.* **59**: 110–116. [Medline] [CrossRef]
2. Crosbie, E. J., Einstein, M. H., Franceschi, S. and Kitchener, H. C. 2013. Human papillomavirus and cervical cancer. *Lancet* **382**: 889–899. [Medline] [CrossRef]
3. Dorn, C. R., Taylor, D. O. and Schneider, R. 1971. Sunlight exposure and risk of developing cutaneous and oral squamous cell carcinomas in white cats. *J. Natl. Cancer Inst.* **46**: 1073–1078. [Medline]
4. Dunowska, M., Munday, J. S., Laurie, R. E. and Hills, S. F. K. 2014. Genomic characterisation of *Felis catus* papillomavirus 4, a novel papillomavirus detected in the oral cavity of a domestic cat. *Virus Genes* **48**: 111–119. [Medline] [CrossRef]
5. Geisseler, M., Lange, C. E., Favrot, C., Fischer, N., Ackermann, M. and Tobler, K. 2016. Geno- and seroprevalence of *Felis domesticus* Papillomavirus type 2 (FdPV2) in dermatologically healthy cats. *BMC Vet. Res.* **2**: 1–10. [Medline] [CrossRef]
6. Haga, T., Dong, J., Zhu, W. and Burk, R. D. 2013. The many unknown aspects of bovine papillomavirus diversity, infection and pathogenesis. *Vet. J.* **197**: 122–123. [Medline] [CrossRef]
7. Kessler, Y., Helfer-Hungerbuehler, A. K., Cattori, V., Meli, M. L., Zellweger, B., Ossent, P., Riond, B., Reusch, C. E., Lutz, H. and Hofmann-Lehmann, R. 2009. Quantitative TaqMan real-time PCR assays for gene expression normalisation in feline tissues. *BMC Mol. Biol.* **10**: 1–14. [Medline] [CrossRef]
8. Lange, C. E., Tobler, K., Markau, T., Alhaidari, Z., Bornand, V. and Sto, R. 2009. Sequence and classification of FdPV2, a papillomavirus isolated from feline Bowenoid in situ carcinomas. *Vet. Microbiol.* **137**: 60–65.
9. Manos, M. M., Ting, Y., Wright, D. K., Lewis, A. J., Broker, T. R. and Wolinsky, S. M. 1989. The use of polymerase chain reaction amplification for the detection of genital human papillomaviruses. *Cancer Cell* **7**: 209–214.
10. Mazzei, M., Forzan, M., Carlucci, V., Anfossi, A. G., Alberti, A., Albanese, F., Binanti, D., Millanta, F., Baroncini, L., Pirone, A. and Abramo, F. 2017. A study of multiple *Felis catus* in cat skin lesions in Italy by quantitative PCR. *J. Feline Med. Surg.* doi: 10.1177/1098612x17732255. [Medline]
11. Miller, M. A., Nelson, S. L., Turk, J. R., Pace, L. W., Brown, T. P., Shaw, D. P., Fischer, J. R. and Gosser, H. S. 1991. Cutaneous neoplasia in 340 cats. *Vet. Pathol.* **28**: 389–395. [Medline] [CrossRef]
12. Munday, J. S., Fairley, R. A., Mills, H., Kiupel, M. and Vaatstra, B. L. 2015. Oral Papillomas Associated With *Felis catus* Papillomavirus Type 1 in 2 Domestic Cats. *Vet. Pathol.* **52**: 1187–1190. [Medline] [CrossRef]
13. Munday, J. S. and Peters-Kennedy, J. 2010. Consistent detection of *Felis domesticus* papillomavirus 2 DNA sequences within feline viral plaques. *J.*

- Vet. Diagn. Invest.* **22**: 946–949. [[Medline](#)] [[CrossRef](#)]
14. Munday, J. S., French, A. F., Gibson, I. R. and Knight, C. G. 2013. The presence of p16 CDKN2A protein immunostaining within feline nasal planum squamous cell carcinomas is associated with an increased survival time and the presence of papillomaviral DNA. *Vet. Pathol.* **50**: 269–273. [[Medline](#)] [[CrossRef](#)]
15. Munday, J. S., Dunowska, M., Hills, S. F. and Laurie, R. E. 2013. Genomic characterization of *Felis catus* papillomavirus-3: a novel papillomavirus detected in a feline Bowenoid in situ carcinoma. *Vet. Microbiol.* **165**: 319–325. [[Medline](#)] [[CrossRef](#)]
16. Munday, J. S., Dittmer, K. E., Thomson, N. A., Hills, S. F. and Laurie, R. E. 2017. Genomic characterisation of *Felis catus* papillomavirus type 5 with proposed classification within a new papillomavirus genus. *Vet. Microbiol.* **207**: 50–55. [[Medline](#)] [[CrossRef](#)]
17. Munday, J. S. and French, A. F. 2015. *Felis catus* papillomavirus types 1 and 4 are rarely present in neoplastic and inflammatory oral lesions of cats. *Res. Vet. Sci.* **100**: 220–222. [[Medline](#)] [[CrossRef](#)]
18. Munday, J. S., Gibson, I. and French, A. F. 2011. Papillomaviral DNA and increased p16CDKN2A protein are frequently present within feline cutaneous squamous cell carcinomas in ultraviolet-protected skin. *Vet. Dermatol.* **22**: 360–366. [[Medline](#)] [[CrossRef](#)]
19. O'Neill, S. H., Newkirk, K. M., Anis, E. A., Brahmabhatt, R., Frank, L. A. and Kania, S. A. 2011. Detection of human papillomavirus DNA in feline premalignant and invasive squamous cell carcinoma. *Vet. Dermatol.* **22**: 68–74. [[Medline](#)] [[CrossRef](#)]
20. Stebbins, K. E., Morse, C. C. and Goldschmidt, M. H. 1989. Feline oral neoplasia: a ten-year survey. *Vet. Pathol.* **26**: 121–128. [[Medline](#)] [[CrossRef](#)]
21. Tachezy, R., Duson, G., Rector, A., Jenson, A. B., Sundberg, J. P. and Van Ranst, M. 2002. Cloning and genomic characterization of *Felis domesticus* papillomavirus type 1. *Virology* **301**: 313–321. [[Medline](#)] [[CrossRef](#)]
22. Thomson, N. A., Dunowska, M. and Munday, J. S. 2015. The use of quantitative PCR to detect *Felis catus* papillomavirus type 2 DNA from a high proportion of queens and their kittens. *Vet. Microbiol.* **175**: 211–217. [[Medline](#)] [[CrossRef](#)]
23. Tieben, L. M., Berkhout, R. J., Smits, H. L., Bouwes Bavinck, J. N., Vermeer, B. J., Bruijn, J. A., Van der Woude, F. J. and Ter Schegget, J. 1994. Detection of epidermodysplasia verruciformis-like human papillomavirus types in malignant and premalignant skin lesions of renal transplant recipients. *Br. J. Dermatol.* **131**: 226–230. [[Medline](#)] [[CrossRef](#)]