NOTE

Wildlife Science

Serologic survey of *Brucella* infection in cetaceans inhabiting along the coast of Japan

Kazue OHISHI^{1)*}, Masao AMANO²⁾, Ken NAKAMATSU³⁾, Nobuyuki MIYAZAKI³⁾, Yuko TAJIMA⁴⁾, Tadasu K. YAMADA⁴⁾, Ayaka MATSUDA⁵⁾, Mari OCHIAI⁶⁾, Takashi F. MATSUISHI^{5,7)}, Hajime TARU⁸⁾, Hajime IWAO⁹⁾ and Tadashi MARUYAMA¹⁰⁾

¹⁾Faculty of Engineering, Tokyo Polytechnic University, 1583 Iiyama, Atsugi, Kanagawa 243-0297, Japan

²⁾Graduate School of Fisheries and Environmental Sciences, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan

³⁾Atmosphere and Ocean Research Institute, The University of Tokyo, 5-1-5 Kashiwa, Chiba 277-8564, Japan

⁴⁾National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan

⁵⁾Faculty of Fisheries Sciences, Hokkaido University, 3-1-1 Minato, Hakodate, Hokkaido 041-8611, Japan

⁶⁾Center for Marine Environmental Studies (CMES), Ehime University, 2-5 Bunkyo-cho, Matsuyama, Ehime 790-8577, Japan

⁷⁾Global Institution for Collaborative Research and Education, Hokkaido University, 3-1-1 Minato, Hakodate, Hokkaido 041-8611, Japan

⁸⁾Kanagawa Prefectural Museum of Natural History, 499 Iryuda, Odawara, Kanagawa 250-0031, Japan

⁹⁾Niigata City Aquarium, 5932-445 Nishifunami, Chuo-ku, Niigata, Niigata 951-8101, Japan

¹⁰⁾School of Marine Biosciences, Kitasato University, 1-15-1 Kitazato, Minami, Sagami-hara, Kanagawa 252-0373, Japan

J. Vet. Med. Sci.
82(1): 43–46, 2020
doi: 10.1292/jvms.19-0481

Received: 28 August 2019
Accepted: 5 November 2019
Advanced Epub:
18 November 2019

ABSTRACT. A serologic investigation of *Brucella* infection was performed in 7 species of cetaceans inhabiting along the coast of Japan. A total of 32 serum samples were examined by enzyme-linked immunosorbent assay (ELISA) using *Brucella abortus* and *B. canis* antigens. One serum sample from five melon-headed whales (*Peponocephala electra*) was positive for *B. abortus*. No serum sample showed positive for *B. canis*. The ELISA-positive melon-headed whale serum demonstrated a strong band appearance only against *B. abortus* antigens in Western blot analysis. Many detected bands were discrete, while some of them had a smeared appearance. The present results indicate that *Brucella* infection occurred in melon-headed whale population and the bacterial antigenicity is more similar to that of *B. abortus* than *B. canis*.

KEY WORDS: antibody, *Brucella*, cetacean, whale

Brucella, Gram-negative intracellular bacteria, infect variety of mammals and induce reproductive disorders including abortion [5]. In marine mammals, *B. ceti*, which is preferentially associated with cetaceans, and *B. pinnipedialis*, which is preferentially associated with pinnipeds, have been identified [7]. Compared to terrestrial *Brucella* species, these marine *Brucella* species seem to induce less pathological changes. However, in *B. ceti* infection, there have been substantial number of pathological reports including abnormalities in reproductive organs or nervous system [8, 9]. The abortion-inducing potential of *B. ceti* has been shown in captive and stranded dolphins and porpoises [6, 10, 13].

Extensive microbiological and serologic investigations have been performed on animals mainly from European and American waters, and revealed that many marine mammal species were infected with *Brucella* [8, 9]. In the western North Pacific, long-term and large-scale epidemiological surveys have shown that *Brucella* infection occurred in three species of baleen whales; common minke whales (*Balaenoptera acutorostrata*), Bryde's whales (*B. brydei*), and sei whales (*B. borealis*), and sperm whales (*Physeter macrocephalus*) [17, 20]. Especially, high prevalence of anti-*Brucella* antibodies and granulomatous testes were observed in common minke whales [17, 20]. Serologic evidence of *Brucella* has been found in two pygmy sperm whales (*Kogia breviceps*) stranded on the Japanese coast of the Pacific Ocean [18]. However, the available information is still limited in the Asian waters. A serologic survey was conducted to investigate the status of *Brucella* infection around Japan.

The serum samples used in the present study were summarized in Table 1 and Fig. 1. The blood samples were collected from cetaceans stranded and incidentally caught by a setnet, according to the guideline of the Japanese Fisheries Agency. The blood

*Correspondence to: Ohishi, K.: cie20910@syd.odn.ne.jp

©2020 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License. (CC-BY-NC-ND 4.0: <https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Table 1. Prevalence of serum antibodies to *Brucella* in cetaceans inhabiting the coast of Japan

Species	Year	Location				Positivity rate ^{b)}	
		Prefecture	City	Town	Map site ^{a)}	<i>B. abortus</i>	<i>B. canis</i>
Dall's porpoises (<i>Phocoenoides dalli</i>)	2001			Off Iwate Prefecture	1	0/20	0/20
Melon-headed whales (<i>Peponocephala electra</i>)	2001–2002	Ibaraki	Kamisu	Hasaki	2	1/5	0/5
Harbor porpoise (<i>Phocoena phocoena</i>)	2017	Hokkaido	Hakodate	Usujiri	3	0/3	0/3
Pacific white-sided dolphin (<i>Lagenorhynchus obliquidens</i>)	2017	Hokkaido	Hakodate	Shiokubi	4	0/1	0/1
Minke whale (<i>Balaenoptera acutorostrata</i>)	2001	Kanagawa	Fujisawa	Enoshima	5	0/1	0/1
Hubbs' beaked whale (<i>Mesoplodon carlhubbsi</i>)	2005	Kanagawa	Ninomiya	Yamanishi	6	0/1	0/1
Dwarf sperm whale (<i>Kogia sima</i>)	2007	Niigata	Niigata	Matsuhaminato	7	0/1	0/1

a) Numbers indicate the site in Fig. 1. b) Positive individuals /examined individuals. Absorbance greater than 0.2 at 405 nm is regarded as positive.

samples of Dall's porpoises (*Phocoenoides dalli*) captured by a commercial fishery were also collected. Three harbor porpoises (*Phocoena phocoena*) rescued from a setnet were kept according to the guidelines for regulations on animal experimentation of Hokkaido University and were released after a medical care.

Anti-*Brucella* serum antibody was detected in the enzyme-linked immunosorbent assay (ELISA) and in the Western blot analysis according to the protocol described previously [1, 17]. Briefly, commercially available inactivated *B. abortus* strain 125 (Kaketsuken Co., Kumamoto, Japan) and *B. canis* strain QE-13B (Kitasato Institute Co., Tokyo, Japan) were solubilized and used as antigens in both methods. The sera diluted to 1:100 and horseradish peroxidase-conjugated Protein A/G (Thermo Fisher Scientific Inc., Waltham, MA, USA) diluted to 1:5,000, were used for the primary antibody and for its detection in both methods. In the ELISA, the absorbance value higher than 0.2 at 405 nm, was regarded as positive [1, 16].

One serum sample from a female melon-headed whale (*Peponocephala electra*) (ID: PE57) showed a positive-value (OD=0.34) only for *B. abortus* antigens. However, none of the serum samples showed positive to *B. canis* antigens (Table 1). In Western blot analysis, the ELISA-positive melon-headed whale serum showed a strong response only to *B. abortus* (Fig. 2). Many bands, including the smeary bands, were detected against *B. abortus*, while no band against *B. canis* was detected at this serum concentration (Fig. 2).

The present results indicated that the *Brucella* infection occurred in melon-headed whale population, and the *Brucella* strain

Fig. 1. Sampling sites. Numbers in the map correspond to those of locations in Table 1.

Fig. 2. Western blot analysis of the serum from an ELISA-positive melon-headed whale (ID: PE57) against *Brucella abortus* and *B. canis* antigens. A, *B. abortus* antigen. B, *B. canis* antigen.

was antigenetically closer to *B. abortus* than *B. canis*. The appearance of the smeared bands suggests that the infected *Brucella* are smooth (S) colony type of bacteria containing large O-type polysaccharide in the outer membrane as well as *B. abortus* [23]. These serum responses are similar to those of three species of baleen whales and sperm whales in the western North Pacific, and beluga whales in the Anadyr Firth, Russia [16, 17]. This suggests that S colony type of *Brucella* are prevalent in at least several cetacean species in the western North Pacific.

Melon-headed whales are distributed in the tropical and subtropical waters, and several mass stranding events of melon-headed whales have been recorded on multiple coasts of Japan [2, 12]. In Aoshima town in Miyazaki prefecture, the mass stranding with 135 individuals was recorded for the first time in 1982 [14, 15]. More recently, 171 whales at Tanegashima Island in Kagoshima prefecture and 85 whales at Hasaki town in Ibaraki prefecture, stranded in 2001 and 2002, respectively [4]. Comparison of these animals in term of animals' body size suggests that the melon-headed whales found in various parts of Japan are composed of several geographically different regional populations [4]. It would be interesting to compare the antibody positivity among these regional populations of the species, because maternal transmission is considered to be a major route of *Brucella* in cetaceans [17]. We have previously reported the serologic evidence of *Brucella* infection in two pygmy sperm whales which also inhabit in the tropical and subtropical waters [18, 22]. Because the investigation on *Brucella* infection has been rarely conducted in these waters, it is important to accumulate the epidemiological information through surveillance in Asian waters.

No specific antibodies were found in the examined 20 Dall's porpoises. It is a contrast to common minke whales inhabiting the western North Pacific with a high prevalence of the antibodies (9–38% during 1994–2010), though the habitats of the two species are partially overlapped [3, 17, 19]. Similarly, serologically *Brucella* positive or negative whale species inhabit in the western South Hemisphere. Bottlenose dolphins (*Tursiops truncatus*) inhabiting around the Solomon islands have shown high prevalence of anti-*Brucella* antibodies in their sera [21]. However, Antarctic minke whales (*Balaenoptera bonarensis*) migrating the western South Pacific seem to be *Brucella*-free, because specific antibodies have not been detected in the examined 739 serum samples [11, 17]. These facts may indicate that environmental transmission of *Brucella* is less likely to occur in cetaceans. The present serum samples from Dall's porpoises were collected off Iwate prefecture in 2001, before being seriously damaged by the 2011 Great East Japan Earthquake. The present data would provide a serologic starting line for ongoing or future researches of the species after the disaster. Further studies not only by serology but also by bacterial isolation should be performed in the future. The microbiological and molecular biological characterizations are necessary for understanding the distribution of *Brucella* and its impact on cetaceans in Asian waters.

ACKNOWLEDGMENTS. The authors would like to thank Dr. T. Kuramochi of National Museum of Nature and Science, the staffs of Stranding Network Hokkaido, and many persons involved in the stranding field work for collecting samples. We also thank Mr. H. Shouzushima for a kind cooperation in blood sampling from animals obtained in the commercially fishery. Authors express the thanks to Dr. T. Bando of Institute of Cetacean Research for the earnest discussion.

REFERENCES

1. Abe, E., Ohishi, K., Ishinazaka, T., Fujii, K. and Maruyama, T. 2017. Serologic evidence of *Brucella* infection in pinnipeds along the coast of Hokkaido, the northernmost main island of Japan. *Microbiol. Immunol.* **61**: 114–122. [Medline] [CrossRef]
2. Amano, M. 2015. *Peponocephala electra* (Gray, 1846). pp. 382–383. In: The Wild Mammals of Japan (Ohdachi, S. D., Ishibashi, Y., Iwasa, M. A. and Saitoh, T. eds.), Shoukado, Kyoto.
3. Amano, M. and Hayano, A. 2007. Intermingling of dalli-type Dall's porpoises into a wintering truei-type population off Japan: implications from color patterns. *Mar. Mamm. Sci.* **23**: 1–14. [CrossRef]
4. Amano, M., Yamada, T. K., Kuramochi, T., Hayano, A. A., Kazumi, A. and Sakai, T. 2014. Life history and group composition of melon-headed whales based on mass strandings in Japan. *Mar. Mamm. Sci.* **30**: 480–493. [CrossRef]
5. Corbel, M. J. and Banai, M. B. 2005. Genus *Brucella*. pp. 370–386. In: Bergey's Manual of Systematic Bacteriology. Vol. 2 (Brenner, D. J., Kreig, N. R. and Staley, J. T. eds.), Springer Press, New York.
6. Ewalt, D. R., Payeur, J. B., Martin, B. M., Cummins, D. R. and Miller, W. G. 1994. Characteristics of a *Brucella* species from a bottlenose dolphin (*Tursiops truncatus*). *J. Vet. Diagn. Invest.* **6**: 448–452. [Medline] [CrossRef]
7. Foster, G., Osterman, B. S., Godfroid, J., Jacques, I. and Cloeckaert, A. 2007. *Brucella ceti* sp. nov. and *Brucella pinnipedialis* sp. nov. for *Brucella* strains with cetaceans and seals as their preferred hosts. *Int. J. Syst. Evol. Microbiol.* **57**: 2688–2693. [Medline] [CrossRef]
8. Guzmán-Verri, C., González-Barrientos, R., Hernández-Mora, G., Morales, J. A., Baquero-Calvo, E., Chaves-Olarte, E. and Moreno, E. 2012. *Brucella ceti* and brucellosis in cetaceans. *Front. Cell. Infect. Microbiol.* **2**: 3. [Medline] [CrossRef]
9. Hernández-Mora, G., Palacios-Alfaro, J. D. and González-Barrientos, R. 2013. Wildlife reservoirs of brucellosis: *Brucella* in aquatic environments. *Rev. Off. Int. Epizoot.* **32**: 89–103. [Medline] [CrossRef]
10. Jauniaux, T. P., Brenez, C., Fretin, D., Godfroid, J., Haelters, J., Jacques, T., Kerckhof, F., Mast, J., Sarlet, M. and Coignoul, F. L. 2010. *Brucella ceti* infection in harbor porpoise (*Phocoena phocoena*). *Emerg. Infect. Dis.* **16**: 1966–1968. [Medline] [CrossRef]
11. Kasamatsu, F., Nishiwaki, S. and Ishikawa, H. 1995. Breeding areas and southbound migrations of southern minke whales *Balaenoptera acutorostrata*. *Mar. Ecol. Prog. Ser.* **119**: 1–10. [CrossRef]
12. Marine Mammals Information DATABASE by National Museum of Nature and Science. <https://www.kahaku.go.jp/english/research/db/zoology/mammam/index.php> [accessed on October 1, 2019].
13. Miller, W. G., Adams, L. G., Ficht, T. A., Cheville, N. F., Payeur, J. P., Harley, D. R., House, C. and Ridgway, S. H. 1999. *Brucella*-induced abortions and infection in bottlenose dolphins (*Tursiops truncatus*). *J. Zoo Wildl. Med.* **30**: 100–110. [Medline]
14. Miyazaki, N. 1983. A mass stranding of *Peponocephala electra*. *Animal and Nature* **13**: 2–9 (in Japanese).
15. Miyazaki, N., Fujise, Y. and Iwata, K. 1998. Biological analysis of a mass stranding of melon-headed whales (*Peponocephala electra*) at Aoshima,

- Japan. *Bull Natr. Sci. Tokyo Ser. A* **24**: 31–60.
16. Ohishi, K., Abe, E., Amano, M., Miyazaki, N., Boltunov, A., Katsumata, E. and Maruyama, T. 2018. Detection of serum antibodies to *Brucella* in Russian aquatic mammals. *J. Vet. Med. Sci.* **80**: 1696–1701. [[Medline](#)] [[CrossRef](#)]
17. Ohishi, K., Bando, T., Abe, E., Kawai, Y., Fujise, Y. and Maruyama, T. 2016. Long-term and large-scale epidemiology of *Brucella* infection in baleen whales and sperm whales in the western North Pacific and Antarctic Oceans. *J. Vet. Med. Sci.* **78**: 1457–1464. [[Medline](#)] [[CrossRef](#)]
18. Ohishi, K., Katsumata, E., Uchida, K. and Maruyama, T. 2007. Two stranded pygmy sperm whales (*Kogia breviceps*) with anti-*Brucella* antibodies in Japan. *Vet. Rec.* **160**: 628–629. [[Medline](#)] [[CrossRef](#)]
19. Ohishi, K., Takishita, K., Kawato, M., Zenitani, R., Bando, T., Fujise, Y., Goto, Y., Yamamoto, S. and Maruyama, T. 2004. Molecular evidence of new variant *Brucella* in North Pacific common minke whales. *Microbes Infect.* **6**: 1199–1204. [[Medline](#)] [[CrossRef](#)]
20. Ohishi, K., Zenitani, R., Bando, T., Goto, Y., Uchida, K., Maruyama, T., Yamamoto, S., Miyazaki, N. and Fujise, Y. 2003. Pathological and serological evidence of *Brucella*-infection in baleen whales (*Mysticeti*) in the western North Pacific. *Comp. Immunol. Microbiol. Infect. Dis.* **26**: 125–136. [[Medline](#)] [[CrossRef](#)]
21. Tachibana, M., Watanabe, K., Kim, S., Omata, Y., Murata, K., Hammond, T. and Watarai, M. 2006. Antibodies to *Brucella* spp. in Pacific bottlenose dolphins from the Solomon Islands. *J. Wildl. Dis.* **42**: 412–414. [[Medline](#)] [[CrossRef](#)]
22. Yoshioka, M. 2015. *Kogia breviceps* (Blainville, 1838). pp. 346–347. In: The Wild Mammals of Japan (Ohdachi, S. D., Ishibashi, Y., Iwasa, M. A. and Saitoh, T. eds.), Shoukado, Kyoto.
23. Zygmunt, M. S., Blasco, J. M., Letesson, J. J., Cloeckert, A. and Moriyón, I. 2009. DNA polymorphism analysis of *Brucella* lipopolysaccharide genes reveals marked differences in O-polysaccharide biosynthetic genes between smooth and rough *Brucella* species and novel species-specific markers. *BMC Microbiol.* **9**: 92. [[Medline](#)] [[CrossRef](#)]