Advance Publication

The Journal of Veterinary Medical Science

Accepted Date: 27 July 2021

J-STAGE Advance Published Date: 18 September 2021

©2021 The Japanese Society of Veterinary Science

Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

1 ***[Note] Surgery***

2 **Retrospective study on the outcomes and risk factors of right paramedian**
3 **abomasopexy for right abomasal disorders in 47 dairy cows**

4
5 Kenji TSUKANO¹⁾, Yasunobu NISHI²⁾, Naoki SASAKI²⁾ and Kazuyuki SUZUKI³⁾*

6
7 1) *Minami-Hokkaido Agricultural Mutual Aid Association, 74-2 Higashimae, Hokuto,*
8 *Hokkaido 041-1214, Japan*

9 2) *Joint Faculty of Veterinary Medicine, Yamaguchi University, 1677-1 Yoshida,*
10 *Yamaguchi, Yamaguchi 753-8515, Japan*

11 3) *School of Veterinary Medicine, Rakuno Gakuen University, 582 Midorimati,*
12 *Bunnkyoudai, Ebetsu, Hokkaido 069-8501, Japan*

13
14
15
16 **Correspondence to:** Kazuyuki SUZUKI, DVM, Ph.D.

17 *School of Veterinary Medicine, Rakuno Gakuen University, 582 Midorimachi,*
18 *Bunnkyoudai, Ebetsu, Hokkaido, 069-8501, JAPAN*

19 E-mail: kazuyuki@rakuno.ac.jp

20 Phone and Fax: +81-11-388-4702

21
22
23 **RUNNING HEAD**

24 **RIGHT ABOMASAL DISORDERS IN COWS**

ABSTRACT

Very few epidemiologic studies have verified the utility of the right paramedian abomasopexy (RPA) technique in cows with right abomasal disorders. This study aimed to investigate the outcomes and risk factors for non-survival in the herd within 30 days of surgery in cows with right abomasal disorders who underwent the RPA technique. Forty-seven Holstein cows with right abomasal disorders (25 with right abomasal displacement [RDA] and 22 with right abomasal volvulus [RAV]) were included. Twenty-two cows with RDA (22/25, 88.0%) and 10 cows with RAV (10/22, 45.5%) survived at 30 days post-surgery. Multivariate logistic regression analysis indicated that hyponatremia, hypokalemia, and the presence of abomasal volvulus were the major risk factors associated with non-survival.

KEY WORDS:

cow, mortality, right abomasal displacement, right abomasal volvulus, right paramedian abomasopexy

40 Since the first report of displacement of the abomasum (DA) in 1950 [3], this
41 disorder has become more common in dairy cows. A DA is an abnormal positioning of
42 the abomasum within the abdominal cavity and is classified into three broad categories:
43 left abomasal displacement (LDA), right abomasal displacement (RDA), and right
44 abomasal volvulus (RAV) [21]. In general, cows with simple displacements have good
45 prognosis for returning to a productive life after surgical replacement [2, 13, 20, 23, 24].
46 However, since RAV is an emergency condition with severe hemodynamic compromise
47 [6, 14], cows with RAV have poorer prognosis than cows with simple displacement [20].
48 The goals of treatment for DA are to return the abomasum to its normal position and to
49 prevent the recurrence of displacement.

50 Surgery is known to be the only correction method that ensures satisfactory
51 replacement and prevents recurrence [19]. Many surgical techniques have been employed,
52 including right paralumbar fossa omentopexy (RPFO) [15], left paralumbar fossa
53 abomasopexy [16], left paralumbar fossa omentopexy [25], right paramedian

54 abomasopexy (RPA) [13], toggle pin (bar) suture [17], and 1-step [1] or 2-step [11]
55 laparoscopic abomasopexy. In Japan, RPFO and RPA are the most widely used surgical
56 techniques for cows with DA. Although there are specific indications for RPFO and RPA,
57 which procedure is more advantageous remains controversial. It is well known that in
58 cows with LDA, there is no association between the surgical procedure and longevity
59 [21]. For instance, Fubini *et al.* [13] demonstrated that there was no difference between
60 RPFO and RPA in the rate of a negative outcome within 30 days in cows with LDA.
61 However, the RPA technique is not recommended as the first option for right abomasal
62 disorders [12], because of the difficulty in identifying and manipulating the viscera caused
63 by position in dorsal recumbency. Notably, the relationship of structures is less familiar
64 to most veterinarians using this approach [12]. Indeed, few reports have revealed surgical
65 outcomes with the RPA technique in cows with right abomasal disorders. However, as
66 abomasopexy enables long-lasting correction of displacement [19], the RPA technique
67 has an advantage from the perspective of DA recurrence. Furthermore, the proximity of

the abomasum to the incision makes this approach possible for the smallest surgeon and the largest cow [12]. Thus, the usefulness of the RPA technique in cows with right abomasal disorders should be considered.

To the best of our knowledge, there is very little epidemiologic research to verify the utility of the RPA technique in cows with right abomasal disorders. Thus, the aim of the present study was to investigate the outcomes and identify risk factors for non-survival in cows with right abomasal disorders who underwent the RPA technique. Our results may be useful for veterinary practitioners when selecting surgical techniques for right abomasal disorders.

A retrospective study was conducted using medical records of Holstein cows that underwent surgical treatment for right abomasal disorders at the Minami Hokkaido Agricultural Mutual Aid Association located in northern Japan (140°E, 42°N) between December 2014 and December 2020. The criteria for selection were as follows: 1) surgical treatment was performed to correct RDA or RAV with the RPA technique on the

day of diagnosis; 2) blood analysis results for the day of surgery were available; and 3) the cows had no history of surgical correction of abomasal disorders. Cows presenting with astasia were excluded. Consequently, 47 cases of right abomasal disorders (25 cases of RDA and 22 cases of RAV) were identified. The medical records of these cases were derived from a specific veterinary practice that was equipped with an operating table facility to retain the cow in a dorsal position. In this veterinary practice, surgical correction of DA was performed using the RPA technique in all cases; therefore, surgeons in the veterinary practice were accustomed to this approach. From medical records, data on supportive medical treatments during the operative period, clinical experience of surgeons, and presence or absence of complications were obtained. Furthermore, we investigated the prognosis of cows with right abomasal disorders that underwent the RPA technique; cows were defined as non-surviving if they died or were culled within 30 days of surgery.

All cows underwent a blood analysis immediately before surgery, including

blood sodium (Na), potassium (K), chloride (Cl), hemoglobin (Hb), hematocrit (Ht), and anion gap (AG) values, using an automatic gas analyzer (i-STAT 1, Abbott Lab, Princeton, IL, USA) and i-STAT cartridge (i-STAT CHEM8+ Cartridge, Abbott Lab).

The RPA technique used in this study was as follows. The cows were placed in dorsal recumbence on an operating table. All cows were injected with 50 ml of 2% lidocaine hydrochloride (lidocaine injection 2%, Pfizer Co., Ltd., Tokyo, Japan) at the incision area as a local anesthetic. The skin was incised 10 cm caudal to the sternum and 10 cm to the right of the midline. The surgical incision was 15 cm in length. The abomasum was decompressed if necessary and placed in the correct anatomical position. If correction is not obtained with gas decompression alone owing to volvulus and/or large abomasal fluid volume, the cow's position was changed from dorsal to left lateral recumbency and the pylorus was then pulled out from the incision to expose the body of the abomasum. Abomasotomy was performed in cases with a large amount of fluid in the abomasum. A 6-cm portion of the greater curvature of the abomasum

(approximately 10 cm from the reticulo-abomasal junction and 4 cm from the attachment of the greater omentum) was incorporated into the closure of the USP 5 silk sutures (Zyuuiyou Surgical Silk, Kawasaki-Seibutu Kagakukenkyusho Co., Ltd., Tokyo, Japan) in a simple continuous pattern. Care was taken to avoid penetrating the abomasum mucosa. The peritoneum and rectus muscle were closed with 0.6-mm braided absorbable sutures (Polysorb™, Covidien Japan Co., Ltd., Tokyo, Japan) in a simple continuous pattern, and the subcutaneous tissues were closed with the same 0.6-mm braided absorbable sutures in a Ford interlocking pattern. The skin was also closed with the same 0.6-mm braided absorbable sutures in an intradermal suture pattern.

Statistical analysis was performed using Excel Toukei 2010 (SSRI, Osaka, Japan). Normally distributed data are reported as the mean $\pm$ standard deviation (SD), and non-normally distributed data are expressed as the median (minimum–maximum). Cows were divided into two groups based on the survival period (up to 30 days) from the surgical treatment (survival group or non-survival group). Student's *t*-test or Mann-

Whitney *U*-test after the *F*-test was used for comparisons of blood data between groups.

If there were significant differences in blood analysis values between groups, those values were categorized for multivariate logistic regression analysis based on the reference values: Na (133–139 mM [8]), K (3.6–4.4 mM [8]), Cl (94.7–103.3 mM [8]), Hb (8.3–10.5 g/dl [22]), Ht (26.1–33.5 % [22]) and AG (14–20 mM [7]). Thereafter, the multivariate association between categorized variables (blood analysis values with differences between groups (if any), presence of abomasal volvulus, clinical experience of surgeons [23], and presence of complications] and outcomes of therapy were evaluated by logistic regression analysis with the calculation of odds ratios (ORs) and associated 95% confidence intervals (95% CI). In addition, the overall survival time (OST) was calculated as the time between surgical treatment and non-survival. The OST curves were estimated using Kaplan-Meier statistics for the different groups with or without risk factors for non-survival. The OST was compared between groups using the log-rank test.

The significance level was set at $p < 0.05$.

138 The age, parity, and days postpartum in cows enrolled in this survey were 4 (2–
139 11) years, 3 (1–9), and 34 (2–337) days, respectively. Diagnosis and surgical corrections
140 were performed by 14 different surgeons, who had 5 (2–33) years of clinical experience.
141 The median number of cases operated per surgeon was 3 (1–9). Abomasotomy was
142 performed to excrete the fluid of the abomasum in 1 cow in the survival group and in 4
143 cows in the non-survival group. Supportive medical treatments mostly consist of
144 antimicrobial and intravenous fluid therapies. During the operative period, antibiotics
145 were administered to all cows. Although the total intravenous fluid volume was
146 significantly higher in the non-survival group than in the survival group (survival vs. non-
147 survival: 6.3 [0.0–16.5] vs. 8.0 [2.0–18.0] l, $p = 0.003$), there were no significant
148 differences in the volume of hypertonic saline solution, an intravenous fluid solution used
149 to correct severe electrolyte imbalances and dehydration in cows, during the operative
150 period (survival vs. non-survival: 0.5 [0.0–3.0] vs. 2.0 [0.0–3.0] l, $p = 0.717$). In this
151 survey, 17 of 47 (36.2%) cases had complications: ketosis ($n = 8$, 17.0%), milk fever (n

= 5, 10.6%), mastitis (n = 3, 6.4%), and uterine disease (n = 1, 2.1%). Therefore, the veterinary practitioner also treated complications before and after repositioning surgery until the cows were discharged in a healthy state.

In this survey, 22 cows with RDA (22/25, 88.0%) and 10 cows with RAV (10/22, 45.5%) survived at 30 days from the day of surgical treatment. Therefore, the survival rate of the 47 patients in this survey was 68.1% (32/47). In our survey, 93.3% (14/15) of cows in the non-survival group died within 21 days without recovery after surgery, and the remaining cows were culled 24 days after surgery because the producer was not satisfied with her performance. Previous studies reported a survival rate for right flank laparotomy, such as RPFO and right paralumbar fossa pyloropexy, in cows with RDA and RAV ranging from 85.0% to 95.5% and from 35.0% to 60.0%, respectively, within 30 days following surgery [4, 5, 10]. In this survey, the survival rate within 30 days in cows who underwent the RPA technique was 88.0% for RDA and 45.5% for RAV, which were in good agreement with those of right flank laparotomy. Therefore, the present survey

indicated that RPA techniques are not inferior to right flank laparotomy based on the survival rate in the surgical correction of right abomasal disorders.

We also investigated the major risk factors associated with non-survival in cows with right abomasal disorders that underwent the RPA technique. The blood Na (survival vs. non-survival: 137.2 ± 3.9 vs. 132.4 ± 6.3 mM, $p = 0.003$), K (survival vs. non-survival: 3.5 ± 0.6 vs. 3.0 ± 0.6 mM, $p = 0.005$) and Cl (survival vs. non-survival: 93.8 ± 9.7 vs. 83.9 ± 9.0 mM, $p = 0.002$) concentrations were significantly lower in the non-survival group than in the survival group. The blood AG value was significantly higher in the non-survival group than in the survival group (survival vs. non-survival: 16.1 ± 2.4 vs. 17.9 ± 2.8 mM, $p = 0.032$). Although a significant difference was observed in AG values between the groups ($p = 0.032$), these values remained within the normal range. There were no significant difference in Hb concentration (survival vs. non-survival: 10.7 ± 1.7 vs. 11.8 ± 2.5 g/dl, $p = 0.149$) and Ht (survival vs. non-survival: 31.7 ± 4.5 % vs. 34.5 ± 7.4 %, $p = 0.162$) between the groups. The blood Na, K, and Cl concentrations, abomasal volvulus,

clinical experience of surgeons, and complications were categorized using multivariate logistic regression analysis (Table 1). Multivariate logistic regression analysis indicated that hyponatremia (<133.0 mM, OR 38.20; 95%CI 1.66–879.00; $p = 0.022$), hypokalemia (<3.6 mM, OR 40.60; 95% CI 1.31–1260.00; $p = 0.034$), and presence of abomasal volvulus (OR 10.60; 95% CI 1.05–107.00; $p = 0.045$) were the major risk factors associated with non-survival in cows with right abomasal disorders. However, no differences were found between groups in blood Cl concentrations, clinical experience of surgeons, and the presence of complications.

Figure 1 shows the OST from the day of surgical treatment for cows with right abomasal disorders with or without risk factors for non-survival. At the end of the survey, at 30 days from the day of surgical treatment, all cows ($n = 13$, 100%) without risk factors survived. However, at the end of the survey, 17 of 34 (55.9%) cases indicated that hyponatremia, hypokalemia, and the presence of abomasal volvulus were the major risk factors for mortality. Therefore, based on the survival estimated by Kaplan-Meier analysis,

cows with hyponatremia, hypokalemia, and/or abomasal volvulus had a significantly shorter survival ($p = 0.006$) than those without.

It is well known that since RAV is an emergency condition with severe hemodynamic compromise; cows with RAV have poor prognosis [6, 14]. Indeed, in this survey, the total intravenous fluid volume, which was determined by the degree of dehydration, was significantly higher in the non-survival group than in the survival group during the operative period, indicating that the non-survival group had more severe hemodynamic compromise. Previous studies have revealed that hyponatremia [10, 14], hypokalemia [20], and abomasal volvulus [4, 5, 10] were observed in cows with right abomasal disorders that underwent non-RPA techniques including RPFO [10] with non-survival. Our results indicated that the surgical outcome of the RPA technique was affected by preoperative conditions, such as severe hemodynamic compromise and electrolyte abnormality, similar to that of non-RPA techniques [4, 5, 10, 14, 20].

Generally, because it is believed that the surgical outcome depends on the skill

208 of the surgeon, the clinical experience of the surgeon is a factor that requires consideration.
209 In this survey, surgical corrections were made by 14 different surgeons, who had 5 (2–33)
210 years of clinical experience; however, all surgeons were accustomed to the RPA approach,
211 which is less familiar to most veterinarians [12]. There was no association between the
212 experience of surgeons and longevity in the herd. Therefore, the effects of surgeon skill
213 on the surgical outcomes examined in the present survey were minimal.

214 Kalaitzakis *et al.* [18] reported that the non-survival of cows with LDA who
215 underwent repositioning surgery was related to fatty liver disease. Moreover, ketosis and
216 retained placenta have been reported to be risk factors for herds with a high prevalence
217 of postpartum culling [9]. Since previous studies indicated that concurrent postpartum
218 disorders affect prognosis after LDA surgery, the presence of complications was forced
219 into the logistic regression analysis model in this survey. However, our survey revealed
220 that the presence of complications was not involved in the prognosis of cows with right
221 abomasal disorders that underwent the RPA technique.

222 Since the survival rate and risk factors in the RPA technique are similar to those
223 of the non-RPA technique reported previously, we concluded that RPA techniques are not
224 inferior to non-RPA techniques that are popular among most veterinarians for surgical
225 correction of right abomasal disorders. Our survey showed the usefulness of the RPA
226 technique in cows with right abomasal disorders. Therefore, veterinarian practitioners can
227 select the RPA technique depending on the preference of surgeons or the condition of
228 cows with right abomasal disorders.

229

230 Conflicts of interest

231 The authors have no conflicts of interest directly relevant to the content of this article.

References

1. Babkine, M., Desrochers, A., Bouré, L. and Hélie, P. 2006. Ventral laparoscopic abomasopexy on adult cows. *Can. Vet. J.* **47**:343-348.
2. Baird, A. N., Mohamed, A., Moore, G. E. and Hawkins, J. F. 2017. Comparison of omentopexy versus pyloro-omentopexy for treatment of left abomasal displacement in dairy cows: 87 cases (2001-2005). *J. Am. Vet. Med. Assoc.* **251**: 1182-1187.
3. Begg, H. 1950. Diseases of the stomach of the adult ruminant. *Vet. Rec.* **62**: 797-808.
4. Boulay, G., Francoz, D., Doré, E., Dufour, S., Veillette, M., Badillo, M., Bélanger, A. M. and Buczinski, S. 2014. Preoperative cow-side lactatemia measurement predicts negative outcome in Holstein dairy cattle with right abomasal disorders. *J. Dairy Sci.* **97**: 212-221.
5. Buczinski, S., Boulay, G. and Francoz, D. 2015. Preoperative and postoperative L-lactatemia assessment for the prognosis of right abomasal disorders in dairy cattle. *J. Vet. Intern. Med.* **29**: 375-380.

- 246 6. Constable, P. D., St Jean, G., Hull, B. L., Rings, D. M., Hoffsis, G. F. 1991. Prognostic
247 value of surgical and postoperative findings in cattle with abomasal volvulus. *J. Am.*
248 *Vet. Med. Assoc.* **199**: 892-898.
- 249 7. Constable, P. D., Streeter, R. N., Koenig, G. J., Perkins, N. R., Gohar, H. M. and Morin,
250 D. E. 1997. Determinants and utility of the anion gap in predicting hyperlactatemia in
251 cattle. *J. Vet. Intern. Med.* **11**: 71-79.
- 252 8. Cozzi G, Ravarotto L, Gottardo F, Stefani AL, Contiero B, Moro L, Brscic M, Dalvit
253 P. 2011. Short communication: reference values for blood parameters in Holstein
254 dairy cows: effects of parity, stage of lactation, and season of production. *J. Dairy Sci.*
255 **94**: 3895-3901.
- 256 9. Dubuc, J. and Denis-Robichaud, J. 2017. A dairy herd-level study of postpartum
257 diseases and their association with reproductive performance and culling. *J. Dairy Sci.*
258 **100**: 3068-3078.
- 259 10. Figueiredo, M. D., Nydam, D. V., Perkins, G. A., Mitchell, H. M. and Divers, T. J.

260 2006. Prognostic value of plasma L-lactate concentration measured cow-side with a
261 portable clinical analyzer in Holstein dairy cattle with abomasal disorders. *J. Vet.*
262 *Intern. Med.* **20**: 1463-1470.

263 11. Fiore, F., Musina, D., Cocco, R., Di Cerbo, A. and Spissu, N. 2018. Association
264 between left-displaced abomasum corrected with 2-step laparoscopic abomasopexy
265 and milk production in a commercial dairy farm in Italy. *Ir. Vet. J.* **71**:20

266 12. Fubini, S. L. and Ducharme, N. G. 2017. Farm Animal Surgery, second ed., pp. 223-
267 343, Elsevier, St. Louis, Missouri.

268 13. Fubini, S. L., Ducharme, N. G., Erb, H. N. and Sheils, R. L. 1992. A comparison in
269 101 dairy cows of right paralumbar fossa omentopexy and right paramedian
270 abomasopexy for treatment of left displacement of the abomasum. *Can. Vet. J.* **33**:
271 318-324.

272 14. Fubini, S. L., Gröhn, Y. T. and, Smith, D. F. 1991. Right displacement of the
273 abomasum and abomasal volvulus in dairy cows: 458 cases (1980-1987). *J. Am. Vet.*

- 274 *Med. Assoc.* **198**: 460-464.
- 275 15. Gabel, A. A., Heath, B. R. 1969. Correction and right-sided omentopexy in treatment
276 of left-sided displacement of the abomasums in dairy cattle. *J. Am. Vet. Med. Assoc.*
277 **155**: 632-641.
- 278 16. Gertsen, K. E. 1967. Surgical correction of the displaced abomasums. *Vet. Med. Sm.*
279 *Anim. Clin.* **62**: 679-682.
- 280 17. Grymer, J. and Sterner, K. E. 1982. Percutaneous fixation of left displaced
281 abomasums, using bar suture. *J. Am. Vet. Med. Assoc.* **180**: 1458-1461.
- 282 18. Kalaitzakis, E., Roubies, N., Panousis, N., Pourliotis, K., Kaldrymidou, E. and
283 Karatzias, H. 2006. Evaluation of ornithine carbamoyl transferase and other serum
284 and liver-derived analytes in diagnosis of fatty liver and postsurgical outcome of left-
285 displaced abomasum in dairy cows. *J. Am. Vet. Med. Assoc.* **229**: 1463-1471.
- 286 19. Mather, M. F. and Dedrick, R. S. 1966. Displacement of the abomasum. *Cornell Vet.*
287 **56**: 323-344.

288 20. Meylan, M. 1999. Prognostic indicators in cattle with right-sided displacement of the
289 abomasum and abomasal volvulus. *Schweiz. Arch. Tierheilkd.* **141**: 413-418. [Article
290 in German]

291 21. Niehaus, A. J. 2016. Surgical Management of Abomasal Disease. *Vet. Clin. North Am.*
292 *Food Anim. Pract.* **32**: 629-644.

293 22. Paiano, R. B., Birgel, D. B. and Birgel Junior, E. H. 2020. Influence of peripartum on
294 the erythrogram of Holstein dairy cows. *J. S. Afr. Vet. Assoc.* **91**: e1-e6.

295 23. Petty, R. D. 1981. Surgical correction of left displacement abomasums in cattle: a
296 retrospective study of 143 cases. *J. Am. Vet. Med. Assoc.* **178**: 1274-1276.

297 24. Reynen, J. L., Kelton, D. F., LeBlanc, S. J., Newby, N. C. and Duffield, T. F. 2015.
298 Factors associated with survival in the herd for dairy cows following surgery to
299 correct left displaced abomasum. *J. Dairy Sci.* **98**: 3806-3813.

300 25. Robertson, J. M. and Boucher, W. B. 1966. Treatment of left displacement of the
301 bovine abomasums. *J. Am. Vet. Med. Assoc.* **149**: 1423-1429.

302 **Figure Legend**

303 Figure 1

304 Graphs depicting the overall survival time (OST) from the initiation of surgical
305 treatment in cases with or without risk factors associated with non-survival by Kaplan-
306 Meyer analysis. The OST between groups was compared using the log-rank test. *; $p =$
307 0.006, vs. cases without risk factors.

Fig.1

Table 1. Risk factors for non-survival in surgical correction of right abomasal disorders with right paramedian abomasopexy

Variable	Variable category	n	Odds ratio	95% confidence interval	<i>P</i> -value
Sodium concentration	≥ 133.0 mM	36	1.00	-	-
	< 133.0 mM	11	38.20	1.66, 879.00	0.022
Potassium concentration	≥ 3.6 mM	19	1.00	-	-
	< 3.6 mM	28	40.60	1.31, 1260.00	0.034
Chloride concentration	≥ 94.7 mM	22	1.00	-	-
	< 94.7 mM	25	1.89	0.23, 15.80	0.556
Abomasal volvulus	No	25	1.00	-	-
	Yes	22	10.60	1.05, 107.00	0.045
Clinical experience of surgeon	More than 5 years	26	1.00	-	-
	Under 5 years	21	0.21	0.03, 1.69	0.143
Complications	No	30	1.00	-	-
	Yes	17	0.11	0.01, 1.31	0.081