Advance Publication

The Journal of Veterinary Medical Science

Accepted Date: 23 January 2022

J-STAGE Advance Published Date: 2 February 2022

©2022 The Japanese Society of Veterinary Science

Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

Parasitology

Note

Identification and phylogenetic analysis of *Babesia* parasites in domestic dogs in Nigeria

Haruyuki Hirata¹⁾, Temidayo Omobowale²⁾, Olufunke Adebayo³⁾, Nodoka Asanuma¹⁾, Asako Haraguchi⁴⁾, Yoshiki Murakami¹⁾, Kodai Kusakisako⁴⁾ Keiko Ikeda¹⁾, Mitsuhiro Asakawa¹⁾, Kazuyuki Suzuki¹⁾, Chiaki Ishihara¹⁾ and Hiromi Ikadai⁴⁾*

¹⁾ *School of Veterinary Medicine, Rakuno Gakuen University, Ebetsu, Japan*

²⁾ *Department of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria.*

³⁾ *College of Veterinary Medicine, Federal University of Agriculture, Abeokuta, Nigeria.*

⁴⁾ *School of Veterinary Medicine, Kitasato University, Aomori, Japan.*

*Corresponding author: Hiromi Ikadai

Laboratory of Veterinary Parasitology, School of Veterinary Medicine, Kitasato University
Higashi 23-35-1, Towada, Aomori 034-8628, Japan.

E-mail: ikadai@vmas.kitasato-u.ac.jp

Tel.: +81-176-23-4371

Fax: +81-176-25-0165

Running Head: Epidemiology of *Babesia canis* in Nigeria

ABSTRACT

The present study examined the presence of *Babesia* parasites in 104 domestic dogs in Nigeria. Sequentially, *Babesia* parasites infecting domestic dogs underwent genetic and phylogenetic analyses. The results of nested PCR based on the Piroplasmida 18S rRNA gene illustrated that 13.5% (14/104) of the samples were positive. The obtained positive samples determined the nucleotide sequences of the 18S rRNA genes. In the genetic and phylogenetic analyses, four of five nucleotide sequences were similar to *Babesia canis rossi*, and one sample exhibited a close similarity to a *Babesia* sp. isolated from a raccoon in Hokkaido, Japan. The present study revealed the widespread presence of *B. canis rossi* among domestic dogs in Nigeria.

Keywords: 18S rRNA, *Babesia canis*, domestic dog, Nigeria, phylogenetic analysis

Canine babesiosis is caused by the intra-erythrocytic protozoa of the genera *Babesia* belonging to the order Piroplasmida. *Babesia* classification in dogs is performed microscopically according to the parasite morphology in erythrocytes. Large *Babesia* parasites in canines are usually referred to as *Babesia canis*, whereas small *Babesia* parasites are classified as *B. gibsoni* [3, 8, 14, 25]. Recently, the detection and identification of Piroplasmida in infected domestic animals including dogs and wild animals was enabled by molecular techniques [1-3]. Moreover, the canine *Babesia* parasites include *B. annae*, which was initially classified in the genus *Theileria*, in domestic dogs in Spain [6], as well as *B. conradae* [13] and *Rangelia vitalii* [4, 5, 23]. *B. canis* has been classified into three subspecies by molecular methods: *B. canis canis*, *B. canis rossi*, and *B. canis vogeli* [3, 14]. *B. canis rossi* is the most pathogenic of the three subspecies, and it was first isolated from the side-striped jackal (*Canis adustus*) in Kenya [17]. This protozoan has been detected in sub-Saharan Africa, and it is known to be transmitted by ticks such as *Haemaphysalis leachi* [15-17, 21]. Conversely, *B. canis vogeli* has been detected in southern Europe, Asian countries including Japan, and South Africa [1, 16-18], and its virulence is weaker than that of the other two subspecies. *B. canis vogeli* is transmitted by *Rhipicephalus sanguineus* [15-17]. In Nigeria, the detected *Babesia* spp. include

B. canis rossi and *B. canis vogeli*, as well as other tick-borne pathogens such as *Hepatozoon canis*, *Ehrlichia canis*, and *Rickettsia* spp. [1, 11, 21]. In addition, *Babesia* parasites have been found in dogs in Nigeria [10, 22], Sudan [16], Egypt [20], and South Africa [17], but reports about canine babesiosis in the African continent are sparse. In this study, we collected blood samples from 104 domestic dogs in 2013 to assess the presence of *Babesia* parasites. Sequentially, genetic and phylogenetic analyses of the 18S rRNA gene were performed using *Babesia* parasites obtained from domestic dogs.

In 2013, ethylenediaminetetraacetic acid (EDTA)-anticoagulated blood samples were aseptically collected on Whatman FTA® classic cards (Whatman International Ltd, Maidstone, UK) from 104 domestic dogs that underwent rabies vaccination at the Veterinary Hospital of Federal University of Agriculture, Abeokuta (FUNAAB), in Nigeria. The dogs did not display any clinical signs of parasitemia. Samples were collected in accordance with the Guidelines for the Use of Experimental Animals. All procedures were reviewed and approved by the FUNAAB (approval number: FUNAAB/COLVET/CREC/202/1/09/02). FTA cards spotted with blood specimens from domestic dogs were used for DNA extraction using the phenol–chloroform–isoamyl alcohol (PCI) method [9]. Briefly, 100 µl of red blood cell (RBC) pellets were mixed

with a 10-fold volume of DNA extraction buffer (0.1 M Tris-HCl, pH 8.0; 0.1 M NaCl; 10 mM EDTA containing 1% Sodium dodecyl sulfate (SDS)). The contents of the tube were mixed thoroughly and incubated at 55°C for 2 hr with 50 µg of proteinase K (10 mg/ml), followed by the addition of an equal volume of PCI. Finally, genomic DNA (gDNA) from the aqueous phase was precipitated with ethanol, and the pellet was dried and dissolved in 25 µl of TE buffer (0.1 M Tris-HCl, pH 8.0, 10 mM EDTA). The dissolved gDNA samples were stored at –20°C until use in PCR.

The gDNA extracts from dog blood samples were first screened for the presence of *Babesia* and *Theileria* using previously reported primers [7], which target a 1,616-base pair (bp) portion of the 18S rRNA gene. The 18S rRNA gene was amplified using Piro0F (5'-GCCAGTAGTCATATGCTTGTGTTA-3') and Piro6R (5'-CTCCTTCCTYTAAGTGATAAGGTTTAC-3') for the first-round PCR and Piro1F (5'-CCATGCATGTCTWAGTAYAARCTTTTA-3') and Piro5.5R (5'-CCTYTAAGTGATAAGGTTTCAAACTT-3') for nested PCR as previously described [7, 12]. All PCR products were isolated by 1.0% (w/v) agarose gel electrophoresis in TAE buffer and purified using a GENECLAN kit (BIO 101, Vista, CA, USA). The 18S rRNA gene

88 was successfully amplified from the gDNA of spotted blood samples using nested PCR (data
89 not shown). The prevalence of Piroplasmida infection according to nested PCR using the
90 aforementioned primer set for the 18S rRNA gene was 13.46% (14/104). Genetic and
91 phylogenetic analyses were performed in all positive samples (n = 14) by nested PCR. The 18S
92 rRNA genes were directly determined using a CEQ8000 automated sequencer (Beckman
93 Coulter, Inc., Brea, CA, USA) with the DTCS DNA Sequence kit (Beckman Coulter). DNA
94 sequences were analyzed using Mac Vector software package, version 12.5.1, and their identity
95 was confirmed by comparison to homologous sequences available in the GenBank database
96 using the NCBI Basic Local Alignment Search Tool (BLAST,
97 <https://blast.ncbi.nlm.nih.gov/Blast.cgi>). In a BLAST search analysis, 10 positive were closely
98 related to the genus *Babesia*, and the remaining four were closely related to the genus
99 *Hepatozoon*. We determined that the complete nucleotide sequence of the 18S rRNA genes from
100 10 samples (1,560 bp) with high homology to the genus *Babesia*, followed by genetic and
101 phylogenetic analyses. Six of ten positive samples had identical nucleotide sequences, whereas
102 the other four samples had different nucleoside sequences. We obtained a total of five positive
103 samples. None of these sequences was identical to any sequence in public databases. In a

104 BLAST search of GenBank, four of five 18S rRNA genes were found to be closely related to
105 that of *B. canis rossi* (AB303075, identity: 99.7, 99.5, 99.7, and 99.5%, respectively) detected in
106 dogs in Nigeria, and one positive sample (AB935162) was revealed to be closely related to
107 *Babesia* sp. (AB251610, 99.5%), which was isolated from a wild raccoon in Hokkaido, Japan
108 [12]. All positive samples of 18S rRNA genes were related to the *B. gibsoni* 18S rRNA
109 sequence (AF271081, identity: 99.4%). We registered the nucleotide sequences of the 18S
110 rRNA gene as *Babesia canis rossi* strain N1 (AB935163), *Babesia canis rossi* strain N2
111 (AB935164), *Babesia canis rossi* strain N3 (AB935165), *Babesia canis rossi* strain N4
112 (AB935166), and *Babesia* sp. Nigeria2014 (AB935162) in GenBank. These sequences were
113 aligned using the CLUSTAL W Alignment program [24], and a phylogenetic tree was
114 constructed by the neighbor-joining (NJ) method from the aligned sequences with the
115 phylogenetic analysis in the Mac Vector software package, version 12.5.1. Support for tree
116 nodes was calculated with 1000 bootstrap replicates using the bootstrap tree algorithm [19]. The
117 NJ phylogenetic tree constructed using the 18S rRNA gene sequences separated the *B. canis*
118 *vogeli*, *B. canis canis*, *B. canis rossi*, *Babesia* sp. (raccoon), and *B. gibsoni* groups, respectively
119 (Figure 1). Four of the obtained 18S rRNA gene sequences (AB935163-AB935166) belonged to

the clade of *B. canis rossi* (AB303075), which has been reported in Nigeria. Meanwhile, the remaining sample belonged to the clade of *Babesia* sp. (AB251610) isolated from Hokkaido, Japan (bootstrap value: 99%, Figure 1). Two species of *Babesia* parasites, namely *B. canis rossi* and *B. canis vogeli*, have been detected in domestic dogs in Nigeria [1, 11, 21]. In a survey of canine babesiosis in Nigeria, the prevalence of *B. canis rossi* was 2.0% (8/400) in 2007 [21], 6.6% (12/181) in 2013 [11], and 38% (38/100) in 2014 [1]. Our study indicates that the prevalence of canine babesiosis has increased. We also detected new genotypes, which suggests that the parasitic lifecycle of domestic dogs, ticks, and *Babesia* parasites is widespread in Nigeria. However, few reports of canine babesiosis in Nigeria have been conducted using molecular techniques, and further investigation of the regional, seasonal, and epidemic variations of canine babesiosis in Nigeria and the African continent is needed. In addition, the vector ticks and the pathogen are spreading to new areas because of internationalization, increased trade, and climate change [10, 11]. The possibility of the spread of the disease to new areas in addition to the previously reported areas must be investigated.

CONFLICT OF INTEREST

136 The authors declare that they have no conflict of interest.

137

138 **ACKNOWLEDGMENTS**

139 We thank the late Professor Masayoshi Tsuji for his seminal contributions to this research.

140

141

142

143

144

145

146

147

148

149

150

151

REFERENCES

1. Adamu, M., Troskie, M., Oshadu, D. O., Malatji, D. P., Penzhorn, B. L. and Matjila, P. T. 2014. Occurrence of tick-transmitted pathogens in dogs in Jos, Plateau State, Nigeria. *Parasit Vectors* **7**: 119.
2. Cornillot, E., Dassouli, A., Garg, A., Pachikara, N., Randazzo, S., Depoix, D., Carcy, B., Delbecq, S., Frutos, R., Silva, J. C., Sutton, R., Krause, P. J. and Mamoun, C. B. 2013. Whole genome mapping and re-organization of the nuclear and mitochondrial genomes of *Babesia microti* isolates. *PLoS One* **8**: e72657.
3. Duarte, S. C., Parente, J. A., Pereira, M., Soares, C. M. and Linhares, G. F. 2011. Phylogenetic characterization of *Babesia canis vogeli* in dogs in the state of Goias, Brazil. *Rev. Bras. Parasitol. Vet.* **20**: 274-280.
4. Eiras, D. F., Craviotto, M. B., Baneth, G. and More, G. 2014. First report of *Rangelia vitalii* infection (canine rangelirosis) in Argentina. *Parasitol. Int.* **63**: 729-734.
5. Franca, R. T., Da Silva, A. S., Loretti, A. P., Mazzanti, C. M. and Lopes, S. T. 2014. Canine rangelirosis due to *Rangelia vitalii*: From first report in Brazil in 1910 to current day - A review. *Ticks Tick Borne Dis.* **5**: 466-474.

- 168 6. Garcia, A. T. 2006. Piroplasma infection in dogs in northern Spain. *Vet. Parasitol.* **138**:
169 97-102.
- 170 7. Hirata, H., Kawai, S., Maeda, M., Jinnai, M., Fujisawa, K., Katakai, Y., Hikosaka, K.,
171 Tanabe, K., Yasutomi, Y. and Ishihara, C. 2011. Identification and phylogenetic analysis of
172 Japanese Macaque *Babesia*-1 (JM-1) detected from a Japanese Macaque (*Macaca fuscata*
173 *fuscata*). *Am. J. Trop. Med. Hyg.* **85**: 635-638.
- 174 8. Hunfeld, K. P., Hildebrandt, A. and Gray, J. S. 2008. Babesiosis: recent insights into an
175 ancient disease. *Int. J. Parasitol.* **38**: 1219-1237.
- 176 9. Joseph Sambrook, D. W. R. 2001. Molecular Cloning: a Laboratory Manual, 3rd ed., Cold
177 Spring Harbour Laboratory Press, New York.
- 178 10. Kamani, J., Baneth, G. and Harrus, S. 2019. An annotated checklist of tick-borne
179 pathogens of dogs in Nigeria. *Vet Parasitol Reg Stud Reports* **15**: 100255.
- 180 11. Kamani, J., Baneth, G., Mumcuoglu, K. Y., Waziri, N. E., Eyal, O., Guthmann, Y. and
181 Harrus, S. 2013. Molecular detection and characterization of tick-borne pathogens in dogs
182 and ticks from Nigeria. *PLoS Negl. Trop. Dis.* **7**: e2108.
- 183 12. Kawabuchi, T., Tsuji, M., Sado, A., Matoba, Y., Asakawa, M. and Ishihara, C. 2005.

184 *Babesia microti*-like parasites detected in feral raccoons (*Procyon lotor*) captured in
185 Hokkaido, Japan. *J. Vet. Med. Sci.* **67**: 825-827.

186 13. Kjemtrup, A. M., Wainwright, K., Miller, M., Penzhorn, B. L. and Carreno, R. A. 2006.
187 *Babesia conradae*, sp. Nov., a small canine *Babesia* identified in California. *Vet. Parasitol.*
188 **138**: 103-111.

189 14. Lemos, T. D., Cerqueira Ade, M., Toma, H. K., Silva, A. V., Correa, R. G., Paludo, G. R.,
190 Massard, C. L. and Almosny, N. R. 2012. Detection and molecular characterization of
191 piroplasms species from naturally infected dogs in southeast Brazil. *Rev. Bras. Parasitol.*
192 *Vet.* **21**: 137-142.

193 15. Moraes, P. H., Rufino, C. P., Reis, T., Aguiar, D. C., Meneses, A. M. and Goncalves, E. C.
194 2014. Optimization of a molecular method for the diagnosis of canine babesiosis. *Rev. Bras.*
195 *Parasitol. Vet.* **23**: 105-108.

196 16. Oyamada, M., Davoust, B., Boni, M., Dereure, J., Bucheton, B., Hammad, A., Itamoto, K.,
197 Okuda, M. and Inokuma, H. 2005. Detection of *Babesia canis rossi*, *B. canis vogeli*, and
198 *Hepatozoon canis* in dogs in a village of eastern Sudan by using a screening PCR and
199 sequencing methodologies. *Clin. Diagn. Lab. Immunol.* **12**: 1343-1346.

- 200 17. Penzhorn, B. L. 2011. Why is Southern African canine babesiosis so virulent? An
201 evolutionary perspective. *Parasit Vectors* **4**: 51.
- 202 18. Rene-Martellet, M., Chene, J., Chabanne, L., Chalvet-Monfray, K. and Bourdoiseau, G.
203 2013. Clinical signs, seasonal occurrence and causative agents of canine babesiosis in
204 France: results of a multiregional study. *Vet. Parasitol.* **197**: 50-58.
- 205 19. Saitou, N. and Nei, M. 1987. The neighbor-joining method: a new method for
206 reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406-425.
- 207 20. Salem, N. Y. and Farag, H. S. 2014. Clinical, Hematologic, and Molecular Findings in
208 Naturally Occurring *Babesia canis vogeli* in Egyptian Dogs. *Vet. Med. Int.* **2014**: 270345.
- 209 21. Sasaki, M., Omobowale, O., Tozuka, M., Ohta, K., Matsuu, A., Nottidge, H. O., Hirata, H.,
210 Ikadai, H. and Oyamada, T. 2007. Molecular survey of *Babesia canis* in dogs in Nigeria. *J.*
211 *Vet. Med. Sci.* **69**: 1191-1193.
- 212 22. Sasaki, M., Omobowale, O., Ohta, K., Tozuka, M., Matsuu, A., Hirata, H., Nottidge, H. O.,
213 Ikadai, H. and Oyamada, T. 2008. A PCR-based epidemiological survey of *Hepatozoon*
214 *canis* in dogs in Nigeria. *J. Vet. Med. Sci.* **70**: 743-745.
- 215 23. Soares, J. F., Girotto, A., Brandao, P. E., Da Silva, A. S., Franca, R. T., Lopes, S. T. and

216 Labruna, M. B. 2011. Detection and molecular characterization of a canine piroplasm from
217 Brazil. *Vet. Parasitol.* **180**: 203-208.

218 24. Thompson, J. D., Higgins, D. G. and Gibson, T. J. 1994. CLUSTAL W: improving the
219 sensitivity of progressive multiple sequence alignment through sequence weighting,
220 position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:
221 4673-4680.

222 25. Uilenberg, G., Franssen, F. F., Perie, N. M. and Spanjer, A. A. 1989. Three groups of
223 *Babesia canis* distinguished and a proposal for nomenclature. *Vet. Q.* **11**: 33-40.

224

225

226

227

228

229

230

231

232 **FIGURE LEGEND**

233 Figure 1. Neighbor-joining phylogenetic tree presenting the relationships of 18S rRNA gene
234 sequences from *Babesia* isolates. GenBank accession numbers are presented in the tree. The
235 corresponding *Hepatozoon canis* (AF418558) sequence served as an outgroup. Numbers at the
236 nodes indicate bootstrap support from 1,000 iterations.

Figure 1