JOURNAL
OF THE
Japanese Society of Veterinary Science

Vol. XV.

DECEMBER, 1936

No. 4

**On the Epidemic Equine Encephalitis Which
Occurred in the Year 1935 in Japan.***

By

O. EMOTO, S. KONDO and M. WATANABE.

*(Veterinary laboratory, Ministry of Agriculture
and Forestry, Nishigahara, Tokyo.)*

(Received for publication September 1, 1936)

During the months of June, July and August 1935, an epidemic of equine encephalitis broke out in various parts of our country, amounting to about 900 cases. Clinically, most of them resembled very much the Borna disease reported in Germany or the equine encephalomyelitis found in the U. S. A. The mortality was recorded as about 33%. For the past several years, the Kyusyu island, a southern district of Japan, has been invaded periodically by an endemic malady of horses under the name of cerebrospinal meningitis. From the description of clinical symptoms and anatomical changes reported by many investigators,⁽¹⁾⁽²⁾⁽³⁾⁽⁴⁾⁽⁵⁾⁽⁶⁾ the malady seems to be the same as the encephalitis observed by us in 1935. We have had some opportunity to observe the disease in the field and to study it in the laboratory during 1935, but it is sure that the disease made its first appearance in the Kyusyu district and then spread widely in various parts of Japan. Our study is at present still under way, but we may report briefly the

* Japanese abstract, pp. 219—221

results so far obtained. Since the middle part of June, 1935, the livestock owners and the veterinarians in the Kyusyu island began to pay attention to a peculiar disease affecting the central nervous system of horses. Within two months the malady spread over the country, reached its highest severity from late in July to early in August and disappeared with the onset of cool nights. Few cases were reported in October. According to the official record, 832 horses excluding some military horses were affected and the mortality was, as already pointed out, about 33%. It has been observed that young horses (2~4 years old) were most susceptible, but aged horses could not escape from the attack of the disease. It will be of interest that stallions were highly susceptible and mares came next, geldings being less susceptible than mares.

Climatic conditions appeared to have an influence on the invasion of the disease, the outbreak of which being severer when the weather is abnormally warm and moist. It may be said that there is a certain relation between the climate and the outbreak of the disease as shown in the charts 1.2.3.

It is worthy of note that the first outbreak of the malady followed the onset of abnormally warm and moist weather. There was no observation as regards the natural infection of the disease from animal to animal. The affection due to cohabitation has never been observed, and in spite of transportation of affected horses, the disease had never been introduced into the parts of country in which the disease had never been found.

SYMPTOMS.

It is quite difficult to determine the period of incubation in the naturally infected animals. In a few cases, however, it was shown that the symptoms appeared after a lapse of from 1 to 7 days. It has been found that the incubation time in the small laboratory animals seemed to be 1~4 weeks. In some cases, a temperature rise up to 41°C. preceded the appearance of the symptoms. In the severe acute cases, the horses stagger, fall down,

Chart 1. Max. temperature, min. temperature and average moisture in May, June and July, in Fukuoka prefecture.

Chart 2. Max. temperature, min. temperature and average moisture in May, June and July, in Saga prefecture.

Chart 3. Max. temperature, min. temperature, and average moisture in May, June, and July, in Tochigi prefecture.

are unable to get up, and lie on their sides. Convulsive or swimming motions are commonly seen. In the advanced stage, the animals become rather quiet. In most of cases, the pulse and respiration are not accelerated. Difficulty in swallowing, fibrillar spasm of the muscles of the limbs, continuous or periodic grinding action of teeth, sightlessness, and paresis of tongue and lips are

characteristics shown in those cases.

Two clinical types were commonly observed in the milder cases; the sleepy or encephalic and the walking or spinal ataxic type. In the former, the horse leans with a drooping head against any supporting object in a drowsy somnolent state. A peculiar looseness of the lips with salivation can be observed.

The walking type is characterized by circling motion, paralysis or paresis of the posterior extremities and retention of urine. In such cases the animals walk closely to the fence or wall and turn round when they come in contact with an obstacle.

This walking type is more common than the sleepy type. In both types the appetite diminishes from the beginning of the disease, but it rarely happens that some horses do not lose appetite.

COURSE AND PROGNOSIS.

Most of the affected horses fall victims to the disease within 5 days after the onset of the symptoms. In some severe cases, the horses usually die within 24~48 hours. In milder cases, however, recovery can be expected.

ANATOMICAL CHANGES.

An autopsy of animals killed before complications showed no marked anatomical changes except a slight nephritis.

In a single case examined, the spinal fluid contained some lymphocytes and its specific gravity was 1.011 and pH 7.1.

Beside a slight hyperemia, no lesion was to be found macroscopically in the brain and the spinal cord.

HISTOLOGICAL FINDINGS.

The organs of seven horses were examined histologically, two of them having been killed by bleeding to death and five of them having died after a severe agonal struggle. The organs were fixed

within half an hour or twelve hours after death. The nervous symptoms in these horses continued from several hours to several days. Macroscopically, the nervous system was congested, while no marked lesion was found in other organs.

a) Materials examined

- 1) Brain of "Fukuoka strain" (killed by blood letting)
- 2) " of "Saga strain" (" ")
- 3) " of "Saga No. 1 strain" (died of disease)
- 4) " of "Saga No. 2 strain" (" ")
- 5) " of "Saga No. 3 strain" (" ")
- 6) " of "Tochigi A strain" (" ")
- 7) " of "Tochigi C strain" (" ")

Cultures showed that in the majority of cases the brains were sterile.

b) Fixation.

10% formalin, absolute alcohol, and Zenker's fluid were employed.

c) Staining.

Haematoxylin-eosin, Heidenhain's iron-haematoxylin, phloxin-methylenblue, and Unna-Pappenheim's methylgreenpyronin were used for the staining, and in some materials Giemsa, Lentz's and Mann's stain were used.

d) Histological findings.

Some characteristic findings were found only in the nervous system. We may classify them into two categories, (1) infiltration and (2) degeneration. In the "Tochigi A" strain a severe haemorrhage was found. Both of the histological changes were found in the brain, mesencephalon, cerebellum, spinal cord, and spinal ganglia.

(1) Brain.

The pia mater was infiltrated with mononuclear lymphocytes,

varying with individual cases a trace of the infiltration could be seen, and in other cases three or four layers of infiltrating cells were found. In general, lymphocytes and histiocytes were observed in the infiltrated lesions, but eosinophilic leucocytes and plasma cells were not found. The vessels of the pia mater were often surrounded by "cuffs" consisted of small mononuclears, but the change was very slight.

In the cortex, a localized or diffuse infiltration of lymphocytes accompanied by proliferation and mobilization of the neuroglial cells was seen. They were especially marked in the region of nucleus caudatus, nucleus lentiformis, and hippocampus. The infiltrative elements were lymphocytes rarely combining with large mononuclears and eosinophilic leucocytes, but plasma cells were not found up to this time. Haemorrhages surrounding the vessels were found at times. In the case of the "Tochigi A" strain a severe haemorrhage with a slight infiltration was found, (the infiltration was, however, quite severe in the pons, medulla oblongata and medulla spinalis.) This strain showed a very acute course, but no polymorphonuclear leucocytes were found in the lesion of perivascular infiltration. This strain resembled very much the western type of equine encephalomyelitis. No pronounced neurophagia could be found in the hippocampus, nucleus lentiformis, and lobus frontalis. In these regions the proliferation of neuroglial cells and the mobilization of glial cells were found, and at times so-called "Stäbchenzellen" of Dobberstein⁷⁾ were also found in those areas.

(2) Intranuclear inclusion.

The materials were fixed in Zenker's fluid, formalin, and absolute alcohol and the intranuclear inclusions which are usually associated with Borna disease were examined by means of Lentz's, Mann's-, phloxin-methlenblue-, Heidenhain's iron-haematoxylin-, Giemsa-, and haematoxylin eosin staining. The intranuclear corpuscles were found in the large ganglion cells of the hippocampus.

When sections were stained by means of Lentz's method, the intranuclear corpuscles appeared to be of pink or red colour,

while the nucleolus was shown to be more bluish. A single or a few homogenous corpuscles were found surrounded sometimes or not by characteristic halos, these findings being morphologically identical with those described as specific in Borna disease by Joest-Degen⁽⁸⁾(1909). Some satisfactory results were obtained when the tissues were fixed in Zenker's fluid, but not obtained when other staining methods with other fixatives were employed. These intranuclear corpuscles were found only in the "Tc-strain", but not in the other strains.

(3) Mid-brain and medulla oblongata.

In the mesencephalon and the medulla oblongata, similar infiltration and degeneration took place. Marked changes were also found in these regions and haemorrhages were found in medulla oblongata from some materials. In the "Ta-strain", the infiltration was very severe in the medulla oblongata.

(4) Cerebellum.

The cerebellum showed the infiltration and degeneration which were similar to those of the brain.

In the "Fukuoka-strain", both kind of these changes with haemorrhage were found to be marked. The protoplasm contained many vacuoles.

In the plexus chorioideus ventriculi quarti lateralis of the "Tochigi A-strain", many eosinophilic leucocytes appeared and also haemorrhage was seen.

(5) Spinal cord.

As in the brain material, the meninges of the spinal cord were not severely affected. Only in the "Ta-strain" a slight meningitis was found. On the contrary, some histological changes were always found in all materials from the spinal cords examined. Interstitial or perivascular infiltrations with mononuclear cells were observed, but they were found to be less intense.

Only in the "Ta-strain", a rather intense mononuclear infiltration and hyperemia or haemorrhage took place in the spinal cord

and they were found to be more intense in the posterior horn. Gliaproliferation and neuronophagia were also found in that region. The alterations were occasionally seen in the grey matter of the cord, but rarely in the white matter.

(6) Spinal ganglia.

In the lumbar ganglia of the cord from the "Ta-strain" we have found some perivascular infiltration and severe haemorrhage. So called granule cells were also found in that region. In the ganglia, a perivascular infiltration was also found and some mantle cells were observed to be increased. These histological changes in the peripheral nerves were recognized in the "Ta-strain".

SUMMARY.

The brain showed a slight meningitis with perivascular infiltration which formed nodules in the parenchyma.

There was found degeneration of nerve-cells, and at times haemorrhage especially in the "Ta-strain". Almost all the infiltrative elements were lymphocytes, rarely combining with eosinophilic leucocytes and histiocytes, but no neutrophilic leucocytes were to be seen. In the "Tc-strain", the intranuclear corpuscles described by Joest and Degen were found in some large ganglion cells of the cornu ammonis.

In the cerebellum, there was found a slight infiltration and degeneration. Haemorrhage occurred rather severely in the nucleus dentatus of the "F-strain" in which the degeneration of the nerve-cells was striking.

In the spinal cord of the "Ta-strain", a trace of meningitis and an intense perivascular infiltration were found in the grey matter or in the white matter of the cord, especially in the posterior horn. In this strain, the infiltration and degeneration were more intense than those which occurred in the brains, and haemorrhage was much advanced in the cord and brain, while the infiltration was not so pronounced. The spinal ganglia were intensely affected, showing both infiltration and haemorrhage.

It may be pointed out that the changes were always found in the central and also in the peripheral nervous system and, except the "Ta-strain", they were more intense in the brain than in the cord. The results obtained are shown in the following table.

Table I. Summary of histological findings.

		Trigonum olfactorium			L. pyriformis			Thalamus opticus			Corpora quadrigemina			Pedunculi cerebri			Pons		
		I	B	D	I	B	D	I	B	D	I	B	D	I	B	D	I	B	D
F	Strain													#	#	+	#	#	+
Ta	"	-	+	±	+	+	±										#	#	#
Tc	"																		
S	"	#	-		#	-	#	#	-	+	+	±	+	#	-				
S, No. 1	"	-	+	-	±	+	+	±	-	+	-	-	±	+	-	+	-	-	±
S, No. 2	"																		
S, No. 3	"										#	-	±	#		+	-	+	

		N. lentiformis			N. caudatus			Cornu ammonis			Lobus frontalis			Lobus occipitalis			Lobus temporalis		
		I	B	D	I	B	D	I	B	D	I	B	D	I	B	D	I	B	D
F	Strain	#	-	+				#	-	+	#	-	+						
Ta	"	#	#	#				+	±	+	#	±	#						
Tc	"				#			+	+	#									
S	"	#	-	±				#	-	+	#	±	+	+	+	+			
S, No. 1	"	-	-	-							±	±	±	±	±	±	#	#	+
S, No. 2	"							+	-		-	±	±	#	±	±	#	±	+
S, No. 3	"										#	-	±	+	±	±	-	-	-

		Lobus parietalis			N. dentatus			Medulla oblongata			Medulla spinalis			Nervus periphericus			Plexus chorioideus ventriculi quarti lateralis		
		I	B	D	I	B	D	I	B	D	I	B	D	I	B	D	I	B	D
F	Strain	+	-		+	+	+	+	-	+									
Ta	"				-	+	-	+	+		+	+	+	+	+	+			
Tc	"				+	+		+	+	+	+	-	+						
S	"	+			+	-	+				+	-	+				Eosin	+	+
S. No. 1	"	-	-	-	-	-	-				+	+							
S. No. 2	"	-	-	-							+								
S. No. 3	"	+	-	+	+	-	+	+	-	+	+	+	+						

I: Infiltration by mononuclear cells., B: Haemorrhage,

D: Degeneration of nerve cells.

Experimental transmission of the disease to horse and laboratory animals.

A) Materials

The inoculation materials which we have employed are shown in the following table.

Origin of strains	Tissue	Histological change
a) F-strain (Fukuoka)	Brain	Positive
b) S-strain (Saga)	"	"
c) Ta-strain (Tochigi A)	"	"
d) Tb-strain (Tochigi B)	"	not examined
e) Tc-strain (Tochigi C)	"	Positive
f) H-strain (Hanshin)	"	not examined

B) Experimental animals

The following animals were used.

- Horses (2~8 years old)
- Rabbits (600~2,500 g.)
- Guinea-pigs (250~300 g.)
- Mice (8~25 g.)

C) Method of inoculation.

A small piece of brain tissue was removed aseptically from the dead or killed horse and ground up in a sterile mortar and then emulsified with physiological saline solution or bouillon in a proportion of 1 to 5. This non-filtered emulsion was inoculated intracerebrally into animals.

The doses for rabbit were 0.1 to 0.5ccm., for guinea-pigs 0.5 to 0.1 ccm., for mice 0.025 to 0.05 ccm. and for horses 0.5 to 1.0 ccm.

D) Transmission tests.

- (1) Animals inoculated with the "F-strain".

This strain has been obtained from a horse which was killed in the field 5 days after the onset of the disease and at that time the horse seemed to be recovering. A piece of brain (lobus frontalis) was removed aseptically when the horse was dissected, and an emulsion of the tissue with a diluent about five times its weight was prepared and then was injected intracerebrally into rab-

Table 2. Inoculation of F-Strain to rabbits.

bits. The results obtained are recorded in the following table.

As shown in the table, we were not able to transmit the disease to rabbits with this horse strain.

(2) Animals inoculated with the "S-strain"

The strain was obtained from a horse which had been killed in the field 7 days after the onset of the disease of a very acute course. A piece of brain (lobus frontalis) was removed aseptically when the horse was dissected and an emulsion of the tissue with a diluent about five times its weight was prepared and then was injected intracerebrally into rabbits, but no positive results were obtained.

(3) Animals inoculated with the "Ta-strain".

The horse died in the night 3 days after the onset of the disease with a very acute course. A few pieces of brain (trigunum olfactorium, cornu ammonis and cerebellum) were removed aseptically when the horse was dissected in the following morning.

a) Animals inoculated with the material from trigunum olfactorium.

An emulsion of the tissue with a diluent about five times its weight was prepared and, at once, it was injected intracerebrally into a horse, two mice, two guinea-pigs and two rabbits.

Table 3. Inoculation of Ta-strain (Trigunum olfactorium)

As shown in the table we are inclined to believe that we have succeeded in infecting horse and rabbits with an emulsion of the brain of the affected horse.

The incubation period of the disease in the horse infected by the intracerebral route was 16 days. On the 18th day after the inoculation, paralysis occurred in the hind extremities following a decrease of appetite for two days, consequently the horse lay down on his side and was unable to rise up. Paresis appeared in the lips and in the tongue. The animal was unable to swallow the food taken in the mouth, and salivation was observed. When the stage was advanced a complete loss of sensibility of the skin was observed and the horse had a good sleep. After a week, the horse has gradually recovered, so that we killed the horse by blood letting 52 days after the inoculation. An autopsy of the horse showed no significant changes. Beside a slight change in the liver (nutmeg liver), no marked lesions were macroscopically found in other organs.

On culturing the brain and organs were found to be quite sterile.

Histological changes of the region of the left hemisphere were examined, because the emulsion was inoculated in the right one.

There has been found a localized or diffuse infiltration of lymphocytes accompanied by haemorrhage, proliferation and mobilization of neuroglial cells. They were very marked in the region of nucleus caudatus, trigonum olfactorium and lobus temporalis. Neuronophagia was noted to be very intense in the lobus frontalis, and lobus temporalis. Glial nodules were also found in the trigonum olfactorium. Intense haemorrhage was seen in the cortex and in the cornu ammonis. No alteration in the spinal cord was to be found, and no intranuclear corpuscles were demonstrated in the ganglion cells. But from these points, it can be concluded that this horse was infected by intracerebral inoculation.

Some of the rabbits which were inoculated showed typical symptoms and in their brains similar histological changes to those of affected horses were found. The changes of brain of the rab-

bits resembled very much that of spontaneous encephalitis of rabbits. As regard these points, the following discussion shall be referred to.

b) Animals inoculated with the material from the cornu ammonis of the "Ta-strain".

An emulsion of the tissue with a diluent about five times its weight was prepared immediately after the tissue was removed, it was then injected intracerebrally into a horse, three mice, two guinea-pigs and two rabbits. In these experiments no satisfactory results were obtained.

c) Animals inoculated with the material from cerebellum, pons, pedunculi cerebri, pyriformis and medulla oblongata of the "Ta-strain".

An emulsion was prepared as described in (b) and injected into a horse, four mice, two guinea-pigs, and two rabbits. Some of the typical symptoms appeared in the rabbits, but we were unable to obtain any positive results so far the histological changes of their brain are concerned.

(4) Animals inoculated with the "Tb-strain".

The symptoms observed in the horse (Tb-strain) were quite similar to those of the "Ta-strain". The horse died 3 days after the onset of the disease. The materials were already decomposed, so the emulsion was filtered through Chamberland L1 and then injected into a rabbit. After a few days, convulsion was observed in the neck region, but no histological changes were observed in the nervous system. By using this strain no satisfactory results were obtained.

(5) Animals inoculated with the "Tc-strain".

The material was given to our laboratory from the military veterinary school. The intranuclear corpuscles were found in the ganglion cells of the cornu ammonis.

a) Animals inoculated with the material from the cornu ammonis.

An emulsion was inoculated into a horse, three mice, two

guinea-pigs and two rabbits, but these experiments must be omitted, because the material was already contaminated with pathogenic streptococci.

b) Animals inoculated with the material from the trigonum olfactorium.

Small pieces of trigonum olfactorium were soaked in a 50% solution of glycerine and placed in the ice-box over night. The next morning, an emulsion from the tissue with a diluent five times its weight was prepared and it was immediately injected intracerebrally into a horse, two guinea-pigs, and a rabbit. See the following table.

Table 4. Inoculation of Te-strain (Trigonum olfactorium)

As shown in the table, we succeeded in infecting rabbits with an emulsion from a horse dead of epidemic equine encephalitis. The incubation time of the disease in these rabbits was about 1~4 weeks, however that of the forth generation which showed typical symptoms and changes was about 10 months. Intranuclear corpuscles were found in the large ganglion cells of cornu ammonis.

(6) Animals inoculated with the material from the "H-strain"

By employing this strain, we could not obtain any satisfactory results in rabbits.

SUMMARY.

We may believe that we succeeded in infecting a horse and some rabbits with the emulsion of brain from two cases of horses infected with the epidemic equine encephalitis which was confirmed histologically.

DISCUSSION.

Basing our conclusion upon the epidemiology, symptomatology, pathology and histology, we maintain that the disease which occurred during the course of the year 1935 in this country, should be regarded as an epidemic equine encephalitis.

In spite of having devoted ourselves to the performance of experiments, we could not obtain any satisfactory results regarding inoculation tests. As far as the isolation of the contagious agent and the performance of the cross immunization tests go nothing can be communicated. For this reason, we are quite unable to draw any reasonable conclusion, whether this disease can be considered as a new one or some other which may be compatible with the Borna disease in Germany or the equine encephalomyelitis in the U. S. A. Nevertheless we may discuss the results obtained from our experiments.

From the point of view of epidemiology, symptomatology and pathological-anatomy, the disease has a great resemblance to the Borna disease, though the affected horses died after rather an acute course, viz., within 5~7 days after the appearance of the symptoms, there being of course some exceptional cases.

It has been found by histological examination of all the strains, excluding the "Ta-strain", that the changes were almost similar to those of the Borna disease described by Joest-Degen⁽⁹⁾, Zwick and his collaborators⁽⁹⁾, and by Nicollau and Galloway⁽¹⁰⁾. Among these changes the perivascular infiltration with mononuclears, the degeneration of nerve cells was marked, but the haemorrhage of the small vessels was very slight, and all these changes were recognized to be more intense in the brain than in

the cord.

The intranuclear corpuscles described by Joest-Degen (1909) were found in the large ganglion cells of cornu ammonis only when the tissue was fixed in Zenker's fluid and stained by means of Lentz's method, the inclusion bodies not being found when the tissue was fixed in formalin.

In the brain of the "Ta-strain", however, the haemorrhage occurred rather severely and then perivascular infiltration was found to be slight, while the perivascular infiltration was found rather severe in the spinal cord and spinal ganglia. The intranuclear corpuscles were not still found. Judging from these facts, it may be said that the disease of the "Ta-strain" resembled very much the western type of the equine encephalomyelitis in the U. S. A. described by Meyer, Haring and Howitt⁽¹¹⁾; B. Howitt⁽¹²⁾; K. F. Meyer⁽¹³⁾; O. Larsell, M. Haring and K. F. Meyer⁽¹⁴⁾; and Rosenbdsch⁽¹⁵⁾, although no neutrophilic leucocytes were found in the lesions of perivascular infiltration. In the horse infected with the brain-emulsion of this strain, the inclusion bodies were not found, but severe hyperemia and haemorrhage were found in the brain. It may be considered that, as this strain took an acute course, the haemorrhage was more intense than the perivascular infiltration.

By carrying out our inoculation test, we may believe that we have succeeded in infecting one horse by injecting an emulsion from the horse dead of the disease in the field. We may say that there is no doubt about this infection, because typical symptoms and typical histological changes were found after the inoculation.

As controls, several horses were examined, but such specific changes were not found in the central nervous system. The horses employed as cotrols were healthy, but some of them were sick with some bacterial infection or infectious anaemia, and died of the infection, manifesting some resembling symptoms.

In rabbits, the experiments might prove to be difficult owing to the occurrence of spontaneous encephalitis as described by M. Nishibe⁽¹⁶⁾ and O. Seifried⁽¹⁷⁾, but in our experiments, the disease was very rare viz., only two among about 300 cases. From this

fact, the spontaneous encephalitis has not been taken in consideration. In our experiments, the infection test was regarded as positive when the typical symptoms and the typical histological changes were found, and as doubtful or negative when only one of those symptoms and histological changes were observed. As already described, the granulome was found in the spontaneous encephalitis while not in the animals infected with the emulsion. It must be pointed out that, in only one case, the intranuclear corpuscles were found in the ganglion cells of cornu ammonis taken from the infected rabbit, but the inclusion bodies were found to be absent in the case of spontaneous encephalitis.

In guinea-pigs and rats, we were unable to obtain any satisfactory results regarding the transmission of the contagious agent, and we could not find any reason why they were non-susceptible to the disease.

In mice, a typical case was obtained from a histological point of view, but no interesting symptoms were clinically observed.

In 1934, Max Theiler⁽¹⁸⁾ communicated that there is a spontaneous encephalitis among mice and the histological examination of the brain from the mice affected with the disease showed an interesting picture which resembled that occurring in the brain of the horse affected with the Borna disease or the equine encephalomyelitis of the American type. For this reason, we hesitate to draw a conclusion that the mouse so treated was a victim of an artificial infection.

It is interesting that the incubation period of the affected rabbits was not constant. It is highly probable that the virulence of the contagious agent was very weak or its development in the brain might have been checked by something like an autosterilization, so that the experiment as regards the transmission of the contagious agent from animal to animal was not successful. We were of the opinion that the contagious agent can be regarded as a virus which resembles very much the Borna virus and the equine encephalomyelitic virus, because the epidemiology of the disease and anatomical and histological findings resembled those of the above mentioned diseases and, moreover, the animals are

susceptible to the filtrate from the sterile brain-emulsion made from the affected horse.

In case of an intracerebral inoculation of bacteria, the histological changes were found to be very striking, but differed very much from those found in the case of equine encephalitis. The course of bacterial encephalitis was generally acute and a large number of the neutrophilic leucocytes appeared in the lesions of infiltration. We have also observed neutrophilic leucocytic infiltration in the nervous system of guinea-pigs inoculated with an equine encephalomyelitic virus (given by Miss. Howitt), but it was very slight.

From these facts, it may be estimated that the contagious agent of the epidemic encephalitis is a kind of virus.

CONCLUSION.

(1) We have found two types in the epidemic equine encephalitis which occurred epidemically in this country last summer. The one resembled very much the Borna disease of Germany and the other resembled the equine encephalomyelitis (western type) in U. S. A. But we hesitate to determine whether the epidemic has the nature of the former or that of the latter, for we could not as yet succeed in performing the cross immunization tests.

(2) The virulence of the contagious agent was very weak, but we could not deny that we have succeeded in infecting one horse and a few rabbits with the emulsions prepared from the brains of affected horses.

(3) It may be recognized that a virus is the causative agent of the epidemic equine encephalitis which occurred in Japan in the year 1935 and that the brain of the affected horses is the principal site of the virus.

LITERATURE

- (1) S. Katsushima: *Chōō-Jui-Kai Zasshi* (Japanese). 1898, **5**, 54.
- (2) H. Tokishige: " " " " " **7**, 71.
- (3) " " " " " **8**, 1.
- (4) S. Katsushima: " " " " " 1899, **9**, 1.
- (5) H. Oguni: *Jueki-Chosajo Kōyū-Kai-Ho*, 1926, **5**, 27.
- (6) M. Watanabe and O. Emoto: *Kachiku-Eisei-Kyokai-Ho*, (Japanese), 1935, **3**, 3.
- (7) J. Dobberstein: *Z. f. Inf. d. Haustiere*, 1928, **33**, 290.
- (8) E. Joest and K. Degen: *Z. f. Inf. d. Haustiere*, 1909, **6**, 348.
- (9) W. Zwick, O. Seifried and J. Witte: *Z. f. Inf. d. Haustiere*, 1926, **30**, 42.
- (10) S. Nicolau and A. Galloway: *Medical Research Council, Special Report Series*, No. **121**, 1928.
- (11) K. F. Meyer, M. Haring and B. Howitt: *Science*, 1931, **74**, 227; *J. Amer. Vet. Med. Ass.* 1931, **79**, 376.
- (12) B. Howitt: *J. Inf. Dis.*, 1932, **51**, 493; *J. Immun.*, 1935, **11**, 319.
- (13) K. F. Meyer: *Z. f. Inf. d. Haustiere*, 1934, **35**, 81.
- (14) O. Larsell, M. Haring and K. F. Meyer: *Amer. J. Path.*, 1934, **35**, 81.
- (15) Rosenbusch: *Z. f. Inf. d. Haustiere*, 1934, **47**, 48.
- (16) M. Nishibe: *J. Path. Jap. Soc.*, 1926, **15**, 120.
- (17) O. Seifried: *Z. f. Inf. d. Haustiere*, 1929, **38**, 18.
- (18) Max Theiler: *Science*, 1934, **77**, Aug. 5.

-
- Fig. 1 Microphotograph, Leitz, oc. 5×, ob. 16 mm. Horse of the Tc-strain. Perivascular infiltration with lymphocytes in the nucleus caudatus. Haematoxylin eosin staining.
 - Fig. 2 Microphotograph, Leitz, oc. 5×, ob. 2 mm, apochromat (2× enlarged) Horse of the Tc-strain. Intranuclear corpuscle in the ganglion cell of the cornu ammonis. Lentz's Staining.
 - Fig. 3 Microphotograph, Leitz, oc. 5×, ob. 2 mm, apochromat. Horse of the Tc-strain. Nerve cells of nucleus caudatus showing neuronophagia H.-E.-staining.
 - Fig. 4 Photograph of the rabbit Tc-Bul III. 30 days after intracerebral infection with the brain-emulsion of the rabbit Tc-Bul II of Tc-strain. Paralysis of hind quarters and slight paresis of fore-legs. At this stage, the rabbit laid on the flank and struggled in vain with the fore-legs to recover its normal position.
 - Fig. 5 Microphotograph, Leitz, oc. 5×, ob. 2 mm, apochromat. Rabbit

Fig. 1. Horse Tc Nucleus caudatus. Leitz, ok. 5×, ob. 16 mm.

Fig. 2. Horse Tc Cornu ammonis. Leitz, ok. 5×, ob. 2 mm.

Fig. 3. Horse Tc Nucleus caudatus. Leitz, ok. 5 $\times$, ob. 2 mm.

Fig. 4. Rabbit Tc Bul. III, showing paralysis.

Fig. 5. Rabbit Tc Bul. III, Leitz, ok. 5 $\times$, ob. 2 mm.

Fig. 6. Inoculated Horse No. 126 showing paralysis.

Fig. 7. Inoculated Horse No. 126 Leitz ok. 5 $\times$, ob. 4 mm.

Fig. 8

Tc-Bul III. Vacuole in the cytoplasm of a degenerated neuron in the spinal cord, H.-E.-staining.

Fig. 6 Photograph of Horse pf. 126. 10 days after intracerebral infection with the brain-emulsion from the horse of Ta-strain. Paralysis of the hind quarters. At this stage, the horse was unable to rise up and paresis occurred in the lips and the tongue. He was unable to swallow the food taken in the mouth and salivation was observed. At the advanced stage, the sensibility of the skin was completely lost and the horse had a good sleep.

Fig. 7 Microphotograph. Leitz, oc 5× ob 4 mm, apochromat. Horse pf. 126. perivascular mononuclear infiltration in the trigonum olfactorium. H.-E.-staining.

Fig. 8 Microphotograph of perivascular lymphoid cell infiltration in the brain of a rabbit, which was formerly injected with the filtered material of the brain emulsion of an affected rabbit (Tc. strain of equine epidemic encephalitis, Leitz, oc 5× ob 4 mm. Apochr).

1935 年日本ニ流行セル馬流行性腦炎ニ就テ*

江 本 修， 近 藤 正 一， 渡 邊 守 松

(農林省獸疫調査所)

(昭和 11 年 9 月 1 日 受 附)

1935 年夏期ニ九州ニ初發シ其ノ後漸次本州ニ沿ヒ東ヘ東ヘト相繼イデ多
發セル馬ノ一種ノ神經病ガ流行シテ全國ニ於テ約 900 頭罹患シ斃死率ハ約
30~50%デアツタ然シ此猛烈ナ病氣モ秋ノ終リ頃ニ全ク終熄シテシマツタ

著者等ハ本病ニ就テ種々調査研究シタカラ以下其ノ概略ヲ記述スル事ト
スル

2~4 歳馬ガ最も多ク罹患シ約 5~7 日以内ニ斃死スベキモノハ斃死シタ
而シテ牡ガ最も多ク罹患シタ之等ノ馬ハ他ノ土地カラ該土地ニ移入セラレ
テ而モ始メテノ夏ヲ迎ヘタモノバカリデ氣候の影響ガ本病發生ニ相當重大
ナル要素デアルコトガ想像セシメラレル試ミニ最近 10 年間ノ氣候狀況ヲ
調査シテモ 1935 年ハ非常ニ氣候不順デ 5, 6, 7 及 8 月ノ 4 ケ月ニ就テ
調査シテ見ルニ 1 日ノ最高最低溫度ノ差ガ可ナリ著シク又 1 ケ月ノ中デモ
溫度及濕度ノ高低ハ相當ニ顯著デアツテ之レ等ノ高低ノ非常ニ著シカツタ
後ニ初發ヲ見テキル所カラ察スルニ氣候の影響ハ少クトモ一ツノ要素ト考
ヘ得ラレル

傳染經路ハ全ク不明デアツテ多クノモノハ突然爆發の流行ヲ來シ病例相
互ノ間ニ何等ノ關聯モ認メラレズ又同居感染ノ例モ殆メ認メラレナイ

臨床的ニハ神經症狀ヲ主トシ沈鬱或ハ興奮シ且廻轉運動，步行困難，視
力障礙，口唇四肢ノ麻痺，四肢頸側等ノ強直及振顫，食欲減退等ノ症狀ヲ
示シ急性ノモノハ 24 時間内ニ斃死スルモノ多クハ 1 週間以内ニ斃死シ其レ
以上長引クモノハ治癒スルモノガ多クナツテ居ル

剖檢上肉眼的ニハ殆ド特異性變化ナク只腦膜充血，腎臟及ビ膀胱内ノ含

* 原著歐文 41 頁參照

尿ノ輕度ノ濁濁等ヲ通常認メタノミデアル唯一例福岡馬ニ於テ小腦齒狀核ニ新發ノ點狀出血散發スルヲ見タ

組織學的ニハ必ズ中樞神經系統ニ著明ナル特異性變狀ヲ認メル其ノ病的變化ノ主ナルモノヲ記載スレバ（１）小血管及毛細血管周回ノ淋巴腔ノ擴大竝ニ腔内ニ著明ナル淋巴細胞浸潤ヲ認メル中性多核白血球ハ認メラレナイ（２）神經細胞ノ種々ナル程度ノ退行變性竝ニ神經喰現象乃至膠質細胞ノ結節狀ニ集合セル像ヲ見ル（３）毛細管出血モ相當ニアリ必發トハ限ラナイガ相當注目ニ價スル（４）全例デハナイガ Joest-Degen 小體ヲアンモン角神經細胞核内ニ證明スル事ガ出來タ

次ニ無菌的ニ採取セル腦ノ１片ヲ約５倍ノ食鹽水乳劑トナシ其ノ無菌ナル事ヲ証明シテ後馬、兎、海狸、マウス及ラツテノ腦内ニ接種シテ馬及兎ニ感染サセル事ニ成功シタ

兎ハ 1935 年 4 月日本獸醫學會ニ於テハ 3 代目迄動物通過ニ成功セル事ヲ報告シタガ其ノ後之ヲ Chamberland L. I. デ濾過シテ兎ニ接種シ完全ニ之レニ感染セシメタ

以上ノ研究ノ結果ヲ綜合シテ考察スルニ 1935 年夏季本邦ニ流行狀ニ多發セル馬ノ一新奇病ハ確ニ神經症狀ヲ主徵トスル特殊ノ疾病デアツテ剖檢上肉眼的ノ變化ハ重要デナイガ組織學的ニハ腦實質ニ充血（稀ニ出血ヲ呈ス）、小血管周圍淋巴腔ノ擴大竝ニ其擴大淋巴腔内ニ著明ナル淋巴細胞ノ浸潤、神經細胞ノ種々ナル程度ノ退行變性等ヲ認メ且全例デハナイガアンモン角ノ神經細胞ノ核内ニ Borna 病ノ特徴トスル Joest-Degen 小體ト一致スル小體ヲモ認メタ次ニ試驗動物ニ於ケル接種試驗ノ成績ニ就テハ病獸ノ腦乳劑ヲ腦内ニ接種シテ馬及ビ兎ヲ發症セシムルコトニ成功シタ又兎ニ於テハ其腦乳劑ヲ濾過液ヲ以テシテ尙且兎ヲ感染セシムルコトガ出來タ又 Borna 病小體ト殆一致スル小體ノ見ユル腦ノ乳劑ヲ以テセバ夫レニ感染セル兎ノ腦ニモ矢張同一ノ小體ヲ認メ得ラレルノデテル

又濾液ヲ以テシテ試驗動物ヲ發症セシムルコトニ成功シタ結果ヨリ考ヘテ本病ノ病原體ハ濾過性病毒（Virus）デアルト想像セラレル尙以上ノ試験

ノ結果ヨリ本病ハ濾過性病毒ニ基因スル馬ノ流行性腦炎ト診斷スルコトガ
出來ルト思ハレル

之レヲ獨逸ノ Borna 病、米國ノ傳染性腦脊髓炎等ト比較スルニ本邦ノ
腦炎ハ比較的 Borna 病ニ近イモノノヤウニ思ハレルケレドモマダ今迄ノ
試験丈デハ兩者全然同一ノモノトモ又之レヲ本邦獨自ノモノトモ又本邦ニ
二型アルモノトモ斷定シ難イ

著者等ハ尙 1936 年ニ流行セル同様ノ疾患ニ就テ調査ヲ續行中デアルソ
シテ上記結論ニ益確信ヲ有スルヤウニナツタノデ之レハ次回ニ報告致シタ
ヒト思フテ居ル次第デアル