滿洲農村に於ける鼻疽馬の實態的調査研究

III. 開放性鼻疽馬の運命と新感染馬並に新生仔馬の反應態度について

村瀬信雄, 壽樂明雄, 田韞珠, 清水文康, 韓有庫

(元滿洲國大陸科學院馬疫研究處細菌第一研究室: 處長 小野紀道)

(昭和 27 年 1 月 31 日受付)

I. 緒 言

私共は前 2 報告に於て臨牀検査並に各種免疫反應を以つて同一對象に就いて繰返し實施した調査研究の結果について疫學的の考察を行い、最後にその一部の 32 頭の病理解剖學的検査を行つてマレイン反應及び補體結合反應陽性のものは呼吸器以外の器官にも比較的新鮮な病巢を有する現症の鼻疽ともいふべきものでありマレイン反應のみ陽性のものは比較的病變が呼吸器系統に局限し比較的陳久病巢を有する傾向のあるものであることを指摘し且つこれら免疫反應による顯われと各種環境要約の關係を精査した結果から常在地に於ける本病の經過は極めて慢性で飼料の質量と勞役量又牝馬にあつては妊娠などにより影響せられつゝあるもので、内部病變としては前記兩反應陽性の眞性鼻疽に於ても軽度でしかも早急に重症型に向う性質の窺われないことについて言及した

一方私共は本調査研究の間に於て傳染源として重視すべき開放性鼻疽馬を多數目撃したのであるがこれら開放性鼻疽馬が如何なる運命を辿るものかを知することは本病の本性を知る上にも又實際防疫の立場からも重要と考えて絶えず注目を怠らなかつた

又健康馬が新たに感染する様相即ち常在地に於ける自然傳播速度並に反應陽性馬から生産される幼駒の反應上の態度或は汚染の如何等も如上の觀點から知つて置かなければならないことであるがこれらについての記載は諸外國にも極めて少く唯 GLUCHOW が鼻疽馬から健康幼駒が産せられることについて述べているのみである 又開放性のものがどの位生存するものかについての記載は市川⁽¹⁷⁾によると急性症にあつては數週間、慢性のものでは數ヶ月又は年餘と述べられて居るが、根據ある正確な記載は見當らない

本報告に於てはこれらの諸項目について得られた結果をまとめた

II. 成 績

1. 開放性鼻疽馬に就いて

1943 年 7 月 第 1 回調査検査馬總數 929 頭中 29 頭 (3.1%) の開放性鼻疽馬を發見したが、第 1 報所報の如く本調査研究期間中は何等行政措置を講ずることなく成り行きの儘に任せてその運命を追求することにした その詳細は Table I の通りで、これをまとめると Table II となる

即ち性別に見ると騾馬が $\frac{22}{29}$ で 75.8% を占め、年齡的には 11-15 歳迄のものが

Table I. List of all the horses infected with open-glanders found in the investigations.

Horse No.	Sex	Age	Change in nasal cavity								Mallein eye-test								Compl. fixation test								Agglutination test							
			1st	2nd	3rd	4th	5th	6th	7th	8th	1st	2nd	3rd	4th	5th	6th	7th	8th	1st	2nd	3rd	4th	5th	6th	7th	8th	1st	2nd	3rd	4th	5th	6th	7th	8th
1	♂	12	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
2	"	8	+	+	+	d				+	+	+						+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
3	"	13	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
4	"	9	+	+	S					+	+							+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
5	♀	7	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
6	♀	8	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
7	♂	13	+	+	d					+	+							+	+															
8	"	11	+	S						+	+							+																
9	"	11	+	S						+								+																
10	♀	9	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
11	♂	12	+	+	+	+	+	+	S	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
12	♀	11	+	S						+								+																
13	♂	11	+	+	+	+	d			+	+	+	+					+	+	+	+													
14	♀	16	+	+	-	-	-	-	-	+	+	+	-	-	-	-	-	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	
15	♂	9	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
16	"	17	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
17	"	15	+	+	S					+	+							+	+															
18	"	11	+	d						+								+																
19	"	16	+	d						+								+																
20	"	8	+	+	+	+	+	+	S	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
21	♀	8	+	+	+	+	+	+	m	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
22	"	15	+	+	+	+	+	+	S	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
23	♂	14	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
24	"	17	+	+	S					+	+							+	+															
25	"	11	+	+	+	+	+	d		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
26	"	7	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
27	♀	14	+	d						+								+																
28	♂	16	+	S						+								+																
29	♀	12	+	+	d					+	+							+	+															
30	♂	8	-	+	+	+	+	+	+	+	+	+	⊕	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
31	♀	15	-	-	+	+	d			+	+	+	+					+	+	+	+													
32	"	11	-	-	-	+	+	d		+	+	+	+	+				+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
33	♂	14				+	+	+	S				+	+	+						+	+	+											
34	"	5				+	S						+								+													
35	"	9	-	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
36	"	13				-	-	+	+	+			+	+	+	⊕	+			+	+	+	+	+	+	+	+	+	+	+	+	+	+	
37	"	8	-	-	-	-	+	S		+	+	-	+	+				+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
38	"	9				+	+	+					+	+	+					+	+	+	+	+	+	+	+	+	+	+	+	+	+	
39	♂	7	-	-	-	-	+	S		+	+	-	+	+				+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
40	♀	8	-	-	-	-	+	+	+	+	+	-	-	-	⊕	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
41	"	12				+	+	+					+	+	+							+	+	+										
42	♂	7	-	-	-	-	+	+	+	+	+	-	-	-	+	+	+	-	-	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

Note: d...died s...sold m...missing

50% 餘で最も多いことが分る 鼻汁培養の結果 82.7% ($\frac{24}{29}$) に於て鼻疽菌を検出した 又此の 29 頭中 28 頭が私共の病性区分上 MS (マレイン反應並に補體結合反應陽性) に屬するもので常在地に於ける自然感染開放性鼻疽馬の反應態度が窺い知られる

次にこれら 29 頭の開放性鼻疽馬の運命を追求すると共に、新に出現して来る開放性

鼻疽馬を調査した結果得られた成績が Table III である

即ち第1回調査時の29頭の開放性のものゝ中、16ヶ月後の第7回調査時迄に10頭が賣却されて8頭斃死、非開放性に轉じたもの1例、不明1例となつて9例が最後まで開放性として残り此の間に9頭が新に開放性となり4頭が他の地域から購入され合せて13頭が新しく開放性鼻疽馬として入つて來たのであるが、そのうち斃死2頭賣却4

Table II. Description of 29 horses found with open-glanders in the 1st investigation.

Total	Sex			Age				Culture with nasal discharge		Classification			
	♂	♀	♂	0~5	6~10	11~15	16~20	Positive	Negative	MS	M	S	—
29	22	7	0	0	9	15	5	24	5	28	0	1	0

Table III. Details of the horses infected with open-glanders.

29 Cases in 1st investigation	Details	1st-2nd	2nd-3rd	3rd-4th	4th-5th	5th-6th	6th-7th	Total
	Sold	4	3				3	10
	Died	3	2	1	1	1		8
	Missing						1	1
	Turned to non-open type		1					1
	Remained open	22	16	15	14	13	9	
	Bought			2	2			4
	From non-open type	1	1	2	5			9
	Sold				1	2	1	4
	Died				1	1		2
	Remained open	1	2	6	11	8	7	
	Total	23	18	21	25	21	16	

Note: 1st-2nd denotes transitions of open-glanders during 1st and 2nd investigations.

頭で結局7頭が残り、第1回から残つた9頭と合せて第7回調査時には16頭の開放性鼻疽馬が見られた(997頭に對し1.6%)

Table IIIで特に注目すべきことは3月から5月迄に開放性の出現が最も多いことゝ、約1年6ヶ月で開放性のものゝ中斃死したもの8頭、賣却された10頭の中半數が斃死せるものと見て5頭計13頭、即ち45%弱が斃死の運命を辿るものであることが

考えられ、常在地に於ける開放性鼻疽馬の運命は平均2-3年であるとみて差支えない様である。しかもこれら開放性のものゝ大部分が此の間特に加療などのことなく他の健康馬と同様に勞役に供されて居つたことは注目されてよい。

次に凝集反應について見るならば29例中6例(20.7%)が1:800以下14例(48.2%)が1:1600僅かに8例(27.5%)が1:3200以上であつて次に報告する豫定の第IV報に示す如く健康馬に於ても時期によつては35.6%が1:1600を示すことから本反應の陽性限界については現在のところ第II報で一部論じたように1:3200以上が本病との結びつきを確實に考慮せられるべきだとすると、病性と本反應の關係の一部が解明せられて本反應の應用價值がどの程度のものであるかと窺われる。

2. 新感染馬について

Table IVに見られるように本調査期間即ち1943年7月以降1944年11月迄の16ヶ月間に合計23頭の明かに新感染馬と判定し得られるものが見られたが此の表で特に注目される點は第5回目から第6回目の間即ち5月から7月の間に於ての新感染數が最も多いことであつて第I報に於て論じたところとよく一致して居る。次に前項で一部觸れたように本項で特に注目せられることは凝集反應が此の場合には特異的にマレイン反應並びに補體結合反應と平行して上昇していることである。更に、本報告ではこれらの事を論ずるのが目的でないので詳述は避けるが、此の3種の免疫反應の出現速度について少しく述べてみたい。即ち凝集反應は最も早く出現して其の後急激に減退する傾向がある。その持續期間は私共の例からは平均4ヶ月と推定せられる。例えばTable IVの1號馬を見ると第1回目(7月)1:1600, 第2回目(9月)1:3200, 第3回目(11月)1:3200, 第4回目(翌3月)1:1600であるのを見ると分る。補體結合反應はマレイン反應より幾分早目に出現することは確實のようであるが(5號馬, 8號馬, 11號馬, 16號馬, 19號馬, 21號馬等), まづ殆んど同時と稱して差支えない。前述の如く私共の16ヶ月の調査研究期間中に斃死したものが13頭あるのに對し新感染馬は23頭あること及び次に述べる反應陽性馬より生産された仔馬の本病に對する態度等を對比すると此の地方に於ける鼻疽侵襲の速度は比較的緩慢なことがわかる。

3. 反應陽性馬より産した仔馬の反應

Table Vに見られる様に1944年5月迄に本調査研究の對象馬群に於て98頭の幼駒が生産されたが、(在來種21頭, 改良種77頭)此の中36頭は3月の調査時に生れて居り62頭は5月調査時迄に生れたものであるがまず3月迄に生れた36頭の中反應態度MSの母馬から生れた仔馬は7例でその何れも補體結合反應は母馬と同程度又は幾分低目の反應を示したがマレイン反應は全く陰性であつた。この中2例が5月に於ても尙幾分の抗體を残して居つたが7月に至つて全例陰性となり反應上全く健康であつた。次に3月の調査終了後5月の調査時迄に産した仔馬62頭について見るに反應態度MSの母馬からのものは9例で中1例(3月調査終了直後に産したもの)を除き8例が補體結合反應のみ陽性で此の中5例が7月に於ても尙輕度乍ら陽性を示し11月に

Table V. Reactions observed on the colts born of glanderous mares.

Mares with the reaction of	Colts borne in March (4th investigation)				Colts borne in May (5th investigation)			
	Number	Type of reaction	Examined in		Number	Type of reaction	Examined in	
			Mar.	May			May	July
MS (16)	7	MS	0	0	9	MS	0	0
			7	0			8	0
MS (12)	8	MS	0	0	4	MS	0	0
			0	0			0	0
S (2)	1	MS	0	0	1	MS	0	0
			0	0			0	0
-(68)	20	MS	0	0	48	MS	0	0
			0	0			0	0
Total (98)	36	MS	0	0	62	MS	0	0
			0	0			0	0

Table IV. List of the horses newly infected with glanders.

Period	Horse No.	Original horse No.	Sex	Age	Mallein eye-test					Compl. fixation test					Agglutination test				
1st-2nd	1	71	♂	6	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	2	328	♂	9	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	3	795	♂	7	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	4	877	♂	11	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	5	82	♂	9	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	6	409	♂	5	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	7	823	♂	10	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	8	228	♂	6	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	9	664	♂	8	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	10	40	♂	8	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	11	933	♂	7	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	12	407	♂	4	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	13	237	♂	12	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	14	1081	♀	7	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	15	85	♂	9	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	16	125	♂	14	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	17	87	♂	7	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	18	390	♂	9	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	19	892	♂	5	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	20	1194	♀	10	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	21	180	♂	7	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	22	422	♂	9	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th
1st-2nd	23	1777	♂	5	1st	+	2nd	+	3rd	+	4th	+	5th	+	6th	+	7th	+	8th

Note: 1st-2nd denotes the infection took place during 1st and 2nd investigations.

は全例陰性となつた マレイン點眼反應については前と同様全例はじめから陰性であつた

この事から母馬の血清抗體が胎盤を通じてか或いは母乳を通じてか幼駒の血清中に移行するものであるが、これは大體 2~3 ヶ月にして消失することを知つた 反之マレイン點眼反應の意味に於けるアレルギー性抗體は全く幼駒に移行しないことを知つた 本表から窺い得た最も重要なことは反應陽性馬 (MS) から産した幼駒が何れも病馬である母馬と共に飼育哺乳されて居つて何等感染を受けることなく 6 ヶ月以上も健康に生育し得たことである

III. 總 括

1. 私共は本調査研究開始當初 29 例の開放性鼻疽馬を發見したがこれは全頭數 929 例から見ると 3.2% 強に當る この 29 例について見るに騮馬が最も多く、5 例を除き鼻漏より本菌を證明し、反應上 1 例が S に屬し 28 例が MS に屬したが凝集反應では 1:3200 以上を示したものは 8 例 (27.5%) に過ぎなかつた 年齢的には 11~15 歳迄のものが 50% 強を示した

以上のことは前 2 報告に於て述べた本病に關する見解と全く一致するものであると共に開放性の大部分から純粹に *M. mallei* を分離し得て、菌の檢出不可能であつた例も現地に於いての野外作業であつた關係上雜菌による分離不能であつたことを考慮すると、全例が菌の排出馬として防疫上傳染源として極めて重視さるべきものであることを示すと共に、これら 29 例の開放性鼻疽馬は 1 年 6 ヶ月後には斃死、賣却、不明等による減少で 9 例が残り又新に開放性となつて來たもの等が 7 例加わつて結局 16 例が第 7 回調査時に見られ其の時の總頭數 997 頭に對し 1.6% 強となつていることから全體として此の地方では開放性のものは 1.6% から 3.2% の間を動いているとみられるが、これら開放性のものはこの間他の健康馬と同様勞役に供されている事實は注目すべきことである

2. 次に私共の調査研究期間 (16 ヶ月間) 中に出現した新感染馬は 23 例で、反應上新に感染發症して來る時期としては第 I 報に於て述べたことと一致して 5 月から 7 月の間が最も多い 又これらの例の初期に於いては 3 種の反應が極めてよく平行する事實を知り且つその出現の速度については凝集反應がまづ最初にあらわれ、次で補體結合反應がマレイン反應に幾分先行することを知つた しかし常在地に於ける自然感染馬に於ける凝集反應については其の反應持續期間比較的短く 3~4 ヶ月という結果を示しマレイン反應及び補體結合反應は比較的長期に及ぶこともこれらの例から判斷される

3. 更に反應陽性の母馬からは健康な幼駒を産し得て、6 ヶ月後まで臨牀上又反應上感染發症することなく成育する事實を知つたがこれは非開放性の反應陽性馬は傳染源としては重要な意義をもたないことによるものと解される 反應陽性母馬からの生後 2~3 ヶ月の幼駒の補體結合反應に於ける抗體の證明はそれが胎盤通過か初乳によるものか私

共は分娩直後授乳前の採血を行い得なかつたので遺憾ながら確め得なかつた。これに反しアレルギー性抗體は胎盤を通過しないと共に生後授乳によつても移行せしめ得ないことを認めた。

本稿を終るに臨み御指導御校閲をいただいた越智教授に深謝すると共に小野元處長に敬意を表し又本調査研究に種々御協力を與えられた同僚我妻・鄭の兩氏に感謝する。

(文獻・後出)

EPIZOOTIOLOGICAL OBSERVATIONS ON GLANDERS
IN HORSES IN MANCHURIA. III.
INVESTIGATIONS OF NASAL GLANDERS, NEWLY
INFECTED HORSES AND COLTS BORN
OF THE INFECTED MARES.

N. MURASE, A. JURAKU, J. TIEN, F. SHIMIZU, J. HANG

*(From the former Laboratory of Veterinary Bacteriology, Institute for Horse Diseases,
attached to the Manchurian Science Research Institute, Hsinking, Chief: K. Ono)*

(Received for Publication Jan. 31, 1952)

In the previous papers, it has been concluded that this infectious disease is a typical consumptive one which is largely influenced by the amount of labour and the manner of feeding. The serological reactions and the post-mortem findings obtained were unexpectedly slight and none of the changes were considered fatal.

Findings based on observations made on the history of the cases found with open glanders and on those cases newly infected during our investigations, as well as on the serological behaviours of the colts born of glanderous mares, are referred to in the present paper. The problems treated herein indicate the most important aspects encountered in the control of this infectious disease.

The results obtained are summarized as follows:

1. There were 29 cases found with open glanders out of 929 horses examined in the 1st investigation,—the rate of morbidity being 3.1%. (July, 1943).
2. Among these 29 cases, 22 cases were geldings, and most of them were comparatively aged horses. In addition to this, their immunological behaviours were positive in the mallein test and complement fixation test, except one. (28/29) At the same time, in the agglutination test, their agglutination titres were less than 1:800 in 7 cases, 1:1600 in 12 cases, and more than 1:3200 in 9 cases.

[Japanese Journal of Veterinary Science, Vol. XV, No. 3, 1953]

It is noteworthy that the agglutination titre is not more than 1:800 in such serious open-type cases.

3. In November, 1944, 16 months after the 1st investigation, out of these 29 cases, mentioned above, 8 died, 10 were sold, 1 changed to non-open type, 1 was missing, and other 9 cases remained as before.

Since it is most probable that at least half of the 10 horses sold died, it can be assumed that more than 50% of these 29 cases with open glanders died during the observation period of 16 months. In view of controlling this disease attention should be directed to the fact that the rest might have lived for 2 or 3 years during which period they must have always been disseminating the microorganisms of *M. mallei*.

4. During our investigations, some of those horses classified healthy (—) came in sight as glanderous cases (MS).

Out of nearly 1,000 healthy horses, 21 newly infected cases (2.1%) were detected by our methods (mallein test and complement fixation test), during the period of 16 months and out of these 21 cases, 9 horses developed open-type glanders.

The agglutinating titers of these newly infected cases were more than 1:3200 except for one case.

As to the time that the infection took place, 7 cases were infected during a period ranging from May to July of 1944. It is considered, therefore, that the infection mostly appears periodically, during May to July every year.

5. Careful investigations were undertaken to detect some relation which might exist between the cases with open-type glanders and the newly infected cases, but no appreciable relation was found except for 2 cases.

6. Pregnant mares reacting positively in both tests, gave birth to healthy colts without any difficulties.

These colts showing positive reactions in complement fixation tests, and the agglutination tests made on them showed reactions practically equal, or only slightly lower in degree, to their mothers; also, no sign of positivity in mallein eye-test, within 2–3 months after birth. These antibodies distinguished themselves within the period of 2–3 months.

No sign of infection was recognized even after 8 months had elapsed since their birth.