ELECTRON MICROSCOPY OF AVIAN POX VIRUSES WITH SPECIAL REFERENCE TO THE SIGNIFICANCE OF INCLUSION BODIES IN VIRAL REPLICATION

Masanori TAJIMA and Takeshi USHIJIMA

Nippon Institute for Biological Science, Akebonocho, Tachikawa, Tokyo

(Received for Publication March 24, 1966)

The avian pox viruses have been studied by a number of investigators, using the techniques of thin-section electron microscopy.

MORGAN et al.¹³⁾ distinguished two kinds of viral particles, which seemed to be in different stages of development, in chick chorioallantoic membrane (CAM) infected with fowlpox virus, and proposed a theory of replication based upon the sequestration of viroplasm by incomplete membranes which progressively closed to form viral particles. TAJIMA and KUBOTA^{18,19)} discriminated four forms of viral particles in CAM infected with pigeon pox virus, obtaining such results as supporting the theory of MORGAN et al.¹³⁾ on viral replication. These two groups of authors, however, could not identify electron-microscopically those cytoplasmic inclusions which are comparable to classical inclusion bodies that can be seen by light microscopy. On the other hand, inclusion bodies were identified by EAVES and FLEWETT⁶⁾ and PETERS¹⁵⁾ in CAM infected with fowlpox virus, by ARHELGER et al.¹⁾ in chick scalps infected with the same virus, by HERZBERG and KLEINSCHMIDT⁹⁾ and HERZBERG et al.¹⁰⁾ in histiocytes of canaries infected with canary pox virus, and by BEAVER and CHEATHAM³⁾ in CAM infected with an avian pox virus isolated from a slate-colored junco.

WOODRUFF and GOODPASTURE^{22,23)} demonstrated that the Bollinger body of fowlpox is a minute colony of the etiological agents of this disease. Since then, it has generally been considered that the Bollinger body is a site of viral replication. RANDALL and GAFFORD¹⁶⁾ presumed from their histochemical studies on isolated inclusions that the inclusion must be the site of a very active and complex biochemical nidus. Recently BEAVER et al.⁴⁾, who performed an electron microscopic study on the CAM infected with fowlpox virus, stated that the Bollinger body is the only area within the cell where there is an unquestionable gradual transition between undifferentiated and mature viral particles.

On the contrary, KATO et al.¹¹⁾ have described two types of cytoplasmic inclusions in the ascites free cells of chickens inoculated intraperitoneally with fowlpox virus. They designated as type A the inclusion which corresponds to the Bollinger body, and as type B another cytoplasmic inclusion. They stated that type B inclusion plays a principal role in the process of multiplication of this virus, and apparently is the site providing the nucleoproteins needed for the synthesis of viral elementary bodies, and that the Bollinger body is mere reservoirs of mature virus. It thus appears that the relationship of the inclusion body to viral replication still remains a subject of controversy in spite of a long history of studies on it.

The present paper deals with the ultrastructure and development of viral particles and cytoplasmic inclusions associated with lesions of fowlpox, canary pox, and pigeon

pox, which have been regarded as different viruses or strains of the avian pox group. In addition, it contains discussion on the relationship of inclusions to viral replication.

MATERIALS AND METHODS

Viruses: Two strains, NF-1 and B-36, of fowlpox virus were used for inoculation of eggs and chickens. At the time of inoculation, the NF-1 and B-36 strains were at the 2nd and 5th egg passage, respectively.

Canary pox virus was employed for inoculation of eggs and canaries at its 6th egg passage level after it had been isolated (MIYAMOTO¹²⁾) from a spontaneous outbreak of canary pox among the canaries. The same virus passed 16 times in chick-embryo cell culture was used for tissue culture experiments.

The Nakano strain of pigeon pox virus was employed for egg inoculation at its 4th egg passage level.

Egg- and bird-inoculation: In egg-inoculation, dropped CAMs of 11-day-old chick embryos were inoculated with 0.1 ml of each of 10^{-3} to 10^{-5} dilutions of a suspension of CAM infected with one of the viruses of fowlpox, canary pox, and pigeon pox. The inoculated embryos were reincubated at 35°C for 2 to 6 days.

Four 56-day-old cockerels were inoculated with 1:10 dilution of CAM infected with fowlpox virus by scarification of the comb.

Five adult female canaries derived from a flock which had never been involved in an outbreak of canary pox were inoculated with canary pox virus by gently rubbing a 1:10 dilution of emulsified infected CAM on the freshly defeathered follicles of the breast.

Propagation of canary pox virus in chick-embryo cell culture: The methods described by SARO et al.¹⁷⁾ were applied to the preparation of the cell culture used and the infection of cultures. In this experiment, each tube culture was inoculated with 0.1 ml of 10^{-1} dilution of infected culture fluid. The highest infective titer determined in the culture fluid of these cultures reached $10^{5.0}$ TCID₅₀ per 0.1 ml 5 days after inoculation.

Preparation of specimens for morphological observations: Egg material was harvested daily from 2 to 6 days after inoculation, chicken material 5 and 6 days, and canary material from 9 to 11 days, after inoculation. Discrete pocks or confluent lesions had developed at the site of inoculation by the time of harvesting. In infected cell cultures, harvesting was carried out 2, 3, and 5 days after inoculation when distinct cytopathic effects were observed.

All the materials harvested were fixed with PALADE's fluid¹⁴⁾, dehydrated with graded alcohols, and embedded in methacrylate. Thin sections were cut with glass knives on a JUM type 3 microtome. These sections were examined in a JEM-4C electron microscope, without staining or after staining with lead hydroxide or uranyl acetate by WATSON's method^{20,21)}.

When materials for electron microscopy were harvested, those for light microscopy were collected from the same tissues or cultures, so that interpretation of electron micrographs might be facilitated. They were fixed with formalin or Zenker's fluid and stained with hematoxylin and eosin or by Giemsa's method.

RESULTS

Morphology of Viral Particles

Particles of all the types of avian pox virus described by previous workers were easily identified in any type of cells infected with any virus of avian pox examined. Those viral particles examined of fowlpox, canary pox, and pigeon pox were similar to one another with regard to their size and structure. Moreover, it was impossible to find any difference in these regards between the two strains of fowlpox virus. Based their morphological characteristics, the viral particles were classified into five types.

Incomplete forms: These forms were composed of an incomplete limiting membrane and a moderately dense, finely granular and fibrillar viroplasm (Figs. 1, 8 and 13). They were situated in the cytoplasm of infected cells in close association with matrix inclusions which will be mentioned below. It was interpreted that these forms indicated the first stage of viral condensation.

Spherical forms: These forms were round to oval in shape. They had a complete limiting membrane with double structure and such viroplasm as resembling that of the incomplete forms in appearance and density to electron beams (Figs. 1, 2, 8, 10, 13, 14 and 16). They contained no visible nucleoid. It cannot be denied, however, that some spherical forms may possibly be early intermediate ones which have been cut tangentially and which, therefore, are beside the nucleoid. It is presumed that the spherical forms have been transformed from the incomplete ones as a result of accomplishment of the formation of the limiting membrane. Those forms were present in close association with matrix inclusions and also scattered singly or in clusters all over the cytoplasm.

Early intermediate forms: These forms were the same in morphology as the spherical forms, except that the former had a nucleoid, which was an electron-dense, eccentric internal body measured to be $70 \times 95 \text{ m}\mu$ on the average (Figs. 8 and 16). The nucleoid was separated from the viroplasm by a zone of lesser density. These forms were seen in the cytoplasm of infected cells, together with the other developmental forms.

Later intermediate forms: These forms were a little more elongated in shape than the forms mentioned above. In them, the nucleoid increased in volume and decreased in density, occupying the central position of a particle (Figs. 2 and 14). The zone of low density surrounding the nucleoid became so wide that it gave an impression that the nucleoid had increased in size at the expense of granular or fibrillar material in the viroplasm. These forms were observed in the cytoplasm of host cells, together with the other developmental forms.

Mature forms: It was assumed that these forms might be disk-shaped and their nucleoid take a shape of biconcave disk when observed three-dimensionally. Therefore, it might well be that various pictures were shown according to the differences in plane and angle at which sections had been cut. In a cross-section of the disk, an electron-dense, dumbbell-shaped nucleoid about $208 \text{ m}\mu$ in length occupied the center of a particle. In the nucleoid was seen an axis of lesser density containing granular or filamentous material. The nucleoid was surrounded by a less dense zone, and the outermost side of the particle was covered by an outer coat with multiple layers. Two electron-dense protuberances of the outer coat were seen projecting against both concavities of the nucleoid. When a cross-section was made in such a manner as to be parallel to the flat surface of the disk, the particle and its nucleoid exhibited an outline which was oval or sometimes rectangular (Figs. 7 and 11). The less dense axis of the nucleoid and the less dense zone surrounding the nucleoid in the mature forms were often seen remarkably enlarged in degenerated cells located in the proliferated ectoderm of the CAM of

embryonated hen's eggs and in the outer portion of the epidermic layer of chickens and canaries (Fig. 11). The viral particles contained in a classical inclusion and those lying in the extracellular space had morphology characteristic of the mature particles. In the cytoplasm of infected cells, mature particles were also scattered individually or in clusters, together with the other developmental forms.

It was interpreted that the incomplete forms revealed the first stage of the maturation of viral particles and that the spherical, early intermediate, and later intermediate forms were forms of the transitional stages leading to mature forms which were in the terminal stage of development. Consequently, the four forms mentioned above can be called the immature or developmental forms in contrast to the mature forms.

In Table 1 are shown the average diameters of the developmental and the mature forms of the avian pox viruses observed in thinly sectioned cells. There were differences of some degree in the size of particle among the kinds of viruses and among the viruses grown in host cells of different types. Nevertheless, these differences were not deemed to be significant, since there was a considerable individual variation in the size of particle.

Table 1. Dimensions of Avian Pox Viruses (All Measurements in Millimicrons)

Virus	Average diameter of particle developed in		
	chorionic ectodermal cell	dermal epithelial cell of natural host	cultured chick- embryo cell
Fowlpox			
Developmental form	265×312	241×279	
Mature form	153×225×277	164×252×284	
Canary pox			
Developmental form	245×315	266×303	250×327
Mature form	155×224×290	176×260×320	155×202×263
Pigeon pox			
Developmental form	252×310		
Mature form	140×236×289		

Ultrastructure of Inclusion Bodies

It was possible to distinguish two structures, which were regarded as localized manifestations of viral activity, in the cytoplasm of infected cells by using the electron microscope. One of them was easily identified as a structure corresponding to classical inclusion bodies when studied comparatively by light microscopy (Figs. 4, 7, 9~12 and 17). These bodies were observed in large numbers in all the types of cell examined, except cultured chick-embryo cells infected with canary pox virus. The other structure was not exactly identified by light microscopy, so far as the methods of fixation and staining used in the present studies are concerned, but was observed electron microscopically as areas of dense granular and fibrillar material (Figs. 1, 8, and 13). In many cases, these areas were surrounded by such various developmental forms as mentioned above. These areas resembled the matrix areas observed by GAYLORD and MELNICK⁸⁾ in the cytoplasm of host cells infected with the other members of the pox group. Accordingly, the authors want to propose that this structure be called the matrix inclusion hereafter. The matrix inclusion has often been observed in all the types of host cells infected with the three viruses examined.

In Table 2 is shown the state of occurrence of inclusions of both types in various host cells infected with the avian pox viruses.

The classical inclusion was composed of a large number of mature forms and stroma,

Table 2. Occurrence of Inclusion Bodies in Various Host Cells Infected with Avian Pox Viruses

Host cell	Inclusion body*					
	Fowlpox		Canary pox		Pigeon pox	
	C	M	C	M	C	M
Dermal epithelial cell of natural host	P	P	P	P	.	.
Chorionic ectodermal cell	P	P	P	P	P	P
Cultured chick-embryo cell	.	.	N	P	.	.

* C: classical inclusion, M: matrix inclusion, P: positive, N: negative, and : not tested.

or ground substance, which exhibited a complicated structure and which filled space among these forms. In the stroma were distinguished electron-dense amorphous material, dense granules 10 to 30 m μ in diameter with an irregular outline, threads with a diameter and density of the same order as these dense granules and with the maximum length reaching 300 m μ , and tubulous structures 15 to 45 m μ in diameter and 200 m μ in length.

It seemed that there was a difference in structure between classical inclusions according to the type of host cells in which these inclusions had been developed rather than according to the kind of viruses. In ectodermal cells of the CAM of embryonated hen's eggs, regardless of the virus with which these cells had been infected, classical inclusions were generally rich in electron-dense stroma and particularly had a dense, irregularly-shaped core that had been formed in the center by compact accumulation of amorphous material. Viral particles were rarely or not observed in the core. The core was surrounded by a zone where numberless dense granules and threads were scattered. Outside this zone there was an area rich in viral particles. In this outermost area, viral particles showed the typical structure of the mature forms, and most of them were arranged in such a manner that their long axes might be parallel to the surface of the inclusion. On the other hand, most of the viral particles lying in the granular zone of the inclusion were poorly defined. In them, the nucleoid and the outer coat increased in granularity and lost their fine structure, giving an impression that they were in the process of disintegration (Figs. 5~7, 12 and 17). Some inclusions were found to be covered with a limiting membrane, the origin of which was unknown, partly or entirely at their circumference.

Classical inclusions appearing in dermal epithelial cells of chicken and canary had the same constituents as those mentioned above. It was generally impossible, however, to discriminate a core, a granular zone, and a peripheral zone in them. They had only mature forms which exhibited a typical structure and which were dispersed evenly all over the inclusion. In the inclusion, the dense amorphous material of the stroma was small in amount and filled the space between viral particles loosely. Accordingly, it was possible to differentiate the viral particles and the limiting membrane of the inclusion with ease and distinctness (Figs. 4 and 9~11). Inclusions of large size found in older lesions had a looser structure with less dense stroma, containing sometimes large clefts or vacuoles (Figs. 9 and 11). In inclusions of smaller size and ill-defined aggregations of mature forms, space between viral particles was filled compactly with granular and fibrillar material which resembled that contained in the matrix inclusion described below in size and density. In this space dense amorphous material was scanty or absent (Figs. 3 and 10). Generally, these aggregations of mature forms or inclusions were contained in well-preserved cells in younger lesions or the inner portion of the proliferated epidermic layer. Therefore, it was interpreted that they were in the early stages of inclusion

formation. In cultured cells infected with canary pox virus, the mature forms were scattered throughout the cytoplasm, together with the developmental forms, and there was no formation of any well-defined structure corresponding to classical inclusion body (Fig. 14).

The matrix inclusion appeared in the cytoplasm of the host cell as a dense area consisting of short fibrils 3 to 10 m μ in diameter and granules 10 to 17 m μ in diameter (Figs. 1 and 8). The axis of a large one of these areas reached 3 μ in length. It was possible to find one to several matrix inclusions in the cytoplasm of a single cell. It was not seldom for a cell to contain both matrix inclusion and classical body in its cytoplasm. In dermal epithelial cells of chicken and canary and ectodermal cells of the CAM of embryonated hen's eggs, the matrix area was compact, dense, and well-defined. Elaboration of limiting membranes of viral particles occurred almost exclusively in the peripheral portion of the area (Figs. 1 and 8). On the other hand, cultured cells infected with canary pox virus had a matrix area loose, less dense, and ill-defined, giving the impression that the accumulation of matrix material was small in amount. In these cells, the condensation of limiting membranes of virus particles generally occurred within the area of dense material (Fig. 13).

Cytopathology of Host Cells

A remarkable enlargement was generally observed among infected epithelial cells of the skin of chicken and canary and infected ectodermal cells of the CAM of embryonated hen's eggs. In these cells nuclei were often found distorted. Moreover, they were displaced to the peripheral portions of the cells when inclusions were contained in the cytoplasm. In the nucleus, the nucleolus was swollen, and the tangled-ribbon configuration described by BEAVER and CHEATHAM³⁾ was not infrequently present. The cytoplasm became less dense as a result of dispersion of the organelles. The mitochondria and endoplasmic reticulum were enlarged and underwent varying degrees of disintegration. In the cytoplasm containing inculsions, the mitochondria and other cytoplasmic elements appeared to have been pushed away to the peripheral portion of the cytoplasm.

In chick-embryo cell cultures, cytopathic changes began to appear two days after addition of virus to the cultures. In stained preparations taken at this period, discrete foci containing cells having a retraction of both the cytoplasm and the nucleus, with increasing basophilia were observed. Some of these cells contained such eosinophilic, well-defined structure as apparently resembling classical inclusion. Such a structure as this, however, was not rarely observed in uninoculated controls. During the 3 to 5 days following infection, the cells with the described changes increased in number until the culture was almost completely destroyed. A large number of cells with their cytoplasm damaged and their nucleus distorted were observed by electron microscopy. These changes were generally pronounced in cells containing viral particles in the cytoplasm. Sometimes in these cells were seen such dense structures as presumed to be comparable to the eosinophilic bodies which had been recognized by light microscopy. These structures resembled classical inclusion bodies in size and appearance. No viral particles, however, were identified in these structures, in spite of a careful study was performed on the latter (Fig. 15).

DISCUSSION

It seems that the concept "viral inclusion" has recently been extended with the application of electron microscopy. As pointed out by FLEWETT⁷⁾, the term viral inclu-

sion may well be used to express any localized intracellular manifestation of viral activity. In the present investigation, two types of inclusions, matrix and classical inclusions, were distinguished in the cytoplasm of infected cells.

Matrix inclusions corresponded to the aggregate of fine granular and fibrillar material which was observed for the first time by MORGAN et al.¹³⁾ in cells infected with fowlpox virus, and later by other investigators in cells infected with the other members of the pox group. They were found in all types of host cells infected with the viruses of fowlpox, canary pox, and pigeon pox. Most of the areas of dense granular and fibrillar material were closely associated with the developmental forms. In the present studies, no attempt was made to clarify the sequence of adsorption and penetration of viral particles into host cells. It seems unquestionable, however, that these areas represent the first stage of virus condensation, as was reported by DALES and SIMINOVITCH⁵⁾ who had carried out sequential studies using the vaccinia-L cell system. The incomplete forms which develop around or within the dense material of the matrix inclusion may eventually grow into mature forms, passing through the stages of spherical, early intermediate and later intermediate forms in the order listed. This possible process of transformation can be easily assumed, since many transitional forms are present among the forms mentioned above. BEAVER et al.⁴⁾ found such particles as resembling the aforesaid developmental forms in mesodermal macrophages of the CAM infected with fowlpox virus. From their findings, they presumed that these particles might be degradation product rather than those representing a phase of the life-cycle of fowlpox virus. In this case, however, it will not be able to deny a possibility that fowlpox virus may propagate even in mesodermal macrophages.

The classical inclusion mentioned above is identical with the Bollinger body of fowl pox which has been well known light-microscopically. Its ultrastructure is not different from that described by earlier workers. It was interpreted from the results of the present investigation that the classical inclusion might be formed by the aggregation of viral particles which have developed and passed through the process of maturation in the matrix inclusion. Some of the dense granular material constituting the stroma of an early classical inclusion resemble the granular material of a matrix inclusion morphologically. This fact suggests that it is a waste product derived from the material accumulated in the matrix inclusion for the formation of viral particles. The stroma of a large, probably older classical inclusion had a more complicated structure, containing a large amount of dense amorphous material. In the light of the results obtained by RANDALL and GAFFORD¹⁶⁾, it is presumed that lipid may have been added secondarily to the stroma of the classical inclusion. The present studies gave results suggesting that some of the mature forms contained in the center of a large classical inclusion had undergone degradation. These findings agreed with the observations made by ARHELGER et al.¹⁾

In cultured chick-embryo cells infected with canary pox virus, the maximum infective titer determined in the culture fluid reached $10^{5.0}$ TCID₅₀ per 0.1 ml. No classical inclusions were detected at all in these cells, although matrix inclusions and numerous mature forms were observed by electron microscopy. These results are similar to those obtained by BANG et al.²⁾ who cultivated fowlpox virus in roller-tube cultures of chick fibroblasts or chick epithelial cells. These facts indicate that a classical inclusion is not always a structure indispensable for the development and maturation of the avian pox viruses. SATO et al.¹⁷⁾ reported that cytoplasmic inclusion bodies, stained reddish pink by Giemsa's method, appeared in cultured chick-embryo cells infected with canary pox virus. They did not ascertain, however, whether these inclusion bodies contained viral particles or not. At present, it is impossible to find any exact reason why classical inclusion have failed to develop in those cultured cells, but the following factors may

be available for the explanation of these findings. (1) The number of viral particles per infected cultured cell was apparently smaller than that per host cell of any other type. (2) Most of the infected cultured cells suffered severe damage within five days after infection. Consequently, there may have been no sufficient time for these cells to allow classical inclusions to develop. (3) In infected cultured cells, the amount of granular and fibrillar material accumulated in the matrix inclusion was relatively small. Accordingly, it may well be that it was impossible for this material to produce sufficient waste products to form the stroma of the classical inclusion.

The present investigation could not provide any evidence that the matrix inclusion described in this paper might be identical with the type B body discovered by KATO et al.¹¹⁾ in cells infected with fowlpox virus by means of light microscopy. It seems, however, that both structures resemble each other with regard to a point that either inclusion body plays an important role in the process of viral replication.

SUMMARY

The viruses of fowlpox, canary pox, and pigeon pox grown in various types of host cells were studied by the technique of thin-section electron microscopy. Viral particles were identified in all the types of host cells examined. They were classified into five forms, incomplete, spherical, early intermediate, later intermediate, and mature. It was interpreted that of them, the first four were immature or developmental forms, and that the mature forms were in the terminal stage of development. Two types of cytoplasmic inclusion bodies were distinguished in infected cells. The body of one type was identical with the classical inclusion which was visible under the light microscope, and consisted of a large number of mature forms and electron-dense stroma. It was easily identified in all the types of host cells used, except cultured cells infected with canary pox virus. The body of the other type was represented by a dense cytoplasmic area composed of fine granular and fibrillar material. In general, it was associated with various kinds of developmental forms and designated the matrix inclusion. Matrix inclusions were demonstrated in any type of host cells infected with what-ever avian pox virus. They were considered to play an important role in the process of viral replication. It was interpreted that classical inclusions might be formed by the aggregation of viral particles which accomplished maturation in matrix inclusions and by a secondary addition of dense material, most of which might probably be lipid. There were no distinct differences among the viruses of fowlpox, canary pox, and pigeon pox in the morphology of viral particles or in the apparent mechanism of viral replication. Except for the presence of viral particles and cytoplasmic inclusion bodies, the cellular alterations observed in the infected cells could not be considered as specific for viral infection.

ACKNOWLEDGMENTS

The authors wish to thank Dr. J. NAKAMURA, director, Nippon Institute for Biological Science, for his advice and Drs. T. MIYAMOTO and T. MOTOHASHI, staff members of the same institute, for their assistance.

This report is dedicated to Dr. S. YAMAGIWA, the president of Obihiro Zootechnical University (1962—), Professor Emeritus in Hokkaido University, Ex-professor in the Department of Comparative Pathology, Faculty of Veterinary Medicine, Hokkaido University.

The outline of this study was presented before the Fifty-Fifth Meeting of the Japanese Society of Veterinary Science, Tokyo, April 8, 1963.

REFERENCES

- 1) ARHELGER, R. B., DARLINGTON, R. W., GAFFORD, L. G. and RANDALL, C. C. (1962): *Lab. Invest.*, **11**, 814~825.
- 2) BANG, F. B., LEVY, E. and GEY, G. O. (1951): *J. Immunol.*, **66**, 329~345.
- 3) BEAVER, D. L. and CHEATHAM, W. J. (1963): *Amer. J. Path.*, **42**, 23~39.
- 4) BEAVER, D. L., CHEATHAM, W. J. and MOSES, H. L. (1963): *Lab. Invest.*, **12**, 519~530.
- 5) DALES, S. and SIMINOVITCH, L. (1961): *J. biophys. biochem. Cytol.*, **10**, 475~502.
- 6) EAVES, G. and FLEWETT, T. H. (1955): *J. Hyg. (Lond.)*, **53**, 102~105.
- 7) FLEWETT, T. H. (1956): *J. Hyg. (Lond.)*, **54**, 393~400.
- 8) GAYLORD, W. H. and MELNICK, J. L. (1953): *J. exp. Med.*, **98**, 157~171.
- 9) HERZBERG, K. and KLEINSCHMIDT, A. (1959): *Zbl. Bakt., I. Abt. Orig.*, **174**, 1~27.
- 10) HERZBERG, K., KLEINSCHMIDT, A. and LANG, D. (1960): *Zbl. Bakt., I. Abt. Orig.*, **179**, 308~324.
- 11) KATO, S., HAGIWARA, K., BABA, E., SATO, Y. and KAMAHORA, J. (1955): *Virus*, **5**, 318~324 (in Japanese with English abstract).
- 12) MIYAMOTO, T. (1959): *Jap. J. vet. Sci.*, **21**, 77 (Abstract in Japanese).
- 13) MORGAN, C., ELLISON, S. A., ROSE, H. M. and MOORE, D. H. (1954): *J. exp. Med.*, **100**, 301~308.
- 14) PALADE, G. E. (1952): *J. exp. Med.*, **95**, 285~298.
- 15) PETERS, D. (1959): *Zbl. Bakt., I. Abt. Orig.*, **176**, 259~295.
- 16) RANDALL, C. C. and GAFFORD, L. G. (1962): *Amer. J. Path.*, **40**, 51~62.
- 17) SATO, T., SUGIMORI, T. and MATUMOTO, M. (1962): *Jap. J. exp. Med.*, **32**, 263~277.
- 18) TAJIMA, M. and KUBOTA, Y. (1956): *Virus*, **6**, 308~317 (in Japanese with English abstract).
- 19) TAJIMA, M. and KUBOTA, Y. (1957): *NIBS Bull. biol. Res.*, **2**, 48~57.
- 20) WATSON, M. L. (1958a): *J. biophys. biochem. Cytol.*, **4**, 475~478.
- 21) WATSON, M. L. (1958b): *J. biophys. biochem. Cytol.*, **4**, 727~730.
- 22) WOODRUFF, C. E. and GOODPASTURE, E. W. (1929): *Amer. J. Path.*, **5**, 1~10.
- 23) WOODRUFF, C. E. and GOODPASTURE, E. W. (1930): *Amer. J. Path.*, **6**, 713~720.

禽痘ウイルスの電子顕微鏡的研究 特にウイルス増殖における封入体の意義について

田島正典・牛島毅

日本生物科学研究所

(昭和 41 年 3 月 24 日受付)

鶏痘、カナリヤ痘および鳩痘のウイルスを種々の細胞に感染させ、超薄切片法により電子顕微鏡をもって研究した。ウイルス粒子は検索した総ての種類細胞において同定され、それらは不完全、球形、早期中間、後期中間および成熟の 5 型に分類された。これら 5 型のうち前の 4 型は未熟あるいは発育型と、成熟型はウイルス発生の終末段階にある粒子と考えられた。

感染細胞には 2 種の細胞質封入体が観察された。その 1 つは光学顕微鏡で見られる古典的封入体に一致し、多数の成熟型粒子と高電子密度の基質からなり、カナリヤ痘ウイルスに感染した培養細胞を除き、

他の総ての種類宿主細胞において容易に同定された。

他の 1 つはマトリックス封入体と名づけられたもので、微細顆粒状および微小線維性物質からなり、細胞質において、電子密度の高い部域として認められ、通常、種々の発育型粒子と共存した。この封入体は総ての宿主細胞—ウイルス系において証明され、ウイルスの増殖過程において重要な役割を演ずるものと解釈された。

古典的封入体はマトリックス封入体で発生分化した成熟型粒子の集簇と、その大部分はリピドと考えられる高電子密度の物質の二次的加入によって形成

されるもので、ウイルス増殖に対して必須な構造ではないと見られた。

鶏痘、カナリヤ痘および鳩痘のウイルス粒子の間には、それらの形態と発生様式において明確な差異

はないようであった。ウイルス粒子と細胞質封入体の存在を除けば、ウイルス感染に対して特異的と考えられる細胞病変は見出されなかった。

EXPLANATION OF PLATES

Figures 1 to 4 were taken from sections of the comb epidermis of chickens infected with the B-36 strain, and figures 5 to 7 from sections of chick chorio-allantoic membranes infected with the NF-1 strain of fowlpox virus. Figures 8 to 11 were taken from sections of the canary epidermis, figure 12 was from a section of chick chorio-allantoic membrane, and figures 13 to 15 were from sections of cultured chick-embryo cells infected with canary pox virus. Figures 16 and 17 are micrographs taken from sections of chick chorio-allantoic membranes infected with the Nakano strain of pigeon pox virus.

PLATE I

- Fig. 1. A portion of the cytoplasm of an epidermic cell 5 days after inoculation. A large matrix inclusion is shown in the center of the picture. Several incomplete limiting membranes are being formed in the periphery of an area of dense granular and fibrillar material. On the upper side of the area are seen many spherical forms. Part of a classical inclusion occupies the lower border of the picture. $\times 24,000$.
- Fig. 2. A portion of the cytoplasm of an infected cell at 5 days, containing a group of developmental forms, most of which have complete limiting membranes. One particle in the lower part of the group contains a dense, large nucleoid centrally located. $\times 44,000$.
- Fig. 3. Part of the cytoplasm of an infected cell at 5 days, containing a group of viral forms, most of which possess the morphology characteristic of the mature forms. They are embedded in dense granular and fibrillar material similar in appearance to that seen in the matrix inclusion. $\times 44,000$.

PLATE II

- Fig. 4. Section of the epidermic layer at 6 days, showing two large classical inclusions. Numerous mature forms are clearly visible within the inclusions. Viral forms in various stages of differentiation are also scattered, either singly or in clusters, throughout the cytoplasm surrounding the inclusions. $\times 5,000$.
- Fig. 5. A portion of an ectodermal cell 6 days after inoculation, containing a large, irregularly shaped inclusion body. The inclusion appears to be composed of two dense, amorphous central areas surrounded by a less dense cortical zone which contains numerous mature forms. The nucleus is seen in the lower right corner. $\times 13,000$.

PLATE III

- Fig. 6. Part of an ectodermal cell at 6 days, containing another type of classical inclusion. The center of the inclusion is occupied by amorphous material in which dense granules and threads are scattered. Viral forms located near the center of the body are poorly defined, possessing a dense, granular outer coat. The nucleus is present in the lower left corner. $\times 26,000$.
- Fig. 7. Higher magnification micrograph of a portion of the cortical zone of a classical inclusion similar to that shown in Fig. 6. Viral forms contained exhibit the morphology characteristic of the mature forms. Dense threads and tubular structures are clearly visible lying among the viral forms. $\times 64,000$.

PLATE IV

- Fig. 8. Part of the cytoplasm of an epidermic cell 9 days after inoculation, containing two matrix inclusions. Incomplete forms are shown forming around areas of dense granular and fibrillar material. Around the inclusions are seen many developmental forms, one of which contains a dense eccentric nucleoid. $\times 33,000$.
- Fig. 9. Section of the epidermic layer at 11 days, containing four large, round classical inclusions. Numerous viral forms are evenly distributed throughout the inclusions. The host cells appear to be at an advanced stage of degeneration. $\times 4,000$.

PLATE V

- Fig. 10. A portion of an epidermic cell at 11 days, containing a classical inclusion presumed to be at an early stage of its development. All viral forms contained possess the morphology characteristic of the mature forms. They are embedded in dense granular components which bear some resemblance in appearance to those contained in matrix inclusions. Several developmental forms are seen in the lower left corner. The nucleus is in the upper right corner. $\times 22,000$.
- Fig. 11. A portion of a classical inclusion is shown at a higher magnification. Two mature forms near the lower border of the picture have been sectioned transversely and the others sagittally. In viral forms cut along an appropriate plane, a characteristic dumbbell-shaped nucleoid occupies their center and is surrounded by a less dense zone. The less dense zone is surrounded by a dense granular layer which has been thickened on either side of the viral form. The outer coat of the viral form appears to consist of multiple membranes. $\times 100,000$. Inset shows two mature forms seen in another classical inclusion contained in a markedly degenerated cell in the superficial layer of the epidermis. The inner core of the nucleoid and the clear space surrounding the nucleoid have been remarkably expanded, giving an impression that these forms have undergone some drying effect. $\times 80,000$.

PLATE VI

- Fig. 12. A large classical inclusion in the cytoplasm of an ectodermal cell at 5 days. Numerous osmophilic granules and amorphous material occupy the central area. Viral forms are mainly distributed in the peripheral zone. $\times 19,000$.
- Fig. 13. A portion of a cultured cell sampled 3 days after infection and containing a matrix inclusion. Developmental forms in various stages of differentiation are embedded in dense fibrillar material. Most of the incomplete membranes appear to have occurred within the dense material. Several swollen mitochondria lie around the inclusion. The nucleus occupies the left border of the picture. $\times 19,000$.

PLATE VII

- Fig. 14. Part of the cytoplasm of a cultured cell sampled 3 days after infection and containing viral forms which are in various stages of differentiation. Every transitional form between the spherical and the mature forms can be observed. $\times 30,000$.
- Fig. 15. Part of the cytoplasm of a cultured cell at 3 days, containing a large dense structure reminiscent of a classical inclusion. Such structure, however, contains no recognizable viral forms. It was also observed in uninoculated controls. $\times 15,000$.

PLATE VIII

- Fig. 16. A portion of an ectodermal cell 4 days after inoculation, containing various viral forms in the cytoplasm. They appear to be in the process of coalescence with such irregularly shaped,

dense, amorphous material as seen near the center of the field. Near the left border of the picture are seen several swollen mitochondria. The nucleus is in the upper right corner. $\times 19,000$.

- Fig. 17. Part of the cytoplasm of an ectodermal cell at 4 days, containing an irregularly shaped, dense classical inclusion. The greater part of the inclusion is occupied by osmiophilic amorphous material. Viral forms are visible only in a narrow peripheral zone of the inclusion. $\times 26,000$.