BRIEF NOTE

**Hepatosplenic Myelosis in Naturally Occurring Mouse Hepatitis
Virus Infection in the Nude Mouse**

Takuo ISHIDA, Teruko TAMURA, Katsumoto UEDA
and Kôzaku FUJIWARA

*Department of Animal Pathology, Institute of Medical Science,
University of Tokyo, Minato-ku, Tokyo 108*

(Received for publication April 26, 1978)

The etiologic association of mouse hepatitis virus (MHV) with wasting syndrome in the athymic nude mouse has been well documented, and involvement of lymphoid and hemopoietic tissues resulting in a modification of immune response has been reported [5-9]. This communication describes bone marrow involvement with extensive extramedullary hemopoiesis in nude mice naturally infected with MHV.

Adult male and female nude [nu/nu, BALB/c] mice and their haired heterozygous littermates [nu/+, BALB/c] were obtained from a national institute colony where a high incidence of wasting in nude mice was recognized. Twenty-seven nude mice were necropsied, and hepatic lesions and splenomegaly were recognized in 10 mice. Hepatic lesions were seen in another group of 4 mice without prominent splenomegaly. Grossly, the liver showed irregular white spots 1 to 2 mm in diameter on the surface, and granular appearance in some cases. No gross lesion was observed in the 9 haired littermate mice supplied simultaneously.

Histopathology revealed hepatic multi-

focal necrosis with multinucleated giant cells and eosinophilic bodies almost identical to those reported in MHV-NuU infection [8]. By direct immunofluorescence [4], MHV antigen was demonstrated in the hepatocytes as well as multinucleated giant cells at the margin of necrotic foci. Furthermore, the affected livers showed varying degrees of extramedullary hemopoiesis with a number of myeloid cells appearing in the sinusoids and around large blood vessels. Degenerated multinucleated giant cells were occasionally seen within those hemopoietic foci (Figs. 1, 2).

In the spleen, the red pulp was found greatly enlarged and mostly replaced by myeloid cells mainly of granulopoietic series. A considerable increase in number of megakaryocytes was also observed, and many of them appeared dyspoietic (Figs. 3, 4). Multinucleated giant cells and their degenerative figures were seen in the red pulp. Karyorrhexis and karyolysis of megakaryocytes were also noted. The area of the white pulp was reduced, and necrosis of lymphocytes and reticular cells was observed. Plasma cells were frequently seen

短報: マウス肝炎ウイルス自然感染ヌードマウスの肝脾骨髓症について: 石田卓夫・田村照子・上田雄幹・藤原公策 (東京大学医科学研究所獣医学研究部)

in the vicinity of the central arteries.

The above hepatosplenic hemopoiesis was almost always accompanied by bone marrow lesions. The changes in the marrow consisted of anaplasia, necrosis and syncytium formation (Fig. 5). The lesions in the paracortical area of the lymph nodes were similar to those reported previously [8]. Syncytium formation and reticular cell degeneration were commonly seen, while lymphoid necrosis was rare. No pneumonia was shown in the wasted nude mice.

Serum samples were taken from 16 nude mice and 9 haired littermates, and they were tested for complement fixing (CF) antibodies by the microtiter method [2]. No samples from nude mice were positive for CF antibodies against either MHV or Sendai virus. However, in the haired mice, 4 out of 9 (44%) were positive for MHV and other 2 (22%) were positive for Sendai virus.

Clarified 10% tissue homogenate prepared from the affected livers and spleens of the nude mice was inoculated into 48-hr culture of SR-CDF₁-DBT (DBT⁺) cells [3, 4]. A virus which produced syncytium formation in DBT cells was isolated from the livers and spleens, although the titers in the spleen were much lower than those in the liver. By direct immunofluorescence, MHV antigen was demonstrated in DBT cells infected with the isolated virus. Immunodiffusion [4] revealed cross reaction between the isolate, MHV-NuU and MHV-2.

To study the pathogenicity of the isolated virus, 5 nude mice [CLEA:nu/nu, BALB/cA] and 10 haired littermates [CLEA:nu/+, BALB/cA], from an MHV-free colony, were inoculated with 1×10^5 plaque forming units/0.2 ml [3] of the liver isolate at 5th passage on DBT cells. Four nude mice died with wasting syndrome at

10 to 20 days after infection all having hepatitis and bone marrow lesions as observed in the spontaneous cases. In these mice, MHV antigen was demonstrated in the liver, spleen, lymph node and bone marrow. Hepatosplenic hemopoiesis was observed in 3 mice died at 15 and 20 days after infection. The surviving one was killed at 30 days after inoculation, and it manifested focal hepatic necrosis and splenic syncytium formation. The haired littermates showed no clinical signs nor died, but had CF antibodies to MHV at titers of 1:5 or higher when examined 14 and 21 days after infection. There were no positive cases for Sendai virus antibodies.

These results suggest that the present isolated virus, tentatively designated as MHV-Kyoto strain, is capable of producing bone marrow disorders as well as hepatitis. Furthermore, coincidence of hepatosplenic hemopoiesis and marrow damages suggests that those extramedullary hemopoiesis may be related to hemopoietic disorders in the bone marrow. These situations may simulate hepatosplenic myelosis in the man, an entity constructing the myeloproliferative syndrome [1]. This human syndrome, in many cases, is known to be associated with a marked increase in dyspoietic megakaryocytopoiesis, and the present cases with unexplained enhanced megakaryocytopoiesis in the spleen seem to be a mimetic syndrome in mice. The myeloproliferative disorders are currently considered not merely as a compensatory process but as an expression of stimulation on dormant mesenchymal stem cells, although the mechanism is not known. The infection in mice with the newly isolated MHV may provide a good experimental model for studying the pathogenesis of the myeloproliferative syndrome.

References

- [1] Damesheck, W. (1951). *Blood*, **6**, 372-375.
- [2] Fujiwara, K. (1971). In *Defining the Laboratory Animal*, Proceedings of the IVth International Symposium on Laboratory Animals in Washington, D.C., Schneider, H. A., editor, National Academy of Science, Washington, D.C., 77-92.
- [3] Hirano, N., Fujiwara, K., Hino, S., and Matsu-
moto, M. (1974). *Arch. Virusforsch.*, **44**, 298-
307.
- [4] Ishida, T., Taguchi, F., Lee, Y. S., Yamada, A.,
Tamura, T., and Fujiwara, K. (1978). *Lab.*
Anim. Sci., **28**, 269-276.
- [5] Scheid, M. P., Goldstein, G., and Boyse, E. A.
(1975). *Science*, **190**, 1211-1213.
- [6] Tamura, T., Machii, K., Ueda, K., and Fuji-
wara, K. (1978). *Microbiol. Immunol.*, **22**, 557-
564.
- [7] Tamura, T., Taguchi, F., Ueda, K., and Fuji-
wara, K. (1977). *Microbiol. Immunol.*, **21**, 683-
691.
- [8] Tamura, T., Ueda, K., Hirano, N., and Fuji-
wara, K. (1976). *Jap. J. exp. Med.*, **46**, 19-30.
- [9] Ward, J. M., Collins, M. J., Jr., and Parker,
J. C. (1977). *Lab. Anim. Sci.*, **27**, 372-376.

Explanation of Figures

- Fig. 1. Extensive extramedullary hemopoiesis in
the liver of a naturally infected nude mouse.
Hematoxylin and eosin (HE) stain. $\times 100$.
- Fig. 2. Perivascular hemopoiesis in the liver of a
naturally infected nude mouse. HE stain.
 $\times 200$.
- Fig. 3. A large number of myeloid cells along the
trabeculae in the spleen of a naturally infected
nude mouse. HE stain. $\times 200$.
- Fig. 4. Considerable granulopoiesis with dyspoietic
megakaryocytes in the red pulp of the spleen
in a naturally infected nude mouse. HE stain.
 $\times 400$.
- Fig. 5. Anaplastic femoral marrow with a de-
generated multinucleated giant cell in a natu-
rally infected nude mouse. HE stain. $\times 400$.