BRIEF NOTE

**Nodular Goiters of Three Mares and Their Foals,
Induced by Feeding Excessive Amount of Seaweed**

Kiyoshi MIYAZAWA, Shigekatsu MOTOYOSHI and Kazuya USUI

*Department of Medicine, School of Veterinary Medicine, Faculty of Agriculture,
University of Tokyo, Bunkyo-ku, Tokyo 113*

(Received for publication May 11, 1978)

Clinical cases of nodular goiters of 3 mares and their foals in a farm, probably induced by the administration of excessive iodine are reported. Mares were 7, 8 and 8 year-old and foals 7, 13 and 14 days after parturition at the time of consultation. Of these foals, 2 were male and 1 was female.

These mares were kept on ration consisting of 2 Kg of hay, 2.7 Kg of wheat bran, 2 Kg of oats, salt, calcium and over 60 g of feed additive which was mainly consisted of seaweed. The amount of iodine contained in this feed additive was 520–580 mg/Kg. As the results mares were fed over 30 mg (about 60 μ g/Kg body weight) of iodine daily throughout their gestation period. Daily requirement of the horse for iodine is 2–4 μ g/Kg body weight according to Maynard [6], but the maximum quantity of iodine which does not cause trouble is not clear in any report.

Clinical findings: Size of goiters was from the thumb tip to a hen egg in mares and from a hen egg to a fist in foals. By palpation, they were hard in consistency, but did not show pain. There were no abnormality in temperature, pulse and any other clinical findings.

Blood examination: RBC, Hb, Ht, WBC,

differentiation of white cells, total protein, serum protein electrophoretic pattern, cholesterol, ALP, GOT, GPT, LDH, Ca and P were determined with no abnormal deviation from the control mares and foals of the same age. In order to examine the function of thyroid gland, PBI, T_4 and T_3 in serum were assayed. Since there is rather a wide variation of these hormones in the blood in mares and foals in reference [1–5], no value among the mares and foals was beyond the normal range. The biopsy of thyroid was not carried out. With the above findings the termination of feeding of feed additives was immediately indicated.

After 2 months these goiters markedly reduced in sizes and at the same time they became soft in consistency. The similar blood examination was carried out with no abnormality again. From those findings of thyroid hormones, this goiter was not functional at all. Thus we diagnosed this as nodular goiters which was induced by feeding the excessive amount of iodine. The upper limit of iodine, which will not cause trouble, is certainly below 60 μ g/Kg body weight.

短報：海草給与によると思われる馬の甲状腺腫大例について：宮沢清志・本好茂一・臼井和哉（東京大学農学部家畜内科学教室）

Table 2. Blood examination in the foals

		Clinical case			Control		
		a	b	c			
RBC	$10^4/\text{mm}^3$	740	944	707			1003
Hb	g/dl	10.1	13.3	10.1			11.0
Ht	%	29.0	31.5	29.0			31.0
MCV	μ^3	39.2	33.4	41.0			30.9
MCH	$\mu\mu\text{g}$	13.6	14.1	14.3			11.0
MCHC	%	34.8	42.2	34.8			35.5
WBC	$10^2/\text{mm}^3$	59	91	79			94
Baso	%	1.0	0	0			
Eosino	%	0	0.5	0			
Band	%	9.0	7.5	9.0			
Seg	%	51.0	53.5	55.0			
Lympho	%	35.0	36.5	30.0			
Mono	%	4.0	2.0	6.0			
Protein T.	g/dl	6.0	6.0	6.3			6.4
Albumin	g/dl	3.32	3.18	3.29			3.48
α_1 -glob.	g/dl	0.10	0.16	0.07			0.13
α_2 -glob.	g/dl	0.47	0.76	0.59			0.56
β_1 -glob.	g/dl	0.71	0.72	0.79			0.71
β_2 -glob.	g/dl	0.38	0.38	0.28			0.55
γ -glob.	g/dl	1.03	0.80	1.28			0.97
A/G		1.24	1.12	1.10			1.19
Cholest. T.	mg/dl	198	212	242			248
F.	mg/dl	38	76	75			50
E.R.	%	80.7	64.0	69.3			79.8
ALP	KAu/dl	86.5	52.8	36.2			
GOT	Ku/ml	44.0	29.5	32.0			
GPT	Ku/ml	7.5	6.0	8.5			
LDH	Wu/ml	650	670	560			
Ca	mg/dl	10.4	7.0	6.2			
P	mg/dl	7.5	7.9	8.5			

Table 1. Blood examination in the mares

		Clinical case			Control	
		A	B	C	1	2
RBC	$10^4/\text{mm}^3$	738	1092	763	1010	999
Hb	g/dl	14.5	18.1	13.5	17.0	15.6
Ht	%	38.0	49.0	31.5	49.0	44.0
MCV	μ^3	51.5	44.9	41.3	48.5	44.0
MCH	$\mu\mu\text{g}$	19.6	16.6	17.7	16.8	15.6
MCHC	%	38.2	36.9	42.9	34.7	35.5
WBC	$10^2/\text{mm}^3$	105	100	75	81	44
Baso	%	1.0	2.0	1.0	1.0	3.0
Eosino	%	7.0	0	4.0	5.0	8.5
Band	%	2.0	9.0	1.0	1.0	0
Seg	%	36.0	41.0	57.5	64.0	40.5
Lympho	%	53.5	47.0	33.5	23.5	43.5
Mono	%	0.5	1.0	3.0	5.5	4.5
Protein T.	g/dl	6.0	6.8	6.4	6.4	6.0
Albumin	g/dl	3.59	3.48	3.33	3.17	3.36
α_1 -glob.	g/dl	0.07	0.12	0.08	0.20	0.20
α_2 -glob.	g/dl	0.50	0.62	0.70	0.60	0.61
β_1 -glob.	g/dl	0.89	1.26	0.85	0.45	0.62
β_2 -glob.	g/dl	0.46	0.61	0.40	1.00	0.65
γ -glob.	g/dl	0.48	0.72	1.03	0.99	0.55
A/G		1.49	1.05	1.08	0.93	1.27
Cholest. T.	mg/dl	90	108	90	85	112
F.	mg/dl	22	29	24	20	11
E.R.	%	75.8	73.1	73.8	76.5	90.2
ALP	KAu/dl	20.8	21.4	17.0	14.9	19.1
GOT	Ku/ml	57.5	51.0	44.0	49.0	48.5
GPT	Ku/ml	2.0	1.0	3.5	4.0	3.5
LDH	Wu/ml	470	470	370	315	385
Ca	mg/dl	5.5	13.2	11.3	12.5	8.7
P	mg/dl	2.8	4.5	2.9	3.7	3.9

Table 3. Assay of thyroid hormone in the mares

		Clinical case			Control	
		A	B	C	1	2
PBI	$\mu\text{g/dl}$	1.9	2.0	1.8	2.0	2.0
T ₄	$\mu\text{g/dl}$	1.2	3.0	0.4	1.8	2.0
T ₃	ng/dl	96	40	35	88	72
T ₄ +T ₃	$\mu\text{g/dl}$	1.3	3.0	0.4	1.9	2.1

Table 4. Assay of thyroid hormone in the foals

		Clinical case			Control
		a	b	c	1
PBI	$\mu\text{g/dl}$	9.9	5.4	5.2	13.4
T ₄	$\mu\text{g/dl}$	9.1	6.3	4.8	17.5
T ₃	ng/dl	424	296	308	463
T ₄ +T ₃	$\mu\text{g/dl}$	9.5	6.6	5.1	18.0

References

- [1] Baker, H. J., and Lindsey, J. R. (1968). *J. Amer. vet. med. Ass.*, **153**, 1618-1629.
- [2] Gribble, D. H. (1967). *In Equine Medicine and Surgery*, Amer. Vet. Pub. Inc., Wheaton, 442-457.
- [3] Irvine, C. H. G. (1967). *J. Endocrinol.*, **39**, 313-320.
- [4] Irvine, C. H. G. (1967). *Amer. J. vet. Res.*, **28**, 1687-1692.
- [5] Kallfelz, F. A., and Lowe, J. E. (1970). *J. Amer. vet. med. Ass.*, **156**, 1888-1891.
- [6] Maynard, L. H., and Loosli, J. K. (1956). *Animal Nutrition*, McGraw-Hill Book Comp. Inc., New York.

Explanation of Figure

Fig. 1. Nodular goiters of the foal.