Susceptibility of Fecal Anaerobic Bacteria from Pigs and Chickens to Five Polyether Antibiotics for Growth Promotion

Yoshimi BENNO¹⁾, Kimiko ENDO¹⁾, Nobue SHIRAGAMI¹⁾, and Tomotari MITSUOKA^{1,2)}

¹⁾The Institute of Physical and Chemical Research, Wako, Saitama 351-01, ²⁾Department of Biomedical Science, Faculty of Agriculture, the University of Tokyo, Bunkyo, Tokyo 113, Japan

(Received 4 November 1987/Accepted 16 March 1988)

ABSTRACT. The minimum inhibitory concentrations (MIC) of 100 strains of anaerobic bacteria, isolated from the intestines of healthy pigs and chickens, for five polyether antibiotics including cationomycin (CNM), lysocellin (LSC), monensin (MNS), portmicin (PRM) and salinomycin (SNM) were determined by an agar-dilution technique. CNM, MNS, PRM, and SNM, at a concentration of 6.25 µg/ml, were active against 41 to 56% of the *Bacteroides fragilis* group, whereas LSC showed less activity at the same concentration. All strains of genera *Fusobacterium*, *Selenomonas*, *Mitsuokella* and *Megasphaera* were resistant to the five polyether antibiotics. *Eubacterium* spp. were susceptible to the antimicrobial agents (MICs of 1.56 to 3.12 µg/ml) except for MNS. CNM and PRM inhibited all strains of *Bifidobacterium* spp. at lower concentrations, but LSC, MNS and SNM did not. CNM, PRM and SNM, at a concentration of 1.56 µg/ml, were active against all strains of *Lactobacillus* spp. whereas LSC and MNS did not inhibit them at the same concentration. *Clostridium perfringens* was also susceptible to five polyether antibiotics tested (MICs of 1.56 to 12.5 µg/ml). All strains of *B. fragilis* from pigs were resistant to the five polyether antibiotics but those from chickens were inhibited at lower concentrations.—**KEY WORDS:** chicken, intestinal bacterium, pig, polyether antibiotic, susceptibility.

Jpn. J. Vet. Sci. 50(3): 783–790, 1988

Antimicrobial agents have been widely used as food additives to improve growth and feed conversion in poultry, swine and cattle [31]. Although the mechanisms by which the feed additives promote growth of animals are unknown, the action of the agents may cause changes in the microbial populations within the intestinal tract [10, 31].

Polyether antibiotics are a diverse group of ionophores [27], most of which are produced by the genus *Streptomyces* [9, 13, 23]. Recently several new polyether antibiotics, produced by *Actinomycetes* other than *Streptomyces*, e.g. *Streptoverticillium* [18], *Nocardioopsis* [19], *Nocardia* [21], and *Actinomadura* [24], have been reported. It is known that polyether antimicrobial agents appear to reduce growth depressing toxins produced by intestinal bacteria, inhibit coccidial infection, and enhance the absorp-

tive efficiency and utilization of nutrients by animals [16, 25, 28]. However, *in vitro* susceptibility of intestinal anaerobes from animals to polyether antibiotics has not been investigated.

The purpose of the present study was to provide information on the susceptibility of anaerobic bacteria isolated from the intestines of healthy pigs and chickens to five polyether antibiotics used for growth promotion.

MATERIALS AND METHODS

Bacteria tested. A total of 100 strains of anaerobes from the feces of healthy pigs and chickens shown in Table 1 was used. These animals were not exposed to the polyether antibiotics throughout the study, while pigs were exposed to virginiamycin. The isolated strains consisted of the following species: 27

Table 1. Bacterial strains tested

Species	No. of isolates	
	Pigs	Chickens
<i>Bacteroides</i>		
<i>B. fragilis</i> group ^{a)}	10	17
<i>B. suis</i>	13	
<i>Bacteroides</i> spp.	2	3
<i>Fusobacterium</i>		
<i>F. necrophorum</i>	1	1
<i>Selenomonas</i>		
<i>Selenomonas</i> spp.	2	
<i>Mitsuokella</i>		
<i>M. multiacidus</i>	1	
<i>Eubacterium</i>		
<i>E. aerofaciens</i>	4	3
<i>Eubacterium</i> spp.	3	2
<i>Bifidobacterium</i>		
<i>B. pseudolongum</i>	1	1
<i>B. thermophilum</i>	4	2
<i>Propionibacterium</i>		
<i>P. acnes</i>	2	
<i>Clostridium</i>		
<i>C. perfringens</i>		3
<i>Peptostreptococcus</i>		
<i>P. anaerobius</i>	1	1
<i>P. asaccharolyticus</i>	4	
<i>P. magnus</i>	1	
<i>P. productus</i>	1	
<i>Peptostreptococcus</i> sp.	1	
<i>Megasphaera</i>		
<i>M. elsdenii</i>		1
<i>Lactobacillus</i>		
<i>L. acidophilus</i>	6	3
<i>L. fermentum</i>	1	1
<i>L. salivarius</i>	3	

a) Including *B. distasonis*, *B. fragilis*, *B. thetaiotaomicron*, and *B. vulgatus*.

strains of the *Bacteroides fragilis* group (13 of *B. fragilis*, 8 of *B. vulgatus*, 5 of *B. distasonis*, 1 of *B. thetaiotaomicron*), 13 of *B. suis*, 3 of the *B. buccae-oris* group, 2 of *B. pyogenes*, 2 of *Selenomonas* spp., 1 of *Fusobacterium necrophorum*, 2 of *F. varium*, 1 of *Mitsuokella multiacidus*, 7 of *Eubacterium aerofaciens*, 1 of *E. moniliforme*, 1 of *E. limosum*, 3 of *Eubacterium* spp., 6 of *Bifidobacterium thermophilum*, 2 of *B. pseudolongum*, 2 of *Propionibacterium acnes*, 3 of *Clostridium perfringens*, 4 of

Peptostreptococcus asaccharolyticus, 2 of *P. anaerobius*, 1 of *P. magnus*, 1 of *P. productus*, 1 of *Peptostreptococcus* sp., 1 of *Megasphaera elsdenii*, 9 of *Lactobacillus acidophilus*, 3 of *L. salivarius* and 2 of *L. fermentum*. Three species of *B. thermophilum* P2-91, *C. perfringens* ATCC 13124, and *L. acidophilus* ATCC 4356 were used as standard strains. The strains were maintained on Eggerth-Gagnon liver agar slants [22] filled with CO₂ gas and stored at 4°C. Transfers were usually made at 1-month intervals.

Antibiotics. The polyether antibiotics tested were cationomycin (CNM), lysocellin (LSC), monensin (MNS), portmicin (PRM), and salinomycin (SNM). They were supplied by Kaken Pharmaceutical Co., Ltd., Tokyo.

Antimicrobial susceptibility. The agar dilution method was used. Briefly, antimicrobial solutions were freshly prepared and added to *Brucella* agar (Difco, Detroit) supplemented with 5% defibrinated horse blood. The inoculum, a 24-hour culture in peptone-yeast extract-Fildes solution-glucose [14] broth to match McFarland standard No. 5, was dispensed using an automatic replicating system (Sakuma Co. Ltd., Tokyo) to deliver approximately 10⁸ colony-forming units. Control plates without drugs were inoculated before and after each set of drug-containing plates. Final concentrations of antibiotics tested ranged from 0.05 to 100 µg/ml. Plates were incubated in an anaerobic steel-wool jar [26] filled with 85% nitrogen gas, 10% carbon dioxide and 5% hydrogen gas at 37°C for 48 hrs. The lowest antimicrobial concentration that produced no visible bacterial growth was considered as the minimum inhibitory concentration (MIC).

RESULTS

The MIC values of five polyether antibio-

Table 2. Susceptibility of fecal anaerobic bacteria from pigs and chickens to five polyether antibiotics for growth promotion

Group and species	No. of strains tested	Cumulative percentage MIC at increasing antibiotic concentrations ^{a)}											
		≤0.05	0.1	0.2	0.39	0.78	1.56	3.12	6.25	12.5	25	50	≥100
Cationomycin													
<i>Bacteroides</i>	(45)		4.4	6.7		15.6		17.8	44.4	48.9	68.9	73.3	100
<i>B. fragilis</i> group	(27)					14.8		18.5	51.9	55.6	63.0	66.7	100
<i>B. suis</i>	(13)		15.4	23.1					30.8	38.5	84.6	92.3	100
Other <i>Bacteroides</i> spp.	(5)								20.0		60.0		100
<i>Fusobacterium</i> spp.	(3)												100
<i>Selenomonas</i> spp.	(2)												100
<i>Mitsuokella multiacidus</i>	(1)												100
<i>Eubacterium</i> spp.	(12)			23.1	92.3	100							
<i>Bifidobacterium</i> spp.	(8)			14.3	28.6	85.7	100						
<i>Propinoibacterium acnes</i>	(2)				50.0	100							
<i>Clostridium perfringens</i>	(3)				33.3	66.7			100				
<i>Peptostreptococcus</i> spp.	(9)		33.3		88.9	100							
<i>Megasphaera elsdenii</i>	(1)												100
<i>Lactobacillus</i> spp.	(14)			14.3	57.1	85.7	100						
Lysocellin													
<i>Bacteroides</i>	(45)		2.2	4.4	6.6				22.2	48.9	64.4	82.2	100
<i>B. fragilis</i> group	(27)								14.8	48.1	59.3	70.4	100
<i>B. suis</i>	(13)		7.7	15.4	23.1				46.2	69.2	76.9	100	
Other <i>Bacteroides</i> spp.	(5)										60.0	100	
<i>Fusobacterium</i> spp.	(3)												100
<i>Selenomonas</i> spp.	(2)												100
<i>Mitsuokella multiacidus</i>	(1)												100
<i>Eubacterium</i> spp.	(12)				7.9	30.8	61.5	100					
<i>Bifidobacterium</i> spp.	(8)						25.0	50.0			87.5		100
<i>Propinoibacterium acnes</i>	(2)					50.0	100						
<i>Clostridium perfringens</i>	(3)						33.3	66.7		100			
<i>Peptostreptococcus</i> spp.	(9)			33.3	55.6	100							
<i>Megasphaera elsdenii</i>	(1)												100
<i>Lactobacillus</i> spp.	(14)					21.4	42.9	92.9	100				
Monensin													
<i>Bacteroides</i>	(45)	2.2		6.7		8.9		26.7	40.0	55.6			100
<i>B. fragilis</i> group	(27)			3.7				22.2	40.7	55.6			100
<i>B. suis</i>	(13)	7.7		15.4		23.1		38.5	46.2	53.8			100
Other <i>Bacteroides</i> spp.	(5)							20.0		60.0			100
<i>Fusobacterium</i> spp.	(3)												100
<i>Selenomonas</i> spp.	(2)												100
<i>Mitsuokella multiacidus</i>	(1)												100
<i>Eubacterium</i> spp.	(12)							30.8	46.2	84.6	92.3		100
<i>Bifidobacterium</i> spp.	(8)							12.5		37.5			100
<i>Propinoibacterium acnes</i>	(2)							100			100		
<i>Clostridium perfringens</i>	(3)							33.3		66.7			
<i>Peptostreptococcus</i> spp.	(9)		33.3			55.6	100						
<i>Megasphaera elsdenii</i>	(1)												100
<i>Lactobacillus</i> spp.	(14)							14.3	28.6	100			

Table 2. Continued.

Group of species	No. of strains tested	Cumulative percentage MIC at increasing antibiotic concentrations ^{a)}											
		≤0.05	0.1	0.2	0.39	0.78	1.56	3.12	6.25	12.5	25	50	≥100
Portmicin													
<i>Bacteroides</i>	(45)	2.2		6.6	11.1		13.3	28.9	35.6	46.7	75.6	80.0	100
<i>B. fragilis</i> group	(27)	3.7		7.4			11.1	37.0	40.7	51.9	66.7	70.4	100
<i>B. suis</i>	(13)			7.7	23.1				30.8	38.5	92.3		100
Other <i>Bacteroides</i> spp.	(5)								20.0	60.0		80.0	100
<i>Fusobacterium</i> spp.	(3)												100
<i>Selenomonas</i> spp.	(2)												100
<i>Mitsuokella multiacidus</i>	(1)												100
<i>Eubacterium</i> spp.	(12)				30.8	61.5	100						
<i>Bifidobacterium</i> spp.	(8)			25.0			87.5		100				
<i>Propinoibacterium acnes</i>	(2)					100							
<i>Clostridium perfringens</i>	(3)						33.3		100				
<i>Peptostreptococcus</i> spp.	(9)		33.3		88.9	100							
<i>Megasphaera elsdenii</i>	(1)												100
<i>Lactobacillus</i> spp.	(14)			7.1	21.4	42.9	100						
Salinomycin													
<i>Bacteroides</i>	(45)	2.2	4.4	8.9		11.1	17.8	37.8	62.2	80.0	82.2	86.7	100
<i>B. fragilis</i> group	(27)					7.4	18.5	29.6	55.6		77.8	88.9	100
<i>B. suis</i>	(13)		7.7	23.1				46.2	76.9	92.3	100		
Other <i>Bacteroides</i> spp.	(5)							60.0		100			
<i>Fusobacterium</i> spp.	(3)												100
<i>Selenomonas</i> spp.	(2)												100
<i>Mitsuokella multiacidus</i>	(1)												100
<i>Eubacterium</i> spp.	(12)			7.7	61.5	92.3				100			
<i>Bifidobacterium</i> spp.	(8)				25.0	37.5	50.0			87.5			100
<i>Propinoibacterium acnes</i>	(2)				100								
<i>Clostridium perfringens</i>	(3)				33.3	66.7				100			
<i>Peptostreptococcus</i> spp.	(9)		33.3	55.6	88.9		100						
<i>Megasphaera elsdenii</i>	(1)												100
<i>Lactobacillus</i> spp.	(14)		7.1	42.9	92.9	100							

a) µg/ml.

tics, including CNM, LSC, MNS, PRM, and SNM, for a total of 100 strains of anaerobic bacteria isolated from the intestine of healthy pigs and chickens are shown in Table 2. Within *Bacteroides* spp., 41, 41, 52, and 56% of the *B. fragilis* group tested were inhibited by MNS, PRM, CNM, and SNM, respectively, at MIC of 6.25 µg/ml, whereas LSC showed less activity on these microorganisms at the same concentration. *B. suis* from pig feces had an MIC pattern with two peaks except for MNS, 23% at MIC of 0.2 to 0.4 µg/ml and 77% at MIC of 6.25 to 100

µg/ml. All strains belonging to genera *Fusobacterium*, *Selenomonas*, *Mitsuokella*, and *Megasphaera* were resistant to the five polyether antibiotics tested. With the exception of NMS, all *Eubacterium* spp., which are a beneficial in animal production, were inhibited by CNM and PRM at a concentration ranging from 1.56 to 6.25 µg/ml. LSC, MNS, and SNM, however, were less active against the microorganisms at the same concentration. *Propionibacterium acnes* was inhibited by all antibiotics at 0.78 to 3.12 µg/ml. *C. perfringens* which produces

Table 3. Susceptibility of fecal *Bacteroides fragilis* from pigs and chickens to five polyether antibiotics for growth promotion

Antibiotic	Source	MIC ($\mu\text{g/ml}$) ^{a)}			
		Range	MIC ₅₀	MIC ₇₅	MIC ₉₀
Cationomycin	Pigs	25 – ≥ 100	≥ 100	≥ 100	≥ 100
	Chickens ^{c)}	0.78 – 12.5	6.25	6.25	12.5
Lysocellin	Pigs	12.5 – ≥ 100	≥ 100	≥ 100	≥ 100
	Chickens	6.25 – 50	12.5	12.5	50
Monensin	Pigs	12.5 – ≥ 100	≥ 100	≥ 100	≥ 100
	Chickens	3.12 – 12.5	0.25	12.5	12.5
Portmicin	Pigs	12.5 – ≥ 100	≥ 100	≥ 100	≥ 100
	Chickens	0.2 – 12.5	3.12	12.5	12.5
Salinomycin	Pigs	6.25 – ≥ 100	50	≥ 100	≥ 100
	Chickens	1.56 – 12.5	6.25	12.5	12.5

a) Concentration of anti-microbial agents at showing growth inhibition of over 50%, 75%, or 90%.

b) 8 strains was tested.

c) 5 strains was tested.

growth depressing toxins was inhibited by the polyether antibiotics tested at a concentration ranging from 1.56 to 12.5 $\mu\text{g/ml}$. All strains of *Peptostreptococcus* spp. were susceptible to the polyether antimicrobial agents, with the exception of CNM, at MIC of about 1.56 $\mu\text{g/ml}$. *Lactobacillus* spp. which predominate in the intestinal microflora of healthy pigs and chickens were inhibited by CNM, PRM, and SNM at MIC of 1.56 $\mu\text{g/ml}$, whereas these microorganisms revealed susceptibility to LSC and NMS at a concentration ranging from 6.25 to 12.5 $\mu\text{g/ml}$.

Table 3 shows the susceptibility of *B. fragilis* isolated from the intestines of healthy pigs and chickens to five polyether antibiotics used for growth promotion. All strains of *B. fragilis* from pigs were resistant to the polyether antibiotics but those from chickens were inhibited by these antibiotics at a concentration ranging from 0.2 to 12.5 $\mu\text{g/ml}$.

DISCUSSION

It is widely known that polyether antibio-

tics exhibit both antimicrobial and anticoccidial activities [5, 6, 15]. Polyether antimicrobial agents, including NMS and SNM, are added to feeds as growth promoters in Japan, but CNM, LSC and PRM are not used in this way. In the near future, however, three polyether antibiotics may be permitted for use as a food additives.

The criteria of susceptibility as used here with the growth promoting agents are tentative because criteria for the evaluation of susceptibility and resistance have never been established. Dutta and Devriese [7, 8] reported that *C. perfringens* strains isolated from the feces of chickens, cows and pigs were inhibited by NMS at MIC of 4 $\mu\text{g/ml}$ and this MIC was established as the susceptible value. *In vitro* sensitivity levels of the polyether antibiotics against the intestinal isolates with growth-depressing activity may be interpreted as indicating sensitivity to the growth promotor. It is of interest to determine what effects, if any, polyether antibiotics exert on the intestinal microflora. Further studies are required to determine the effects of polyether antibiotics on the composition of intestinal microflora in the

animals.

The existence of growth-depressing activity which can be influenced by antimicrobial agents used for growth promotion has been noted mainly among the gram-positive intestinal flora, e.g. *Clostridium perfringens* [20], *Streptococcus faecalis* [11, 12], *S. faecium* and *S. faecalis* subspecies *liquefaciens* [1]. In the present study, five polyether antibiotics were active *in vitro* against most of the species belonging to the genera *Eubacterium*, *Bifidobacterium*, *Propionibacterium*, *Clostridium*, *Peptostreptococcus*, and *Lactobacillus*. However, they did not act uniformly and their activity spectra differed markedly.

Decuypere *et al.* [4] and Vervaeke *et al.* [30] found that virginiamycin and spiramycin suppress growth of lactobacilli and streptococci in the small intestines of pigs, thereby influencing carbohydrate metabolism in a way apparently beneficial to the host. In this study, *Lactobacillus* spp. were inhibited by polyether antimicrobial agents with the exception of MNS at MIC ranging from 0.2 to 6.25 $\mu\text{g/ml}$. Thus, effects of polyether antibiotics on lactobacilli were expected.

It is widely known that pathogenicity of *Eimeria tenella* is promoted by the presence of *C. perfringens* in the intestine of chickens [3, 17, 29]. Our previous study [2] demonstrated that SNM and MNS inhibited *C. perfringens* isolated from broiler intestines at lower concentrations. One of three strains of *C. perfringens* tested in this study was resistant to SNM and MNS at higher concentrations. The existence of polyether antibiotic-resistant *C. perfringens* strains may soon be increased by the use of polyether antimicrobial agents.

On the other hand, gram-negative anaerobic bacteria were less affected by five polyether antibiotics. Although the strain tested was a few, the genera *Fusobacterium*, *Mitsuokella*, *Selenomonas*, and *Megas-*

phaera were not inhibited at concentrations of more than 100 $\mu\text{g/ml}$. Some isolates of the *B. fragilis* group, however, did not show a uniform susceptibility and revealed different activity spectra. CNM and SNM exhibited less activity on the *B. fragilis* group with an MIC₅₀ of 6.25 $\mu\text{g/ml}$, MIC₇₅ ranging from 12.5 to 25.0 $\mu\text{g/ml}$, and MIC₉₀ of more than 100 $\mu\text{g/ml}$. In addition, the concentrations at which the *B. fragilis* group were resistant to LSC, MNS, and PRM were markedly higher than those for CNM and SNM. Although the polyether antibiotics are characterized by lower activity against anaerobic gram-negative bacilli as described previously [9, 23], the *B. fragilis* group, which predominates in the intestines of animals, had different activity spectra.

In the present study the susceptibilities to five polyether antibiotics were noted to vary with the animal source of *B. fragilis*. All strains of *B. fragilis* isolated from chickens, but not all of the *B. fragilis* strains isolated from pigs were sensitive to the five antimicrobial agents. The pigs and chickens, from which *B. fragilis* was isolated, had not been exposed to the polyether antibiotics. Although the pigs were exposed to virginiamycin, the reason why the isolates revealed different susceptibilities is unknown. Further studies are required to confirm which anaerobic bacteria have acquired or natural resistance to polyether antibiotics.

REFERENCES

1. Barnes, E. M., Maed, G. C., Impey, C. S., and Admas, B. W. 1978. The effect of dietary bacitracin on the incidence of *Streptococcus faecalis* subspecies *liquefaciens* and related streptococci in the intestines of young chicks. *Br. Poult. Sci.* 19: 713-723.
2. Benno, Y., Endo, K., and Mitsuoka, T. 1988. Isolations of fecal *Clostridium perfringens* from broiler chickens and their susceptibility to eight antimicrobial agents for growth promotion. *Jpn. J. Vet. Sci.* 50: 832-834.
3. Bradley, R. E., and Radhakrishnan, C. V. 1973.

- Coccidiosis in chickens: obligate relationship between *Eimeria tenella* and certain species of cecal microflora in the pathogenesis of the disease. *Avian Dis.* 17: 461-476.
4. Decuyper, J., Henderickx, H. K., and Vervaeke, I. 1973. Influence of nutritional doses of virginiamycin and spiramycin on the quantitative and topographical composition of the intestinal flora of artificially reared piglets. *Zentralbl. Bakterirol. Parasitenkd. Infektionskr. Hyg., Abt. I. Orig.* A223: 348-355.
 5. Danforth, H. D., Ruff, M. D., Reid, W. M., and Miller, R. L. 1977. Anticoccidial activity of salinomycin in battery raised broiler chickens. *Poult. Sci.* 56: 926-932.
 6. Danforth, H. D., Ruff, M. D., Reid, W. M., and Johnson, J. 1977. Anticoccidial activity of salinomycin in floorpen experiments with broilers. *Poult. Sci.* 56: 933-938.
 7. Dutta, G. N., and Devriese, L. A. 1980. Susceptibility of *Clostridium perfringens* of animal origin to fifteen antimicrobial agents. *J. Vet. Pharmacol. Ther.* 3: 227-236.
 8. Dutta, G. N., and Devriese, L. A. 1981. Macrolide-lincosamide-streptogramin resistance patterns in *Clostridium perfringens* of animal origin. *Antimicrob. Agents Chemother.* 19: 274-278.
 9. Ebata, E., Kasahara, H., Sekine, K., Inoue, Y. 1975. Lysocellin, a new polyether antibiotic. I. Isolation, purification, physicochemical and biological properties. *J. Antibiotics* 28: 118-121.
 10. Eyssen, H., and De Somer, P. 1963. Effect of antibiotics on growth and nutrient absorption of chicks. *Poult. Sci.* 42: 1373-1379.
 11. Eyssen, H., and De Somer, P. 1967. Effect of *Streptococcus faecalis* and a filtrable agent on growth and nutrient absorption in gnotobiotic chicks. *Poult. Sci.* 46: 323-313.
 12. Fuller, R., Coates, M. E., and Harrison, G. F. 1979. The influence of specific bacteria and a filtrable agent on the growth of gnotobiotic chicks. *J. Appl. Bacteriol.* 46: 335-342.
 13. Haney, M. E. Jr., and Hoehn, M. M. 1968. Monensin, a new biologically active compound. I. Discovery and isolation. *Antimicrob. Agents Chemother.* 7: 349-352.
 14. Holdeman, L. V., Cato, E. P., and Moore, W. E. C. 1977. *Anaerobic Laboratory Manual*, 4th ed., Anaerobe Laboratory, Virginia Polytechnic Institute and State University, Blacksburg, Va.
 15. Itagaki, K., Tsubokura, M., and Otsuki, K. 1974. Studies on methods for evaluation of anticoccidial drugs in vitro. *Jpn. J. Vet. Sci.* 36: 195-202.
 16. Ito, F., Takano, S., Okazaki, M., Iizumi, A., Murakami, T., Seima, T., Noda, K., Tomita, S., Takase, M., Masuda, K., and Kudo, H. 1984. Effects of feed additive salinomycin on growth of pigs. *Jpn. J. Zootech. Sci.* 55: 619-627 (in Japanese with English summary).
 17. Johansson, K. R., and Sarles, W. B. 1948. Bacterial population changes in the ceca of young chickens infected with *Eimeria tenella*. *J. Bacteriol.* 56: 635-647.
 18. Kitame, F., Utsushikawa, K., Kohama, T., Saito, T., Kikuchi, M., and Ishida, N. 1974. Laidlomycin, a new antimycoplasmal polyether antibiotic. *J. Antibiotics* 27: 884-888.
 19. Kusakabe, Y., Takahashi, N., Iwagaya, Y., and Seino, A. 1987. Portmicin, a new antibiotic. *J. Antibiotics* 40: 237-238.
 20. Lev, M., and Forbes, M. 1959. Growth response to dietary penicillin of germ free chicks and of chicks with a defined intestinal flora. *Br. J. Nutr.* 13: 78-84.
 21. Liu, C. -M., Hermann, T. E., Downey, A., Prosser, B. La T., Schildknecht, E., Palleroni, N. J., Westley, J. W., and Miller, P. A. 1983. Novel polyether antibiotics X-14868A, B, C, and D produced by a *Nocardia*. Discovery, fermentation, biological as well as ionophore properties and taxonomy of the producing culture. *J. Antibiotics* 36: 343-350.
 22. Mitsuoka, T., Segi, T., and Yamamoto, S. 1965. Eine verbesserte Methodik der qualitativen und quantitativen Analyse der Darmflora von Menschen und Tieren. *Zentralbl. Bakterirol. Parasitenkd. Infektionskr. Hyg. I. Abt. Orig.* A195: 455-469.
 23. Miyazaki, Y., Shibuya, M., Sugawara, H., Kawaguchi, O., Hirose, C., and Nagatsu, J. 1974. Salinomycin, a new polyether antibiotic. *J. Antibiotics* 27: 814-821.
 24. Nakamura, G., Kobayashi, K., Sakurai, T., and Isono, K. 1981. Cationomycin, a new polyether ionophore antibiotic produced by *Actinomadura* sp. nov. *J. Antibiotics* 34: 1513-1514.
 25. Nakashima, T., Masuno, T., Sakauchi, R., and Hoshino, S. 1982. Effect of salinomycin on feed efficiency, ruminal and blood characteristics of steers. *Jpn. J. Zootech. Sci.* 53: 541-546 (in Japanese with English summary).
 26. Parker, C. A. 1955. Anaerobiosis with iron wool. *Aust. J. Exp. Biol. Med. Sci.* 33: 33-37.
 27. Pressman, B. C., Harris, E. J., Jogger, W. S., and Johnson, J. H. 1967. Antibiotic-mediated transport of alkaliions across lipid barriers. *Proc. Natl. Acad. Sci. U.S.A.* 70: 1949-1956.
 28. Shumard, R. F., and Callender, M. E. 1968. Monensin, a new biologically active compound. VI. Anticoccidial activity. *Antimicrob. Agents Chemother.* 7: 369-377.
 29. Stutz, M. W., Johnson, S. L., Judith, F. R., and Muir, L. A. 1983. Effect of the antibiotic thiopeptin on *Clostridium perfringens* and growth and

- feed efficiency of broiler chicks. *Poult. Sci.* 62: 1633-1638.
30. Vervaeke, I., Decuyper, J. A., Dierick, N. A., and Henderickx, H. K. 1979. Quantitative *in vitro* evaluation of the energy metabolism influenced by virginiamycin and spiramycin used as a growth promoters in pig nutrition. *J. Anim. Sci.* 49: 846-856.
31. Visek, W. J. 1978. The mode of growth promotion by antibiotics. *J. Anim. Sci.* 46: 1447-1469.

要 約

豚および鶏糞便由来嫌気性菌の発育促進用ポリエーテル系抗生物質に対する感受性について：辨野義己¹⁾・遠藤希三子¹⁾・白神伸江¹⁾・光岡知足^{1,2)} (¹⁾理化学研究所・²⁾東京大学農学部実験動物学教室)——発育促進用ポリエーテル系抗生物質、カチオノマイシン (CNM)、ライソセリン (LSC)、モネンシン (MNS)、ポートミシン (PRM) およびサリノマイシン (SNM) に対する健康豚および鶏糞便由来嫌気性菌100菌株の最小発育阻止濃度 (MIC) を寒天希釈法によって測定したところ、*Bacteroides fragilis* グループに属する菌株の41~56%はCNM, MNS, PRM および SNM の6.25 μ g/mlの濃度で抑制されたが、LSCには同一濃度で抑制されなかった。*Fusobacterium*, *Selenomonas*, *Mitsuokella* および *Megasphaera* の全供試菌株はポリエーテル系抗生物質に耐性を示した。*Eubacterium* はMNSを除く抗生物質に1.56~3.12 μ g/mlの濃度で抑制された。*Bifidobacterium* の全菌株はCNM および PRM に対して低濃度で抑制されたが、LSC, MNS および SNM には弱い耐性を示した。また、*Clostridium perfringens* は全供試薬剤に1.56~12.5 μ g/mlで抑制され、*Lactobacillus* はCNM, PRM および SNM の1.56 μ g/mlで抑制されたが、LSC および MNS に対して12.5 μ g/mlで抑制された。豚糞便由来 *B. fragilis* はポリエーテル系抗生物質に対して耐性を示したが、鶏糞便由来 *B. fragilis* は低濃度で抑制された。