<外人講演>

9 & 10. Liver function test in hepatic amoebiasis, based on
274 clinical casesVikit Viranuvatti*, Tranakchit Harinasuta*, Ukrist Plengvanich*, Pisiu Choungchareon* and
Vikul Viranuvatti†

Bangkok, Thailand.

Hepatic amoebiasis is a common disease in Thailand. Intra-thoracic involvement caused by rupture of the abscess producing various pathological conditions were reported from time to time (Kangsadal & Tandhan and 1952, Harinasuta 1950, Prijanonda & Yongchaigudh 1960). Other rare complications such as amoebic pyopericardium, (Buri et al, 1955), chronic amoebic peritonitis (Pansuwasn, 1957) and vesical amoebiasis (Harinasuta, 1955) were also encountered. In the other parts of the world numerous works were published. Despite extensive work done in this subject few words were mentioned about the liver function tests in hepatic amoebiasis. In this country Harinasuta first studied hippuric acid synthesis by using oral sodium bonzoate and found that impairment of hippuric acid was found in 95% of cases. Later Viranuvatti (1954, 1959) found that Iodine test was abnormal in 94.6% of amoebic liver abscess and was more sensitive than cephalin cholesterol flocculation, thymol flocculation and thymol turbidity test. This test is of diagnostic value in differentiation of various hepatic diseases.

Purpose of study It is the purpose of the authors to study the liver function of hepatic amoebiasis in a large series in order to evaluate the sensitivity, significance and value of each test in the diagnosis of this morbid condition.

Material and method According to Kean (1955), amoebic hepatitis clinically comprises :
1. an enlarged tender liver associated with fever, 2. therapeutic response to anti-amoebic therapy,
3. a moderate leucocytosis, 4. a history of intestinal amoebiasis.

Cases in this series were collected from 1951 to 1959 comprising only those admitted in the medical department, Siriraj Hospital. Bangkok, Thailand. Cases designated as amoebic liver abscess are those cases which yielded characteristic and sterile pus on aspiration whereas amoebic hepatitis included cases with enlarged and tender liver, fever, moderate leucocytosis with predominant polymorphonuclear cells and therapeutic response to anti-amoebic therapy alone. The history of intestinal amoebiasis could not be obtained in all cases as a large number of cases of colonic amoebiasis do not necessarily produce dysenteric symptoms. There were 200 cases of amoebic liver abscess and 74 cases of amoebic hepatitis subjected to this study. During the early period of study only few liver function tests could be performed, in the latter years serum enzymes study, paper electrophoresis and other seroflocculation tests were done.

Result There were 240 males and 34 females in this series resulting in the ratio of 8 to 1. The commonest age group afflicted with this illness was from third to fifth decade. The highest age in the series was 71 and the lowest 17 (see table 1).

* From the division of gastroenterology and tropical medicine Medical department, Siriraj hospital, University of Medical Sciences, Bangkok, Thailand.

† Dean, School of Medical technology, University of Medical Sciences, Bangkok, Thailand.

Table 1.

Age in years	Amoebic liver abscess			Amoebic hepatitis			Total No. of cases.
	Total	M.	F.	Total	M.	F.	
10-19	2	1	1	2	1	1	4
20-29	21	17	4	8	7	1	29
30-39	46	41	5	24	19	5	70
40-49	72	65	7	17	12	5	89
50-59	37	35	2	18	17	1	55
60-69	21	20	1	4	3	1	25
70- and over	1	1	—	1	1	—	2
Total	200	180	20	74	60	14	274

The result of liver function studies was divided into two groups i. e. amoebic liver abscess and amoebic hepatitis according to the criteria described previously.

Table 2. Value of serum albumin and globulin in 192 cases of amoebic liver abscess and 40 cases of amoebic hepatitis

Serum protein in gm. %	Amoebic liver abscess No. of cases		Amoebic hepatitis No. of cases	
	Albumin	Globulin	Albumin	Globulin
1-1.9	4	27	1	11
2-2.9	68	70	8	14
3-3.9	94	69	20	12
4-4.9	21	21	8	3
5-5.9	5	5	2	—
6-upwards	—	—	1	—

Serum albumin and globulin determination was done in 192 cases of amoebic liver abscess and 40 cases of amoebic hepatitis totalling 232 cases. In amoebic liver abscess the average albumin level was 3.1 and globulin 2.8 with average total serum protein of 5.9 gm.%. The maximum value of albumin was 5.2 and the minimum value 1.5 whereas the maximum value of globulin was 5.4 and minimum 1.0 gm. %. The value of albumin globulin ratio was less than 1 in 76 cases of amoebic liver abscess (39.6%) and 8 cases of amoebic hepatitis (20.0%). In amoebic hepatitis, determination of serum protein was done in 40 cases. The average serum albumin level was 3.3 and globulin 2.1 with average serum protein of 5.4 gm. %.

Soroflocculation tests Iodine tests. Strong Lugol's solution or Iodine tests was performed in 170 cases of hepatic amoebiasis by the method previously described (Viramvatti, 1959). There were 126 cases of amoebic liver abscess and 44 cases of hepatitis.

From the result obtained, 83.7% of amoebic liver abscess showed abnormal Iodine tests, the percentage dropped to 79.6% in amoebic hepatitis. This result differed slightly from the series in our previous publication (Viramvatti, 1959) which showed 94.5% positivity in liver abscess and only 33.3% in hepatitis. However, this series is larger particularly in amoebic hepatitis group. On account of difficulty in clinical differentiation between abscess and hepatitis, many cases in this series might as well be an easy cases of small abscess, thus the abnormal result

Table 3. Result of Iodino test in 170 cases of hepatic amoebiasis

Result	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
Negative	16	9	25
positive	110	35	145
4 plus	42	6	48
3 plus	20	7	27
2 plus	26	13	39
1 plus	22	9	31
Total	126	44	170

was almost as high as in the liver abscess group. It is our belief that the result in this large series is more reliable than the previous report.

Cephalin cholesterol flocculation test Cephalin cholesterol flocculation test was done in 109 cases of hepatic amoebiasis by the method of Hanger. (1936). There were 81 cases of amoebic liver abscess and 28 cases of amoebic hepatitis.

Table 4. Result of cephalin cholesterol flocculation test in 109 cases of hepatic amoebiasis

Result	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
Negative	20	8	28
1 plus	26	5	31
2 plus	20	9	29
3 plus	9	4	13
4 plus	6	2	8
Total	81	28	109

In amoebic liver abscess the result was abnormal (more than 1 plus) in 35 out of 81 cases (43.2%) and in amoebic hepatitis the test was abnormal in 15 out of 28 cases (53.6%). In both groups the cephalin cholesterol flocculation test was abnormal in 50 out of 109 (46.0%). The two groups could not be compared statistically as the series of hepatitis was too small.

Thymol turbidity test Thymol turbidity test was estimated by Shank and Hoagland's modification of Maclagan's technique (1946) in 158 cases of hepatic amoebiasis. There were 122 cases of amoebic liver abscess and 36 cases of amoebic hepatitis.

Table 5. Result of thymol turbidity test in 158 cases of hepatic amoebiasis

Value in units	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
0- 4.9	74	23	97
5- 9.9	33	9	42
10-14.9	11	4	15
15-19.9	3	—	3
20-upward	1	—	1
Total	122	36	158

Thymol turbidity test was found to be abnormal (5 units upwards) in 48 out of 122 cases of amoebic liver abscess (39.3%) and in 13 out of 36 cases of amoebic hepatitis (36.1%). It was abnormal in 61 out of 158 cases of hepatic amoebiasis (38.8%). The maximum value of thymol turbidity test in this series was 20.1 units. This was a cases of amoebic liver abscess of 9×9 cm. in diameter yielding 260 c. c. of pus on the first aspiration. The average mean value of thymol turbidity in amoebic liver abscess group was 10.05 and 0.5 units in hepatitis group.

Zinc sulfate turbidity test Zinc sulfate turbidity test was done by the method of Kunkel and his associates (1954) in 80 cases of hepatic amoebiasis. There were 57 cases of amoebic liver abscess and 23 cases of amoebic hepatitis.

Table 6. Result of zinc sulfate turbidity test in 80 cases of hepatic amoebiasis

Value in units	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
0-11.9	18	8	28
12-14.9	13	7	20
15-19.9	10	5	15
20-24.9	6	2	8
25-29.9	4	1	5
30-34.9	4	—	4
35-upwards	2	—	2
Total	57	23	82

The abnormal value of more than 12.0 units was found in 39 out of 57 cases of amoebic liver abscess (68.5%), and in 15 out of 23 cases of amoebic hepatitis (65.2%). In both groups the test was abnormal in 54 out of 80 cases (57.5%). The maximum value of zinc sulfate turbidity was 55.0 units in amoebic liver abscess group and 25.0 units in hepatitis group. The mean value was 28.3 units in liver abscess and 13.5 in hepatitis group.

Copper acetate test Copper acetate test was determined by the method of Zellek and Frade in 21 cases of hepatic amoebiasis. The value of more than 1 plus was considered abnormal. All cases subjected to this study were liver abscess. Negative result was obtained in 2 cases, 1 plus in 4, plus in, 5, 3 plus in 7, and 4 plus in 3. Abnormal result of more than 1 plus was found in 155 out of 21 cases (71.4%).

It is generally accepted that seroflocculation tests are based on the alteration in various serum protein fractions. In amoebic liver abscess hepatic pathology shown extensive liver cell necrosis produced by proteolytic enzyme of *Entamoeba histolytica* results in deranged liver function reflected by abnormal seroflocculation tests and altered albumin globulin level. It is the purpose of the authors to study the correlation between the size of liver abscess and the result of serum protein level and various seroflocculation tests. The size of liver abscess cavity was measured by intracavitary instillation of radio-opaque Iodine compound (Pyelosil 70%, Dionosil 70% and Lipiodol) and P-A film of the liver region was taken immediately after instillation. All cases subjected in this study showed only one cavity. The width and length of the cavity was measured on the X-ray films and recorded.

From this table it is well seen that all seroflocculation tests bear no relationship with the size of liver abscess. The maximum size of cavity in this series was case No. 1 which 2 plus Iodine test, negative cephalin cholesterol flocculation test, 3.0 units of thymol turbidity and zinc

Table 7. Relationship between the value of serum protein, seroflocculation tests and size of amoebic liver abscess as determined by measurement of cavity by intra-cavitary instillation of radio-opaque media

No.	Size by width and length in cm.	A/G	Iodine	C. C. F	T. F	T. T.	Zinc So ₄
1	21 × 11.7	2.9/3.0	2 +	Neg.	—	3.0	14.0
2	15 × 15	4.1/1.7	1 +	1	—	1.0	4.0
3	14 × 13.5	2.9/4.5	4 +	—	—	—	—
4	16 × 12	2.1/3.0	3 +	—	—	—	—
5	13 × 13	2.8/4.4	4 +	2	—	8.0	5.0
6	14.5 × 12.8	3.5/2.8	3 +	4	3	—	—
7	13 × 12.4	3.3/3.8	2 +	—	—	—	—
8	14.5 × 10.8	—	4 +	2	—	8.0	5.0
9	12 × 12.0	—	4 +	—	—	—	—
10	14 × 11	—	1 +	—	—	—	—
11	12 × 12.8	3.1/3.3	2 +	3	—	—	10.25
12	12.3 × 8.4	2.1/3.8	3 +	1	—	6.0	18.0
13	10 × 10	3.4/2.6	—	1	—	3.5	14.5
14	13 × 5	2.2/3.1	4 +	2	—	0	5.0
15	9 × 8	3.1/3.7	2 +	1	—	4.0	20.0
16	7.5 × 7.5	—	—	neg	—	3.0	16.0
17	9.3 × 5.0	2.3/2.4	1 +	1	—	6.0	13.0
18	6 × 7	3.0/4.8	4 +	4	—	7.0	21.0
19	10.5 × 5	3.1/3.1	2 +	1	—	5.0	25.0
20	9.8 × 8.4	2.9/3.7	4 +	2	—	6.0	20.0
21	12 × 9.4	4.1/4.5	4 +	1	—	3.0	14.0
22	8 × 6	—	3 +	2	—	7.0	21.0
23	4.5 × 3.6	4.8/3.7	1 +	3	—	2.0	6.0
24	11.5 × 7.8	3.0/3.3	2 +	—	—	—	—
25	8.7 × 8.7	3.2/1.8	4 +	4	—	4.12	—
26	7.8 × 11.5	3.0/3.3	2 +	—	—	—	—
27	10 × 12	2.9/3.8	1 +	—	—	2.45	—
28	5.5 × 6.4	—	4 +	—	—	—	—
29	6.5 × 5	—	4 +	—	—	8.5	—
30	5 × 6	3.1/1.7	—	1	1	—	—
31	8.5 × 8.5	—	—	2	1	—	—
32	13 × 10	2.3/3.5	—	2	1	—	—
33	14 × 8	—	—	neg	—	6.0	—
34	9.5 × 6.6	4.3/3.5	—	2	1	—	—
35	6 × 4.5	2.8/2.7	—	2	neg	—	—
36	14 × 9	5 /2.6	2 +	—	—	—	—
37	8.7 × 8.1	3.9/2.3	1 +	2	—	5.0	22.0
38	11 × 7	3.1/2.3	4 +	neg	—	8.0	55.0
39	5.5 × 5	5.0/2.8	1 +	2	—	4.0	4.0
40	13.5 × 5.5	3.6/4.6	1 +	—	—	—	21.2
41	12 × 8.1	3.0/3.2	2 +	—	—	—	—
42	11 × 10.5	—	neg	—	—	1.7	9.7
43	11 × 11	3.9/1.3	neg	—	—	—	5.9
44	8.8 × 5	2.9/2.6	3 +	2	3	—	—
45	9.0 × 9	3.2/3.4	4 +	neg	neg	—	—
46	9.5 × 8	2.4/1.1	4 +	—	—	—	—
47	8 × 8	3.2/1.2	4 +	—	—	—	—
48	9.7 × 6	3.1/1.1	—	3	3	—	—
49	14 × 8.5	1.6/5.2	4 +	—	—	5.8	—
50	6.1 × 6.2	3.3/2.0	2 +	—	—	—	—
51	6 × 6.5	4.5/3.3	1 +	—	—	2.65	—
52	9 × 9	3.5/1.5	1 +	—	—	20.1	—

sulfate turbidity of 14.0 units; this case had 21×11.7 cm. size cavity. The smallest cavity in this series was case No. 23 with 4.5×3.6 cm. cavity and showed 1 plus Iodine test, 3 plus cephalin cholesterol flocculation test, 2.0 units thymol turbidity and Zinc sulfate turbidity of 6.0 units. Case No. 38 showed highest value of zinc sulfate turbidity i.e. 55 units, 4 plus Indine test, 8.0 units thymol turbidity but with negative Iodine test had a moderate size cavity of 11×7 cm. It was also observed in this series that the duration of illness showed no correlation with the result of the seroflocculation tests.

It is also conceivable that the size of the cavity does not necessarily reflect the sole pathologic process developing in the liver. The residual normal liver tissue presented, the degree of hepatitis presented in hepatic tissue beside necrotic "abscess" area also play an important role in the result of liver function as determined by these seroflocculation. Our experience with liver biopsy in amoebic liver abscess usually revealed almost normal liver tissues presented in the areas closed to the necrotic areas thus thus minimal liver function impairment may be observed in the presence of a large abscess. On the contrary, small liver abscess with associated diffused hepatitis or hepatic degeneration may cause marked or extreme impairment of liver profile.

Glucose tolerance test Glucose tolerance test by oral method was done in 24 cases of hepatic amoebiasis. In 20 cases of amoebic liver abscess the test was normal in 17 and abnormal in 3, In 4 cases of amoebic hepatitis the test was normal in 3 abnormal in 1.

Jaundice Jaundice was present in 39 cases out of 200 cases of amoebic liver abscess accounting for 19.5% in this series. In amoebic hepatitis, jaundice was present in 21 out of 74 cases (28.4%). Maximum icterus index was 100 in amoebic liver abscess group and 37.7 in hepatitis group. (see table)

Table 8. Table showing value of icterus index in 274 cases of hepatic amoebiasis

Value of icterus index in units.	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases
0- 8.9	161	53
9-14.9	12	12
15-19.9	9	3
20-24.9	5	2
25-29.9	4	1
30-upwards	9	3
Total	200	74

From this table it is well seen that jaundice was present in about 1 in 5 cases of hepatic amoebiasis although in most instances was of mild degree. However deep jaundice could be present.

Paper electrophoresis study Paper electrophoretic study was performed in only 13 cases of amoebic liver abscess using Whatman filter paper stained with Bromphenol blue (Viranuvatti and Viranuvatti.). The result showed elevation of gamma globulin in 8 cases, α_2 in 2 and globulin in cases.

Serum enzyme determination Serum glutamic oxaloacetic transaminase and serum glutamic pyruvic transaminase were determined in 24 cases of amoebic liver abscess and one case

of amoebic hepatitis. The normal value of SGO-T was 8-40 units and SGP-T 5-35 units. The value of SGO-T and SGP-T were read spectrophotometrically using Sigma prepared substrate as reagent. The values of SGO-T and SGP-T in 24 cases of amoebic liver abscess were as follows.

Table 9. Value of SGO-T and SGP-T in 24 cases of amoebic liver abscess

Value of SGO-T in units.	No. of cases	Value of SGP-T in units.	No. of cases
0- 19	5	0- 14	9
20- 39	9	15- 34	11
40- 59	2	35- 59	3
60- 79	2	60- 79	—
80- 99	5	80- 99	—
100-119	—	100-119	1
120-upwards	1	120-upwards	—
Total	24	Total No. of cases	24

The highest value of SGO-T activity was 185 units and the lowest value was 7 units. There were 10 cases (41. %) with abnormal result (higher than 40 units) and 14 cases with normal result. In all instances the value of abnormal SGO-T level were slightly higher than normal in contrast to much higher levels found in infectious hepatitis. The value of SGP-T was abnormal in only four cases. The highest value in this series was 115 units and the lowest value 1.0 units. The abnormal values were all slightly elevated. The SGO-T value was higher than SGP-T in 19 out of 24 cases.

It is evident in this series that value of SGO-T and SGP-T is normal or slightly elevated in amoebic liver abscess, the amount of pus aspirated from the cavity or size of the abscess did not necessarily correlate with the degree of elevation of the enzyme level.

Illustrative case A male Chinese was admitted on October 12, 1959 with pain in the right lower chest and fever for 2 weeks. Examination revealed an enormously enlarged liver. Aspiration yielded 2000 c.c. of pus from the right lobe of liver. Liver function study showed SGO-T of 80 units, Iodine test 1 plus, C.C.F. 1 plus, T.T. 1.0 units, Zinc sulfate turbidity 16.0 units and B.S.P. 20.5% retention.

This case showed slightly elevated SGO-T with normal SGP-T level. The B.S.P. test showed marked dye retention in the presence of slight impairment of seroflocculation tests. In the case with highest level of SGO-T in this series (185 units) and SGP-T level of 115 units only 100 c.c. of pus was obtained by aspiration. The duration of illness bore no correlation with the degree of elevation of SGO-T and SGP-T in this series.

It was mentioned by many authors (Molander et al, Wroblewski et al) that SGO-T activity seems to be extremely sensitive index of hepatocellular integrity. The extent of any necrotising process upon the liver parenchyma appears to be reflected in the level of enzyme activity. However, the result of this study did not confirm the statement mentioned. This may be attributable in some part to that the pathology of amoebic liver abscess is focal, not diffused like those seen in viral hepatitis. The autolysed liver tissue forming abscess in cavity does not produce or release any enzyme to the circulation. Thus SGO-T and SGP-T determination in amoebic liver abscess is not so valuable or sensitive in the diagnosis of amoebic liver abscess.

SGO-T and SGP-T determination was done in only one case of amoebic hepatitis and in this case SGO-T was 126 and SGP-T 57 units. Only slight elevation of enzymatic level was observed.

Serum alkaline phosphatase Serum alkaline phosphatase determination was done in 75 cases of hepatic amoebiasis comprising 53 cases of amoebic liver abscess and 22 cases of amoebic hepatitis. The value was expressed in Bodansky units.

Table 10. Table showing the value of alkaline phosphatase in 75 cases hepatic amoebiasis

Alkaline phosphatase in Bodansky units.	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
0- 4.9	11	2	13
5- 9.9	15	8	23
10-14.9	12	7	19
15-19.9	9	1	10
20-24.9	1	2	3
25-29.9	3	1	4
30-upwards	2	1	3
Total	53	22	75

The abnormal value (more than 5 B.U.) was found in 42 out of 53 cases (79.25%) of amoebic liver abscess and 20 out of 22 (90.91%) in amoebic hepatitis. Value of more than 15 units commonly found in extra-hepatic type of obstructive jaundice was found in 15 out of 53 cases (28.30%) in amoebic hepatitis and 5 out of 22 (22.7%) in amoebic hepatitis. The average value of alkaline phosphatase in amoebic liver abscess was 17.5 B.U. and 15.9 B.U. in amoebic hepatitis. The maximum value was 38.6 in abscess and 32.9 in hepatitis whereas the minimum value was 1.4 in abscess and 6.02 B.U. in hepatitis.

From the result obtained in this study, alkaline phosphatase in hepatic amoebiasis was abnormal in 62 out of 75 total cases (82.7%). Rather high level of alkaline phosphatase was commonly found. The average value in liver abscess and hepatitis is higher than 15 B.U. comparable to that found in extra-hepatic obstructive jaundice although in most instances of cases in this series jaundice was not present. The origin of alkaline phosphatase is not yet definitely known at present. Liver cells seem to play a part in the production and delivery of this enzyme. Damaged liver cells by *Entamoeba histolytica* may cause the release of enzyme into the circulation causing the rise of enzymatic level in large number of cases.

Serum amylase Serum amylase was determined by Somogyi's method expressed in term of milligrammes of glucose liberated after digestion of starch by enzyme presented in 100 c.c. of serum. The normal level was under 100 mg. Twenty eight cases were subjected to this study.

Table 11. Table showing the result of serum amylase in 28 cases of hepatic amoebiasis

Serum amylase in mg. %	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases
0- 99.9	6	3
100-179.9	10	6
180-299.9	1	—
300-upwards	2	—
Total	19	9

In amoebic hepatitis the value were within normal limit, in amoebic liver abscess, the value was abnormal in 3 with very high value of more than 300 mg. % in 2 instances, the ma-

ximum value of which was 387.5 mg. In this case the symptoms, signs and other laboratory findings of pancreatitis or other possible cause producing hyperamylasemia was absent. The cause of elevation of serum amylase was not known.

Bromsulfalein excretion test Bromsulfalein excretion test using 5 mg. of dye per kg. body weight in 45 minutes was performed. There were 31 cases of hepatic amoebiasis comprising 21 cases of liver abscess and 10 cases of hepatitis.

Table 12. Table showing B.S.P. test in 31 cases of hepatitis amoebiasis

B. S. P. retention %	Amoebic liver abscess No. of cases	Amoebic hepatitis No. of cases	Total No. of cases
0- 4.9	2	—	2
5- 9.9	6	5	11
10-14.9	5	3	8
15-19.9	3	1	4
20-24.9	2	1	3
25-29.9	2	—	2
30-and upwards	1	—	1
Total	21	10	31

Intravenous cholecystography Intravenous cholecystography was done in 16 cases of amoebic liver abscess and 2 cases of amoebic hepatitis.

One case of amoebic liver abscess showed no dye retention and one case with deep jaundice showed maximum dye retention of 95%. Nineteen cases out of 21 cases (90.5%) in amoebic liver abscess group showed abnormal result (more than 5% retention) and in all cases of amoebic hepatitis B.S.P. excretion test was abnormal. However, moderate retention of dye was found generally. Eleven out of 21 cases of liver abscess and 8 out of 10 cases of amoebic hepatitis showed 5 to 14.9% retention. In this small series, the relationship between size of liver abscess and degree of dye retention did not exist. Visualisation of gall bladder with good concentration of dye was found in 15 out of 16 cases (93.8%). No excretion was observed in one case with presence of deep jaundice and icterus index of 66.8. In 2 cases of amoebic hepatitis, both were icteric with icterus index of 12.5 and 48.6 respectively. Intravenous cholecystogram in these two cases showed no dye excretion.

It is therefore evident that in non-icteric cases of amoebic liver abscess, intravenous cholecystography showed excretion of dye in all instances. This finding attracts our attention as the result is in contrary to the finding in primary carcinoma of liver without jaundice. Intravenous cholecystography in most instances showed no excretion of dye (Viranuvatti and Stapanakul, (1958), Viranuvatti and Plengvanich, (1960). Normal liver cells presented in the areas away from the abscess cavity seems to function well in the excretion of the dye whereas in the primary carcinoma of liver the presence of predominant liver cells and few healthy normal cells results in impaired excretion of radio-opaque dye. This result, in our experience is very significant in assisting us in the differential diagnosis between amoebic liver abscess and some cases of carcinoma of liver with clinical symptoms and signs simulation amoebic liver abscess or the "liver abscess" type of primary carcinoma of liver encountered from time to time in our institution. It is our experience that besides other laboratory findings which are usually not helpful, intravenous cholecystography without dye excretion in the abscess of jaundice usually reflects malignancy of the liver.

Overall tests of liver function and their significance in amoebic liver abscess

In this series, many liver function tests routinely available in most hospitals were done. It is evident that Iodine test was the most sensitive test in the seroflocculation group giving the highest degree of positivity. Copper acetate test showed second in the degree of abnormal result, next was Zinc sulfate turbidity (68.5%). All of these tests were based predominantly on high gamma globulin and low serum albumin resulting in close percentage of abnormal result. Cephalin cholesterol flocculation test and thymol turbidity test with different mechanisms producing abnormal result showed much lower percentage of abnormality and seemed to be not sensitive in the detection of amoebic liver abscess. This finding was in agreement to our earlier work published (Viranuvatti, 1959). Alkaline phosphatase determination also showed high percentage of abnormal result (79.25%). SGO-T, SGP-T and intravenous cholangiography were not sensitive in the detection of liver abscess in this series. Bromsulphalein showed highest degree of abnormality, however only moderate degree of dye retention was generally found as stated previously. Glucose tolerance test was not sensitive as it was abnormal in only 15%. From this series Iodine test, alkaline phosphatase determination, bromsulphalein excretion test were the most sensitive test in detection of amoebic liver abscess.

DISCUSSION It is rather astonishing that the liver function study in the diagnosis of amoebic abscess of liver was rarely mentioned in literatures despite numerous publication of the disease. In the few papers in which liver function tests were mentioned the series were too small and the revealing tests were usually omitted. Recently, Brem (1955) reported the liver function tests in 10 cases of amoebic liver abscess. Megill and Killough (1958) reported the result of liver function tests in 5 cases of liver abscess and 41 cases of amoebic hepatitis. The result were rather similar but not consistent with ours. The serum protein changes in amoebic liver abscess and hepatitis showed low total protein, albumin and globulin. The albumin and globulin ratio was lowered in the abscess group than the hepatitis group. The ratio was less than 1 in 39.6% of liver abscess and 20% in hepatitis. The duration of illness did not correlate with the alternation in serum protein, however the lowered albumin globulin ratio seemed to correlate with the size of the abscess i.e. the reverse albumin globulin ratio predominantly were seen in large abscess of more than 1000 c.c. This is conceivable on the basis of destruction of a large portion of functioning hepatic tissue. In seroflocculation tests, abnormal values were seen in a large proportion of cases on the contrary to the findings in Brem's series in which all cases showed normal cholesterol flocculation test and slight impairment of thymol turbidity test in 2 out of 10 cases. In Magill and Killough's series, cephalin cholesterol flocculation test was abnormal in 40% of hepatitis and 100% of liver abscess, however, it was also abnormal in 40% of intestinal amoebiasis and this test could not be considered to be of specific value in their study. This finding is agreeable to the result in our previous study (Viranuvatti, 1959). With reference to thymol turbidity test our abnormal result was much higher in amoebic liver abscess (39.3%) and hepatitis (36.1%). Whereas in Magill and Killough's series the abnormal thymol turbidity test was found to be 10% only in hepatitis and all normal in 5 cases of liver abscess. Indine test was of highest diagnostic value as it was positive in 83.7% of liver abscess and 78.6% of amoebic hepatitis. Iodine test was found to be positive of only 1+ in 2 out of 12 cases (16.7%) of intestinal amoebiasis in the previous study by Viranuvatti (1959). Zinc sulfate turbidity test and copper acetate test showed high degree of positivity, however comparative study with the result in intestinal amoebiasis was not yet adequate to warrant any discussion on the sensitivity of these tests. Serum glutamic oxaloacetic transaminase and serum glutamic pyruvic transaminase were not a good index of necrosis caused by *Entamoeba histolytica* as previously stated in this paper. It is also of the author's opinion that high value of SG-T and SGP-T may as well exclude hepatic amoebiasis in suspected non-icteric case. Regarding serum alkaline phosphatase and bromsulphalein

excretion test, our result was in agreement to others that serum alkaline phosphatase is usually high with retention of bromsulfalein excretion (Magill and Killough, Brem). Elevation of alkaline phosphatase and retention of B.S.P. unaccompanied by jaundice as observed in their group were also rather consistent findings in our series excluding icteric cases. The divergence of liver function tests is very likely dependent upon the obstruction of a part of the intra-hepatic bile duct without extensive or diffused cellular damage. Freeman et al (1938). Schiffman and Wilkelman (1939) demonstrated in experimental animal that ligation of a portion of the intra-hepatic bile duct produced striking rise in the serum alkaline phosphatase with no increase in serum bilirubin. Gutman (1940) ascribed this to the fact that bilirubin is readily excreted by the kidneys while phosphatase is not. In our series the degree of alkaline phosphatase level and B.S.P. were not proportionate necessarily to the size of liver abscess or extent of the lesion. The average value of alkaline phosphatase was slightly higher in hepatitis than in liver abscess. Serum amylase determination in this series was not significant in the diagnosis of hepatitis or liver abscess. Glucose tolerance test was abnormal in low percentage of cases and therefore was of no diagnostic significance. Hyperbilirubinemia could be found in a fair percentage and in some instance deep jaundice was also present. This showed that jaundice not exclude the possibility of amoebic affection of the liver although in most instances jaundice is absent. Paper electrophoretic study in this series was too small to warrant any conclusion of its value in the diagnosis. However, from our result here, in cases suspected clinically of having amoebic liver abscess, the positive Iodine test especially of high degree, elevated alkaline phosphatase, bromsulfalein retention, good excretion of dye intravenous cholecystography and slightly elevated or normal SGO-T, SGP-T, are very good laboratory criteria for the diagnosis of amoebic liver abscess. The diagnostic probability was almost certain in our series.

Summary and conclusions Liver function tests commonly used in the laboratory were performed in 274 cases of hepatic amoebiasis. There were 200 cases of amoebic liver abscess and 74 cases of amoebic hepatitis. In amoebic liver abscess the average serum albumin was 3.1 and globulin 2.8. In amoebic hepatitis the average serum albumin was 3.3 and globulin 2.1. The value of albumin globulin ratio was less than 1 in 39.6% of cases of liver abscess and 20.0% of hepatitis. Iodine test was abnormal in 83.7% of liver abscess and 79.6% of hepatitis. Cephalin cholesterol flocculation test was abnormal in 43.2% of liver abscess group and 53.6% of hepatitis group. Thymol turbidity test was abnormal in 39.3% of liver abscess and 36.1% of hepatitis. Zinc sulfate turbidity test was abnormal in 68.5% of liver abscess and 65.2% of hepatitis. Copper acetate test was abnormal in 71.4% of abscess cases. The result of all sero-flocculation tests studied showed no relationship with the size of cavity of liver abscess as determined by measurement on A-P X-ray film after intracavity instillation of radio-opaque Indine compound or the duration of illness. Oral glucose tolerance test was abnormal in 3 out of 20 cases of liver abscess and 1 in 4 cases of hepatitis. Jaundice was present in 19.5% of amoebic liver abscess and 28.4% of hepatitis. Deep jaundice was occasionally found. Paper electrophoresis study was performed in only 13 cases of amoebic liver abscess, elevation of gamma globulin was observed in 8, alpha₂ in 7 and beta in 1. Serum glutamic oxaloacetic transaminase and serum glutamic pyruvic transaminase were done in 24 cases of amoebic liver abscess. SGO-T level was abnormal in 41.2% and elevation was only slight. SGP-T was slightly higher than normal in 4 out of 24 cases. Serum alkaline phosphatase usually showed high abnormal values in both groups with average value more than 15 B.U. Serum amylase was usually normal. Bromsulfalein excretion test showed abnormal result in 90.5% of amoebic liver abscess group and 100% in hepatitis group. Moderate retention of dye was generally observed. Intravenous cholecystography showed visualisation of gall bladder in all non-icteric cases in contradistinction to our findings in primary carcinoma of liver. From the result of this study, in cases suspected

clinically of having amoebic liver abscess, positive Iodine test especially of high degree, elevated alkaline phosphatase, B.S.P. retention, good excretion of dye after I.v.c, slightly elevated or normal SGO-T and SGP-T are very good laboratory criteria for the diagnosis of amoebic liver abscess.

Acknowledgement The authors wish to express their sincere appreciation to Prof. Presert Kangsadal for his guidance.

*

*

*