Brain plasticity in health and disease

Barbro B Johansson

Division for Experimental Brain Research, Department of Clinical Neuroscience, Lund University, Wallenberg Neuroscience Center, Lund, Sweden

(Received for publication on August 2, 2004)

Abstract. Research during the last decades has greatly increased our understanding of brain plasticity, *i.e.* how neuronal circuits can be modified by experience, learning and in response to brain lesions. Currently available neuroimaging techniques that make it possible to study the function of the human brain *in vivo* have had an important impact. Cross-modal plasticity during development is demonstrated by cortical reorganization in blind or deaf children. Early musical training has lasting effects in shaping the brain. Albeit the plasticity is largest during childhood, the adult brain retains a capacity for functional and structural reorganization that earlier has been underestimated. Recent research on Huntington's disease has revealed the possibility of environmental interaction even with dominant genes. Scientifically based training methods are now being applied in rehabilitation of patients after stroke and trauma, and in the sensory retraining techniques currently applied in the treatment of focal hand dystonia as well as in sensory re-education after nerve repair in hand surgery. There is evidence that frequent participation in challenging and stimulating activities is associated with reduced cognitive decline during aging. The current concept of brain plasticity has wide implication for areas outside neuroscience and for all human life. (Keio J Med 53 (4): 231–246, December 2004)

Key words: neuronal plasticity, development, dystonia, stroke, aging

Introduction

The unique evolution of the human brain is the basis for the development of human culture, science and technology. Our neurons are tuned in close interaction with our environment, our own activities and thoughts. Interaction between genes and environment is evident from the global variations in customs, languages, and culture. Ernesto Lugano, an Italian psychiatrist may have introduced the term plasticity in neurosciences as early as 1906.¹ However, the important studies in this field were initiated by Donald Hebb, who more than half a century ago postulated that neuronal cortical connections are strengthened and remodeled by our experience.² He showed that rats that were allowed to run around freely in his house were better learners and had better memory capacity than rats housed in the laboratory.³ Many studies have shown that an activity-stimulating environment has multiple effects on the brain including increasing the number of neuronal connections.^{4–9}

Dendritic spines, *i.e.* tiny protrusions on the dendrites (Fig. 1)¹⁰ are the primary postsynaptic targets of excitatory glutamergic synapses in the mature brain, and they have been proposed as primary sites of synaptic plasticity.^{11,12} The dendritic tree is covered with a variety of excitable synaptic channels operating on different time scales and with activity-dependent sensitivity enabling a sophisticated neuronal plasticity.¹³ The spine cytoskeleton consists of actin filaments, and video recordings from hippocampal neurons expressing actin tagged with fluorescent protein have shown that the shape of spines can change rapidly,¹⁴ events that are accompanied by changes in calcium influx and decay.^{15,16} With a technique enabling *in vivo* images of neurons expressing green fluorescent protein, the lifetime of superficial pyramidal spines in the mouse barrel cortex has been observed to vary greatly.¹⁷ About 20% of spines disappeared from one day to the next, a loss that was balanced by formation of new spines. About 60% of dendritic spines persisted for at least 8 days, and of those 17% had disappeared one month after the

Presented at the 1345th Meeting of the Keio Medical Society in Tokyo, February 10, 2004.

Reprint requests to: Dr. Barbro B Johansson, Wallenberg Neuroscience Center, BMC A13, SE- 221 84 Lund, Sweden, telephone: +46-46-222 0621, facsimile: +46-46-222 0626, e-mail: Barbro.Johansson@neuro.lu.se

Fig. 1 Pyramidal neurons in cortical layer III in male 4 months' old spontaneously hypertensive rats housed in standard environment (S), or housed in an enriched environment (E). Standard housing = 3 rats in each cage; enriched housing = 8 rats in larger cages with opportunities for various activities. The environmental enrichment significantly increases dendritic branching and number of dendritic spines (insertions). (Reproduce from, Johansson BB, *et al*: In: Bazan NG, *et al* eds, *Maturation Phenomenon in Cerebral Ischemia IV*, Berlin Heidelberg, Springer Verlag, 2001; 77-23, Copyright © (2001), with permission from Springer Verlag). Scale bar = 25 μ m; in the insertions scale bar = 5 μ m.

imaging began. Serial-section electron microscopy of imaged dendritic segments revealed retrospectively that spine sprouting and retraction are associated with synapse formation and elimination. Trimming every other whisker, which induced a rapid and robust remodeling of whisker representation, increased the pool of transient spines present for only a day or less in the cortical representation areas of the trimmed whiskers.¹⁷ Thus, experience-dependent plasticity of cortical receptive fields was accompanied by increased synapse turnover. There may, however, be regional differences in the plasticity of dendritic spines. Using a similar *in vivo*-technique in the visual cortex the majority of spines in layer V pyramidal neurons were more stable.¹⁸ Axons and dendrites need guidance for proper orientation, and a number of guidance molecules or surface receptors have been identified. An expression of a common attractant in the neuropil, and activation of its corresponding receptors in axons of one neuron and dendrites of another, might be essential for neuronal network formation during development and plasticity.¹⁹

As will be exemplified in the following sections,

in addition to strengthening contacts between individual neurons, cortical representation areas, "cortical maps" can be modified by sensory input, experience, training (Fig. 2)²⁰ and in response to brain lesions. Synaptic plasticity and number and turnover of dendritic spines in cortical horizontal connections are likely to underlie cortical map reorganization.^{21,22} Activity-dependent modification of synaptic connection and reorganization of adult cortical areas are thought to involve long-term potentiation (LTP) and long-term depression (LTD), mechanisms by which information is stored in the mammalian central nervous system.^{23,24} Despite 30 years research the detailed mechanisms remain unclear.²⁵ LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites.²⁶ In a recent study LTP, induced by intracellular electrical micro stimulation, expanded movement representation and induced dendritic hypertrophy in rat sensorimotor cortex.²⁷ Glutamate, the main excitatory neurotransmitter plays a crucial role, and cortical map reorganization in the primary somatosensory cortex can be prevented by blockade of N-methyl-d-

Fig. 2 The somatosensory cortex of the hand in a monkey before (A) and after (B) repetitive specific tactile stimulation of the tip of the second finger (black areas). (Reproduce from, Jenkins WM, *et al*: J Neurophysiol 1990; 63: 82–104, Copyright © (1990), with permission from The American Physiological Society).

aspartate (NMDA)-receptors. Gamma-aminobutyric acid (GABA)-A receptor antagonists facilitate LTP induction, and the induction can be blocked by GABA-A receptor agonists.^{28,29} Also AMPA receptors play an important role.³⁰ Transmitters released by diffuse neuromodulatory systems that originates in subcortical regions and send long projection axons to the cortex, *i.e.* noradrenalin (locus coeruleus), acetylcholine (nucleus basalis), dopamine (lateral tegmentum), and serotonin (raphe nuclei), can modify the threshold for activity-dependent synaptic plasticity, maybe through a facilitation of NMDA receptor-gated process.^{31–34} Another modulator of cerebral cortex synaptic function is nitric oxide³⁵ that has been shown to interact with dopamine and acetylcholine to induced corticostriatal synaptic plasticity.³⁶ The molecular mechanisms behind activity-dependent modification of synaptic connections and reorganization of cortex in the adult brain are complex and still not fully understood, and they are likely to differ with the regions studied.^{22,37}

Skill learning but not strength training induces cortical reorganization.^{38,39} Exercise induces angiogenesis but does not alter movement representations in the cortex.⁴⁰ It has recently been reported that cortical synaptogenesis and motor map reorganization occur during late but not early phase of motor skill learning, and that synaptogenesis precedes map reorganization.

It was proposed that motor map reorganization and synapse formation do not contribute to the initial acquisition of motor skills but represent the consolidation of motor skill that occurs during late stages of training.⁴¹ There is increasing evidence that non-neuronal cells, in particular astrocytes, have essential roles in plasticity. Astrocytes are responsible for regulating the synaptic environment and for maintaining appropriate levels of neurotransmitters and neurotrophins, and they influence neuronal signaling, number of synapses and play a role in non-synaptic transmission.^{42–46}

Neural stem cells, defined by their ability to both self-renew and to differentiate to produce mature progeny cells and neurons, are present in the sub-ventricular zone and in the hippocampus in the adult human brain.^{47,48} Animal studies have shown that neurogenesis can be stimulated by many factors including trophic factors, hormones, drugs, environment and brain lesions. Environmental enrichment enhances the survival of newly formed neurons in the hippocampus of intact rats and mice and enhances spatial memory.^{49–51}

One important factor restricting brain plasticity after brain lesions is that myelin in the central nervous system contains proteins that inhibit axonal outgrowth.^{52,53} The myelin inhibitors share a common receptor and blockage of the receptor promotes CNS

repair and functional recovery.⁵⁴ Some promising results with antibodies against this receptor have been reported after spinal and focal brain lesions in animal studies.^{55–57}

Interestingly, inosine, a naturally occurring metabolite, has likewise been reported to stimulate axonal growth in the rat corticospinal tract,⁵⁸ and to induce axonal rewiring and promote behavioral outcome after focal brain ischemia.⁵⁹ It is an important area of research still at the experimental level.

Neuroimaging Methods Used in the Study of Brain Plasticity

Brain plasticity in the human brain can be studied with several neuroimaging techniques with different temporal and spatial resolution. Thus the temporal resolution varies from minutes in positron emission tomography (PET), seconds in functional magnetic resonance (fMRI) and milliseconds in magnetoencephalography (MEG), which makes MEG best suited for studying temporal spatial patterns of brain activity, that is, where and when the activation starts, and its temporal spread to other regions.⁶⁰ Because of the better spatial resolution in fMRI, the wider availability of the technique and the possibility to combine with other MRI techniques, fMRI has been widely used in studies on brain plasticity.^{61–63}

Transcranial magnetic stimulation, TMS, is another neurophysiological technique allowing studies of brain functions with a millisecond resolution, which has developed into a popular non-invasive tool for studying the human brain.^{64–66} A pulsed magnetic field creates current flow in the brain and can temporarily excite or inhibit synaptic efficiency in specific areas and alter brain function.^{64–66} Multiple overlapping motor representations are functionally connected through an extensive horizontal network. By changing the strength of horizontal connections between motor neurons, functionally different neuronal assemblies can form. Delivery of multiple single pulses of TMS does not leave any lasting effect, but repetitive TMS can produce effect that last for some time after the stimulation period. The cortical motor and sensory representations of each hand exert inhibitory influences on the homonymous representation in the opposite hemisphere.⁶⁷ Cutaneous anesthesia of the right hand enhances spatial acuity and cortical processing in the left hand,⁶⁸ and tourniquet-induced anesthesia of the right hand improves tactile discrimination and grip strength, lasting at least 15 minutes after anesthesia and associated with increased fMRI activation in the right primary motor cortex.⁶⁹ Furthermore, repetitive TMS over the left frontal cortex can enhance analogical reasoning,⁷⁰ and the encoding of a motor memory in the primary motor

cortex can be enhanced by TMS stimulation in healthy volunteers.⁷¹

Brain Plasticity in the Development and in the Adult Human Brain

During the early childhood there is a considerable capacity for cross-modal plasticity⁷² with redundant connections between auditory and visual areas that gradually decrease between 6 and 36 months of age.⁷³ Speech processing and auditory localization activate the visual cortex in congenitally blind humans,^{74,75} and blind humans localize sound sources better than sighted subjects.⁷⁶ Cerebral organization for language functions differs in deaf and hearing subjects, and sign language activates different regions in hearing and deaf signers.⁷⁷ The sensorimotor cortical representation of the reading finger is expanded in blind Braille readers⁷⁸ and fluctuates with the daily reading activity.⁷⁹ Furthermore, tactile acuity is significantly superior in blind subjects, independently of the degree of childhood vision, light perception level, or Braille reading.⁸⁰ Linguistic differences have been demonstrated in cortical organization for language in adult individuals with less hemispheric asymmetry in Chinese than in English and Spanish language.⁸¹

Exposure to musical training in early life shapes the brain, and the brains of musicians have been extensively studied as models of neuroplasticity.^{82–84} Pianists have to memorize long and complex bimanual finger sequences, and to translate musical symbols into motor sequences during sight-reading. Long-term motor training and continued practice of complicated bimanual motor activity lead to changes in the brain structure and representation of the brain cortical motor maps. The anterior part of the corpus callosum, consisting of nerve fibers connecting frontal motor-related regions and prefrontal regions crucial for the coordination of bimanual motor activity, is larger in musicians that started musical training before the age of 7 than in musicians without an early start, or in a control population.⁸⁵ With diffusion tensor imaging, a technique that allows evaluation of the architecture of the brain white matter, adult subjects with musical training since early childhood differ significantly from control subjects in areas suggesting both cognitive and motor effects of early musical training.⁸⁶ Another example is the increased cortical representation of the fingers of the left hand in string players that correlates with the age at which the person had begun to play.^{87,88}

Auditory experiences during early postnatal development are important in shaping the functional development of auditory cortical representation, and musicians have increased areas in the auditory cortex for functions important for music such as timbre.^{89,90} The

functional anatomy of musical perception, *i.e.* brain activation during a passive music listening task, differs significantly between musicians and non-musicians.⁹¹

Although the developing brain is clearly more plastic than the adult brain, neuronal connections are continuously remodeled by experience and by performance of specific and complex movements during motor and cognitive learning, and memory processes involve brain plasticity. Rapid and transient alteration of cortical representation areas are seen during learning tasks, most likely related to unmasking of previously existing connections, perhaps as a result of decreased inhibition and increased synaptic efficacy in existing neuronal circuits.⁹² Flexible short-term modulations are important in the acquisition of new skills, and can lead to structural changes in the intracortical and subcortical networks as the skill becomes more established and automatic.⁹³ Even mental practice on a piano keyboard can change the cortical representation of finger flexors and extensor muscles in adult subjects.⁹⁴

Another example of structural plasticity in the adult human brain comes from a study on London taxi drivers, who have a 2-year-long training period in special localization before qualifying as drivers. The volume of gray matter in the right posterior hippocampus, which stores a spatial representation of the environment, is greater than in controls, and the volume increase correlates significantly with the amount of time spent learning to be and practicing as a licensed taxi driver.⁹⁵ The failure to find an association between hippocampal volume and navigation expertise in other subjects supports the conclusion that the difference in volume is not an innate genetic difference, but reflect the detail and/or duration of training and use of the spatial representation necessary in the daily work of the drivers.⁹⁶

Examples of the interaction between a dominant gene and environment have been demonstrated in Huntington's disease, a late onset progressive genetic disorder characterized by motor dysfunction, personality and cognitive disturbances leading to dementia and premature death. In a transgenic mice model of Huntington's disease, environmental stimulation significantly delayed the onset of the disease and improved life expectancy.^{97–99} In human patients with Huntington's disease the cannabinoid CB1 receptors are lost from the basal ganglia output nuclei prior to the development of other neuropathology,¹⁰⁰ and the improvement during environmental enrichment in an animal model correlated with delayed loss of cannabinoid CB1 receptors.¹⁰¹ Furthermore, differential clinical and motor control function and different age onset have been reported in pairs of monozygotic twins with Huntington's disease, demonstrating that environmental factors can interact even with dominant genes in humans.^{102,103}

Maladaptive Brain Plasticity

Focal hand dystonia, often called writer's cramp or musician's cramp, is a disabling condition that so far has been very difficult to treat with physiotherapy or other methods. Thus, a recent long-term outcome study on 21 violin and viola players with focal dystonia an average 13.8 years after onset of symptoms showed that only 38% had been able to maintain their professional careers.¹⁰⁴ Focal hand dystonia is now considered to be a disorder of maladaptive plasticity. In adult monkeys it has been shown that frequent very rapid time-synchronous movements during vigorous practicing can cause a degradation of the sensory feedback controlling fine motor movements, resulting in fusion of the distinct cortical representation of the individual fingers.¹⁰⁵ The abnormal sensory representation interferes with motor control, and abnormal motor control strengthens the sensory abnormality, and the positive feedback loop reinforces the dystonic condition. The sensory representation abnormalities parallel focal hand dystonia in the primate model.¹⁰⁶ Patients with focal hand dystonia have impaired intracortical inhibition and abnormal cortical activations during movements with affected muscles,^{107–109} and the regional cerebral blood flow correlates with the severity of symptoms.¹¹⁰ Compared to control individuals, patients with dystonia have significantly reduced levels of the inhibitory neurotransmitter GABA in the sensorimotor cortex and lentiform nuclei contralateral to the affected hand as demonstrated with *in vivo* magnetic resonance spectroscopy.¹¹¹ Based on the hypothesis that successful treatment should be aimed at restoring the normal representation areas of sensory feedback from the hand, methods based on relearning principles have been developed, and substantial improvement has been reported from three centers including a one year follow up study.^{112–115} The clinical improvement is associated with altered somatosensory cortical organization.¹¹⁵ A recent voxel-based morphometry study has demonstrated increase in the gray matter in the hand representation area of primary somatosensory cortex, and to a lesser extent in primary motor cortex in patients compared to controls, suggesting that there might be a hereditary component predisposing for the syndrome.¹¹⁶

Embouchure dystonia is a focal task-specific disorder in professional brass and woodwind players involving abnormal non-coordinated movements and involuntary muscle contraction of lip, jaw, and tongue muscles used to control the flow of air into a mouthpiece while playing.¹¹⁷ It is often so disabling that the musicians have to limit or give up their occupation. There is evidence that abnormal somatosensory reorganization contributes to the disorder.¹¹⁸ Thus, in a study comparing 8 patients

and 8 control subjects, the patient's digit and thumb representations were shifted in a lateral direction towards the lip representational zone. Furthermore, the patient's upper lip showed decreased sensitivity compared to their lower lips, an asymmetry that was absent in controls.

A strong correlation between the subjective strength of tinnitus and reorganization of the auditory cortex in individuals with hearing loss and tinnitus has led to the proposal that this is another examples of maladaptive plasticity.¹¹⁹

Brain Plasticity due to Lack of Sensory Input to an Intact Brain

Lack of sensory input due to amputation, peripheral nerve lesions or local anesthesia, leads to rapid cortical remodeling with a distorted cortical representation and enlarged and overlapping cortical receptive fields.^{120–123} By simultaneous recording of extracellular activity from somatosensory cortex, the thalamus, and trigeminal brainstem area in adult rats, an immediate and simultaneous transient reorganization was produced by subcutaneous lidocaine injection, demonstrating that peripheral sensory deafferentation triggers a system-wide reorganization, and that spatio-temporal cortical plasticity is paralleled by subcortical reorganization.¹²⁴ Studies on monkeys several years after an amputation have indicated that much of the large-scale cortical reorganization that occurs after a major loss of peripheral inputs reflects the sprouting or expansion of afferents from the remaining forelimb stump into deprived territories of the spinal cord and brain stem. Since the topographic representation of the body is greatly magnified in the cortex, the subcortical changes can result in dramatic cortical map changes.^{125,126} Reorganization of the motor efferents to muscles has been demonstrated by a combination of intracortical microstimulation of primary motor cortex and *in vivo* labeling of corticospinal neurons.¹²⁷ Stimulation in the deafferented forelimb cortex evoked shoulder stump, trunk and orofacial movements, whereas stimulation in the deafferented hindlimb cortex evoked hip stump, trunk and tail movements. No unresponsive zones were found in the deprived cortex. The reorganization in sensory afferents along with reorganization of the motor efferents to muscles may provide a basis for phantom sensations of movements in the missing forelimb.¹²⁶ A recent fMRI study from human amputees indicates that, in addition to primary sensory cortex, somatosensory cortex, thalamus, posterior parietal cortex, and prefrontal cortex may be involved in the phantom sensation.¹²⁸ Central maladaptive plasticity has also been proposed to contribute to phantom pain.^{129,130} Repetitive transcranial magnetic stimula-

tion of the contralateral parietal cortex can transiently but not lastingly ameliorate phantom limb pain, supporting the hypothesis that phantom pain at least in part is due to a dysfunctional activity in the parietal cortex.¹³¹

Reconstruction of injured peripheral nerves is a challenging surgical problem. There is no microsurgical technique that makes it possible to reinstate the prelesion axonal connections, and the cortical representation of the reinnervation of the hand will be defragmented.¹³² The axonal growth is slow, and the reinnervation of a hand therefore takes time. Sensory hand training has usually not started until the time reinnervation is taken place. However, because functional reorganization of the cortex occurs rapidly after changes in peripheral input, timing of the onset of sensory-relearning may be critical.¹³³ A new concept is that sensory relearning programs, aimed at refinement of the distorted receptor fields, should start early to try to normalize the hand map and improve processing at a high-order cortical level. This can be done by taking advantage of the fact that sensory deprivation in one sensory system can have effect of the function of the others due to a continued functional interplay between the different association areas in the adult brain.⁷² The hearing sense has a high capacity to discriminate between complex patterns of frequencies, and when touched, individual textures are associated with specific friction sounds. The hypothesis that hearing would be able to take over functions normally devoted to touch after lesions to the hand has led to the development of a Sensor Glove system, which stereophonically transposes the friction sound elicited by active touch. Thus the individual can listen to what the hand is touching and the training can be initiated before the reinnervation of the hand takes place. Preliminary results from a prospective clinical randomized study indicate that use of this re-learning system improves recovery of tactile discrimination in the nerve-injured hand (132).

Lesion-induced and Training Induced Brain Plasticity

Despite permanent tissue loss most surviving stroke patients regain some function with time. In 1973 a neuroanatomist wrote in a paper based on his own experience after a stroke "since regeneration of transected central axons has never been convincingly demonstrated in higher mammals, it seems that one must resort to the assumption that intact fibers take over from the damaged ones".¹³⁴ That is indeed the case and many molecular and neurophysiological mechanisms have been proposed to take part in lesion-induced brain plasticity.^{66,135–142}

That reorganization of cortical representation after brain injury could be the neurophysiological model of

the bases of recovery from cortical lesions was first suggested by Jenkins and Merzernich,¹⁴³ a hypothesis that has been supported by many later studies. A lesion to a small portion of the cortical motor hand representation in monkeys results in a further loss of hand territory in the adjacent undamaged cortex. Retraining of skilled hand use improves behavioral recovery and prevents tissue loss, and the hand representation may expand into regions formerly occupied by representations of the elbow and shoulder.¹⁴⁴ Similarly, cortical reorganization of primary somatosensory cortex is associated with sensorimotor skill recovery after cortical lesions in adult monkeys,¹⁴⁵ and cortical reorganization induced by motor learning is associated with synaptogenesis.¹⁴⁶

The spontaneous posts ischemic functional improvement after an ischemic infarct in the hand representation area of primary motor cortex has been studied in monkeys.¹⁴⁷ An initial post infarct motor impairment of the contralateral hand was followed by a gradual improvement over three months. Intracortical microstimulation mapping at 12 weeks revealed substantial enlargement of the hand representation in the ventral premotor cortex remote from the lesion. The enlargement was proportional to the amount of hand representation destroyed, that is the greater the damage to intracortical pathways, the greater the plasticity in intact areas.¹⁴⁷

In addition to training-induced alteration in cortical maps, there is evidence that the post-ischemic environment can influence functional outcome after brain lesions.^{148–151} Thus, environmental enrichment improves functional outcome after experimental brain infarcts,^{152,153} increases the number of dendritic spines,⁹ alters gene expression pattern,^{154,155} influences the lesion-induced progenitor cell differentiation¹⁵⁶ augments the effect of training,¹⁵⁷ and interacts with drug treatment.¹⁵⁸ Furthermore, it enhances the efficacy of neocortical transplantation.^{159,160} Similar results have been obtained after striatal transplantation in animal models of Parkinson's disease.¹⁶¹

The mechanisms that mediate functional recovery after stroke may differ depending on time after the stroke, lesion site and degree of recovery.^{137,139,140,162} A controversial question has been to what extent changes in the intact hemisphere are responsible for good outcome in human stroke patients. Early PET studies on blood flow distribution during finger movements in a previously paretic hand demonstrated complex patterns of activation with multiple areas of increased activation on both sides, and with large individual variations.^{163,164} Later studies with fMRI suggest that the activation in intact regions in the lesioned hemisphere is most closely related to good recovery in human patients.^{165–168} Thus, Carey *et al.*,¹⁶⁵

who studied the effect of training several years after stroke onset, demonstrated a shift of activation from the intact to the lesion side in connection with improved function. A comparison between patients with poor and good recovery indicated that patients with poor recovery were more likely to recruit a number of motor-related brain regions during a motor task, whereas patients with more complete recovery were likely to have a task-related brain activation that were more similar to control subjects with no stroke.¹⁶⁶ Likewise, results from studies evaluating the effect of transcranial magnetic stimulation on motor performance of the paretic hand of chronic stroke patients indicated that recovered function in a paretic hand of chronic stroke patients relies predominantly on reorganized activity within sensorimotor regions in the affected hemisphere.^{169–171} However, it has been proposed that in some patients with subcortical lesions and severe damage to the corticospinal tract, a slow recovery might be obtained by the utilization of the intact contralateral sensorimotor cortices.¹⁷²

Thus most data today indicate that the substrate for improved functional recovery is mainly in regions ipsilateral to the lesion, and that high activity in the intact hemisphere is connected with less good recovery. In agreement with the concept of inter-hemispheric inhibition described earlier,^{67–69} the primary motor cortex of the intact hemisphere could influence functional recovery through trans-callosal connections. Interestingly, a very recent study indicated an abnormally high inter-hemispheric inhibitory drive from the intact to the affected hemisphere during voluntary movement by the paretic hand.¹⁷³ The authors commented that it is conceivable that this abnormality could adversely influence motor recovery in some patients with subcortical stroke, clearly an issue for future investigations. If so, it would be expected that transient deafferentation of the intact side^{67–69} would improve the motor function on the weak side.

Basic research on brain plasticity has lead to introduction of new techniques in neurorehabilitation. One such method, based on extensive studies in monkeys by Taub and coworkers, is constraint-induced training.^{174–175} Already in 1940 it was observed that monkeys with lesions to one pyramidal tracts failed to use the affected limb, and that the use of the weak limb could be improved if the unaffected limb was restrained.¹⁷⁶ By preventing the patients from using the healthy arm and hand and thus intensify the training, together with what the authors call shaping that includes making a task successively more difficult, the cortical representation maps can change together with improved performance.^{177–178} The major part of the effect is probably caused by the intense training and the relative importance of preventing movement in the intact hand is unclear.¹⁷⁹ A multisite randomized trial to

evaluate the therapeutic success, the influence of post-stroke time before starting the training, and the persistence of any effect over 2 years, is ongoing in the USA.¹⁸⁰ Good effects have been obtained also with techniques such as training with robots that do not require any immobilization of the intact side.^{181–185} The intensity of the training seems to be a crucial factor in the studies reported beneficial effects of specific training.

Training with a mirror is another interesting approach. Arm amputees can experience the perception of movement of a phantom limb while looking at a mirror reflection of the moving, intact arm superimposed on the perceived phantom. To provide illusory visual feedback of movement such use of a mirror has been tried in rehabilitation of hemiparetic patients.^{186–187} Theoretically this method is interesting in light of current knowledge of the so-called mirror neuron system. Seeing a purposeful action activates neurons in the prefrontal regions that are activated both by performing and observation of purposeful actions,^{188–190} neurons that have connections with many regions of the brain and are thought to be essential for the human capacity to imitate and understand other persons intention. Virtual reality is another technique that is under evaluation.¹⁹¹ Mental training activates most of the same circuitry in the brain as real action does,^{94,192} and mental practice is another method preliminary reported to be of value.¹⁹³

Cognitive deficiencies are common among stroke patients and memory problems, attention and ability to concentrate are likely to influence all training. Thus, the effect of mental practice after stroke is related to the working memory capacity of the patients.¹⁹⁴ Most stroke patients have better function in the shoulder and upper arm than in the hand. Since there is competition among body parts for territory in the sensorimotor cortex (Fig. 2), reducing the sensory input from surrounding areas by transient deafferentation of the upper arm during hand motor practice dramatically improved hand motor function, including some activities of daily living.¹⁹⁵

It should be stressed that few of these new techniques have been properly evaluated in larger clinical studies yet. However, the promising neurobiological research is changing the attitude to neurorehabilitation. Collaboration between basic and applied science will continue to be important for further progress. Most of the spontaneous functional sensorimotor improvement occurs within the first 3–6 months after a stroke, and that is the optimal time for rehabilitation. However, specific training can alter cortical representation areas even years after stroke.¹⁶⁵ An important aspect is that training has to be continued to keep up functional gains. Clinical studies indicate that the patient's atti-

tude, activities and social interaction influence functional outcome and quality of life after brain lesions. Rehabilitation strategies that are meaningful for the individual patient are likely to be the most effective.¹⁵⁰

Additional ways of enhancing brain plasticity is being tested in animals, including the trials to find ways to enhance axonal outgrowth mentioned in the introduction.^{52–59} Another area of great interest is whether lesion-induced stem cell proliferation can have beneficial effects after brain lesions. As demonstrated by several recent reviews,^{196–199} many studies have shown a marked increase in stem cell/progenitor cell proliferation in hippocampus and the subventricular zone after global and focal brain ischemia. That endogenous neural precursors can be induced to differentiate into mature neurons in adult mammalian neocortex and form long-distance cortico-thalamic and cortico-cortico connections has been shown in a model of apoptotic selective death of cortical pyramidal neurons.^{200,201} In another model of selective neuronal death, global ischemia, intraventricular infusion of FGF-2 and epidermal growth factor (EGF) during 4 days markedly increased regeneration of hippocampal pyramidal neurons with electrophysiological evidence that the neurons were integrated into the existing brain circuitry, and it was proposed that the new cells contributed to the amelioration of neurological deficits observed in the treated rats.²⁰² After focal brain ischemia, proliferating cells from the subventricular zone migrate to the striatum after large striato-cortical lesions induced by a transient MCA occlusion, a small part of which may develop into medium sized striatal neurons.^{203,204} However, no study has so far demonstrated integration into neuronal circuits or functional effects of neurogenesis after focal brain ischemia. It is possible that cell or tissue transplantation may be more successful, an issue with many aspects that will not be dealt with here.

Aging and Brain Plasticity

To what extent environmental factors can influence the aging processes is controversial. In experimental animals, environmental and social interactions can partially offset age-related decrease and fragmentation of the somatosensory cortical maps.²⁰⁵ 14 days of exposure to an enriched environment in 28-month-old mice significantly enhanced spatial memory and synaptophysin levels in hippocampus and frontoparietal cortex.²⁰⁶ Mice living in an enriched environment from the age of 10 to 20 months had a fivefold higher hippocampal neurogenesis together with significantly better learning ability and exploratory behavior than control rats in a standard environment.²⁰⁷ Twin studies have shown that brain volume, the size of corpus callosum and the size of the ventricles are highly genetically deter-

mined whereas more variability has been found in the gyral and sulcal patterns.^{208–210} In a large twin study about 60% of the variance of the temporal horns and 80% of the variance of corpus callosum were considered to be attributable to genetic components compared to 40% of the variance of hippocampal volume, suggesting that the environment or gene-environmental interactions have greater control in modifying hippocampal size,²¹¹ a conclusion that is consistent with results from an MRI study of hippocampus in monozygotic twins with discordant cognitive abilities.²¹²

Studies with PET and fMRI indicate that elderly people activate more regions than younger even for simple motor task, with no difference in accuracy but increase in reaction time and greater activation in the same areas as activated in the young, and with recruitment of additional cortical and subcortical areas.^{213–215} It has been interpreted to support the notion that an adaptable and plastic motor network is able to respond to age related structural and neurochemical changes in order to maintain performance levels.^{214,215} In an episodic memory-encoding task older adults engaged much of the same neural circuitry as younger subjects. In addition, age-related differences in prefrontal and temporal activity were observed during successful episodic encoding. One difference was a bilateral activity in prefrontal regions in the older subjects compared to a left activation in young.²¹⁶ More bilateral patterns of prefrontal activity in older adults than in younger adults have also been shown during working memory and visual attention tasks, suggesting that older adults may use selective compensatory mechanisms in processing information.²¹⁷

The effect of practice and of cognitive ageing has been studied in a large longitudinal study of elderly community residents in the UK.²¹⁸ Age-related changes were relatively slight between 49 and 70 years and much more marked between 70 and 80 years. However, the population average effect of practice was large relative to the age-related change, and the variation between individuals increased with age. This study provided significant empirical evidence that individuals age at markedly different rates. A large population-based studies of older persons in the USA in a model that controlled for baseline level of cognition, age, sex, race and education, demonstrated that frequent participation in cognitively stimulating activities was associated with significantly reduced cognitive decline in older persons, an effect that remained when the investigators controlled for depressive symptoms and chronic medical conditions.²¹⁹ Furthermore, frequent participation of cognitive stimulating activities appeared to be associated with reduced risk for Alzheimer's disease.^{220,221}

Aging is accompanied by a decline in muscle strength and ability to maintain submaximal force. However, skilled finger movement training improves the ability to control submaximal pinch force, hand steadiness and manual speed also in elderly individuals.²²² An investigation with magnetic resonance spectroscopy in different brain regions before and after 5 weeks of memory training in healthy elderly showed that memory improvement was accompanied by elevation of creatine and choline signals in the hippocampus. Interestingly, those considered to be at risk for neuronal dysfunction because of low neurometabolites at baseline demonstrated the largest increases after training.²²³

A Finnish study in which the participants were grouped according to their own identification with age suggested an association between perceptions of personal aging and physical and psychological well being.²²⁴ Another study on Swedish and Canadian samples of older adults suggested that subjective health should be added to the growing number of individual-different variable that are predictive of episodic memory change in very old age.²²⁵ Together with the recent demonstrations on neurobiological effects of expectation, including activation of endorphin pathways in individuals with pain²²⁶ metabolic activation pattern in depression²²⁷ and dopamine release in Parkinson's disease,²²⁸ the data suggest that individual variations in self-confidence, expectation and attitude towards life events, in addition to genetic factors, may help in explaining the larger inter-individual variations in cognitive performance with increasing age.²¹⁸ Remarkable achievements in the graphic arts, architecture and music have often been accomplished in later life,^{229,230} and as mentioned above, the cognitive variance between individuals at high age is large. Creativity may vary with the specific field evaluated²³¹ or change rather than decline with age.²³² In an extensive sample of more than 1,900 works by 172 classical composers, Simonton compared their last works, "swan songs", with their earlier work as to melodic originality, melodic variation, repertoire popularity, aesthetic significance, listener accessibility, performance duration, and thematic size. The composers' swan songs tended to score lower in melodic originality and performance but higher in repertoire popularity and aesthetic significance.²³³

There are many anecdotal reports that performing musicians and conductors live long and active lives. Legendary conductors like Arturo Toscanini, Leopold Stokowski and Arthur Fiedler have had long and productive careers that has led to the proposal that symphony conductors are a remarkably long-lived group. In a 20-year follow study of 437 active and former conductors of major regional and community symphony and opera orchestras in the United States, published in

the Statistical Bulletin in the US²³⁴ the over all mortality was 38% lower than in the general population with the largest difference (56%) in the 50–59 age range and a 34% reduction in the ages 80 and over. However, it was pointed out that the results on conductors parallel the results of a study demonstrating that the mortality among top corporate executives was much more favorable than the mortality among business executive at all levels of accomplishment in the USA. The conclusion was that the data support the theory that work fulfillment and worldwide recognition of professional accomplishments are important determinants of health and longevity. One personal experience from Japan is that I was present when Asahina Takashi, then 91, conducted the Japan New Philharmonic orchestra in the Sumida Triphony Hall in Tokyo in 1999, an excellent performance that got very long and enthusiastic ovations.

Like conductors, many legendary pianists have performed high up in the ages. The degree of skill maintenance for older expert pianists is predicted by the amount of practice.²³⁵ In a study comparing older expert and amateur pianists both groups showed the normal pattern of age-related reductions in standard measures for general processing speed. Whereas performance on music-related tasks showed similar age-related decline for amateur pianists, the average performance level for the expert pianists was only slightly below that of young expert pianists.²³⁵ When pianists of a wide experience and age range were tested on measures of musical memory and musical perceptual speed, the results indicated that high levels of experience in the older participants partially attenuated the negative effects of age on the memory and speed tasks.²³⁶

Studies on longevity in the population in Okinawa are in agreement with the concept that genetic factors interact with life style, self-confidence and social networks, as well as with diet.^{237,238} The Japanese twins, Kin san and Gin san, who if I am correctly informed became famous after singing in the Japanese TV at the age of 100, were seen in international newspapers at the age of 107 when Gin-san, apparently the most active of the two that lived a little longer, was planting a tree while the sister was looking on sitting in a wheelchair. That is to me an excellent example that old individuals can still enjoy life and that in spite of the same genes there might be some differences that can be influenced by attitude and activities.

Concluding Remarks

Our brain is continuously changing and it is important what we do and what we think. A healthy brain can form new dendrites, spines and synapses throughout

our lifetime even if we need to activate more neural circuits for specific tasks than when we were young. Brain damage is still and will remain a serious problem. However, current basic and clinical data indicate that the possibilities to stimulate brain plasticity after a lesion are larger than earlier believed, and the progress made in neurorehabilitation is promising. The concept on brain plasticity has wide implication also for areas outside neuroscience such as hand surgery.^{132,133} For the first time it seems that we have ways to successfully treat focal hand dystonia.^{112–115} We cannot prevent aging but the way we live out lives may influence the aging process. That the environment and our activities influence our lives is basic human knowledge; what is new is that we are starting to understand a little of how it happens.

References

1. Berlucchi G: The origin of the term plasticity in the neurosciences: Ernesto Lugaro and chemical synaptic transmission. *J Hist Neurosci* 2002; 11: 305–309
2. In: Hebb DO, ed, *The Organization of Behavior: A Neuro-psychological Theory*, New York, John Wiley & Sons, 1949
3. Hebb DO: The effect of early experience on problem solving at maturity. *Am Psychol* 1947; 2: 737–745
4. Bennett EL, Diamond MC, Krech D, Rosenzweig MR: Chemical and anatomical plasticity brain. *Science* 1964; 146: 610–619
5. Rosenzweig MR: Environmental complexity, cerebral change, and behavior. *Am Psychol* 1966; 21: 321–332
6. Volkmar FR, Greenough WT: Rearing complexity affects branching of dendrites in the visual cortex of the rat. *Science* 1972; 176: 1145–1147
7. Globus A, Rosenzweig MR, Bennett EL, Diamond MC: Effects of differential experience on dendritic spine counts in rat cerebral cortex. *J Comp Physiol Psychol* 1973; 82: 175–181
8. In: Kolb B, ed, *Brain Plasticity and Behaviour*, NJ, Lawrence Erlbaum Associates, 1995
9. Johansson BB, Belichenko PV: Neuronal plasticity and dendritic spines: effect of environmental enrichment on intact and postischemic rat brain. *J Cereb Blood Flow Metab.* 2002 Jan; 22(1): 89–96
10. Johansson BB, Belichenko PV: Environmental influence on neuronal and dendritic spine plasticity after permanent focal brain ischemia. In: Bazan NG, Ito U, Marcheselli VL, Kuroiwa T, Klatzo I, eds, *Maturation Phenomenon in Cerebral Ischemia IV*, Berlin Heidelberg, Springer Verlag, 2001; 77–23
11. Calverley RK, Jones DG: Contributions of dendritic spines and perforated synapses to synaptic plasticity. *Brain Res Brain Res Rev* 1990; 15: 215–249
12. Eilers J, Konnerth A: Dendritic signal integration. *Curr Opin Neurobiol* 1997; 7: 385–390
13. Harris KM: Structure, development, and plasticity of dendritic spines. *Curr Opin Neurobiol* 1999; 9: 343–348
14. Svoboda K, Denk W, Kleinfeld D, Tank DW: *In vivo* dendritic calcium dynamics in neocortical pyramidal neurons. *Nature* 1997; 385: 161–165
15. Fischer M, Kaech S, Knutti D, Matus A: Rapid actin-based plasticity in dendritic spines. *Neuron* 1998; 20: 847–854
16. Yuste R, Majewska A, Holthoff K: From form to function: calcium compartmentalization in dendritic spines. *Nat Neurosci* 2000; 3: 653–659

17. Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K: Long-term *in vivo* imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* 2002; 420: 788–794
18. Grutzendler J, Kasthuri N, Gan WB: Long-term dendritic spine stability in the adult cortex. *Nature* 2002; 420: 812–816
19. Kim S, Chiba A: Dendritic guidance. *Trends Neurosci* 2004; 27: 194–202
20. Jenkins WM, Merzenich MM, Ochs MT, Allard T, Guic-Robles E: Functional reorganization of primary somatosensory cortex in adult owl monkeys after behaviorally controlled tactile stimulation. *J Neurophysiol* 1990; 63: 82–104
21. Buonomano DV, Merzenich MM: Cortical plasticity: from synapses to maps. *Annu Rev Neurosci* 1998; 21: 149–186
22. Hickmott PW, Merzenich MM: Local circuit properties underlying cortical reorganization. *J Neurophysiol* 2002; 88: 1288–1301
23. Feldman DE, Nicoll RA, Malenka RC: Synaptic plasticity at thalamocortical synapses in developing rat somatosensory cortex: LTP, LTD, and silent synapses. *J Neurobiol* 1999; 41: 92–101
24. Luscher C, Nicoll RA, Malenka RC, Muller D: Synaptic plasticity and dynamic modulation of the postsynaptic membrane. *Nat Neurosci* 2000; 3: 545–550
25. Malenka RC: The long-term potential of LTP. *Nat Rev Neurosci* 2003; 4: 923–926
26. Frick A, Magee J, Johnston D: LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites. *Nat Neurosci* 2004; 7: 126–135
27. Monfils MH, VandenBerg PM, Kleim JA, Teskey GC: Long-term potentiation induces expanded movement representations and dendritic hypertrophy in layer V of rat sensorimotor neocortex. *Cereb Cortex* 2004; 14: 586–593
28. Hess G, Aizenman CD, Donoghue JP: Conditions for the induction of long-term potentiation in layer II/III horizontal connections of the rat motor cortex. *J Neurophysiol* 1996; 75: 1765–1778
29. Garraghty PE, Muja N: NMDA receptors and plasticity in adult primate somatosensory cortex. *J Comp Neurol* 1996; 367: 319–326
30. Ju W, Morishita W, Tsui J, Gaietta G, Deerinck TJ, Adams SR, Garner CC, Tsien RY, Ellisman MH, Malenka RC: Activity-dependent regulation of dendritic synthesis and trafficking of AMPA receptors. *Nat Neurosci* 2004; 7: 244–253
31. Kilgard MP, Merzenich MM: Cortical map reorganization enabled by nucleus basalis activity. *Science* 1998; 279: 1714–1718
32. Gu Q: Neuromodulatory transmitter systems in the cortex and their role in cortical plasticity. *Neuroscience* 2002; 111: 815–835
33. Bao S, Chan VT, Merzenich MM: Cortical remodelling induced by activity of ventral tegmental dopamine neurons. *Nature* 2001; 412: 79–83
34. Schultz W: Neural coding of basic reward terms of animal learning theory, game theory, microeconomics and behavioural ecology. *Curr Opin Neurobiol* 2004; 14: 139–147
35. Kara P, Friedlander MJ: Dynamic modulation of cerebral cortex synaptic function by nitric oxide. *Prog Brain Res* 1998; 118: 183–198
36. Centonze D, Gubellini P, Pisani A, Bernardi G, Calabresi P: Dopamine, acetylcholine and nitric oxide systems interact to induce corticostriatal synaptic plasticity. *Rev Neurosci* 2003; 14: 207–216
37. Castro-Alamancos MA, Donoghue JP, Connors BW: Different forms of synaptic plasticity in somatosensory and motor areas of the neocortex. *J Neurosci* 1995; 15: 5324–5333
38. Remple MS, Bruneau RM, VandenBerg PM, Goertzen C, Kleim JA: Sensitivity of cortical movement representations to motor experience: evidence that skill learning but not strength training induces cortical reorganization. *Behav Brain Res* 2001; 123: 133–141
39. Plautz EJ, Milliken GW, Nudo RJ: Effects of repetitive motor training on movement representations in adult squirrel monkeys: role of use versus learning. *Neurobiol Learn Mem* 2000; 74: 27–55
40. Kleim JA, Cooper NR, VandenBerg PM: Exercise induces angiogenesis but does not alter movement representations within rat motor cortex. *Brain Res* 2002; 934: 1–6
41. Kleim JA, Hogg TM, VandenBerg PM, Cooper NR, Bruneau R, Remple M: Cortical synaptogenesis and motor map reorganization occur during late, but not early, phase of motor skill learning. *J Neurosci* 2004; 24: 628–633
42. Vernadakis A: Glia-neuron intercommunications and synaptic plasticity. *Prog Neurobiol* 1996; 49: 185–214
43. Chvatal A, Sykova E: Glial influence on neuronal signaling. *Prog Brain Res* 2000; 125: 199–216
44. Araque A, Carmignoto G, Haydon PG: Dynamic signaling between astrocytes and neurons. *Annu Rev Physiol* 2001; 63: 795–813
45. Ullian EM, Sapperstein SK, Christopherson KS, Barres BA: Control of synapse number by glia. *Science* 2001; 291: 657–661
46. Nishiyama H, Knopfel T, Endo S, Itoharu S: Glial protein S100B modulates long-term neuronal synaptic plasticity. *Proc Natl Acad Sci USA* 2002; 99: 4037–4042
47. Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA, Gage FH: Neurogenesis in the adult human hippocampus. *Nat Med* 1998; 4: 1313–1317
48. Gage FH: Mammalian neural stem cells. *Science* 2000; 287: 1433–1438
49. Kempermann G, Kuhn HG, Gage FH: More hippocampal neurons in adult mice living in an enriched environment. *Nature* 1997; 386: 493–495
50. van Praag H, Kempermann G, Gage FH: Neural consequences of environmental enrichment. *Nat Rev Neurosci* 2000; 1: 191–198
51. Nilsson M, Perfilieva E, Johansson U, Orwar O, Eriksson PS: Enriched environment increases neurogenesis in the adult rat dentate gyrus and improves spatial memory. *J Neurobiol* 1999; 39: 569–578
52. Schwab ME: Myelin-associated inhibitors of neurite growth and regeneration in the CNS. *Trends Neurosci* 1990; 13: 452–456
53. Chen MS, Huber AB, van der Haar ME, Frank M, Schnell L, Spillmann AA, Christ F, Schwab ME: Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature* 2000; 403: 434–439
54. Watkins TA, Barres BA: Nerve regeneration: regrowth stumped by shared receptor. *Curr Biol* 2002; 12: R654–656
55. Papadopoulos CM, Tsai SY, Alsbie T, O'Brien TE, Schwab ME, Kartje GL: Functional recovery and neuroanatomical plasticity following middle cerebral artery occlusion and IN-1 antibody treatment in the adult rat. *Ann Neurol* 2002; 51: 433–441
56. Wiessner C, Bareyre FM, Allegrini PR, Mir AK, Frentzel S, Zurini M, Schnell L, Oertle T, Schwab ME: Anti-Nogo-A antibody infusion 24 hours after experimental stroke improved behavioral outcome and corticospinal plasticity in normotensive and spontaneously hypertensive rats. *J Cereb Blood Flow Metab* 2003; 23: 154–165
57. GrandPre T, Li S, Strittmatter SM: Nogo-66 receptor antagonist peptide promotes axonal regeneration. *Nature* 2002; 417: 547–551
58. Benowitz LI, Goldberg DE, Madsen JR, Soni D, Irwin N: Inosine stimulates extensive axon collateral growth in the rat cor-

- ticospinal tract after injury. *Proc Natl Acad Sci USA* 1999; 96: 13486–13490
59. Chen P, Goldberg DE, Kolb B, Lanser M, Benowitz LI: Inosine induces axonal rewiring and improves behavioral outcome after stroke. *Proc Natl Acad Sci USA* 2002; 99: 9031–9036
 60. Hari R, Levanen S, Raij T: Timing of human cortical functions during cognition: role of MEG. *Trends Cogn Sci* 2000; 4: 455–462
 61. Ward NS, Frackowiak RS: Towards a new mapping of brain cortex function. *Cerebrovasc Dis* 2004; 17: 35–38
 62. Matthews PM, Jezzard P: Functional magnetic resonance imaging. *J Neurol Neurosurg Psychiatry* 2004; 75: 6–12
 63. Ashburner J, Friston KJ: Voxel-based morphometry – the methods. *Neuroimage* 2000; 11: 805–821
 64. Hallett M: Transcranial magnetic stimulation and the human brain. *Nature* 2000; 406: 147–150
 65. Rossi S, Rossini PM: TMS in cognitive plasticity and the potential for rehabilitation. *Trends Cogn Sci* 2004; 8: 273–279
 66. Butefisch CM: Plasticity in the human cerebral cortex: lessons from the normal brain and from stroke. *Neuroscientist* 2004; 10: 163–173
 67. Di Lazzaro V, Oliviero A, Profice P, Insola A, Mazzone P, Tonali P, Rothwell JC: Direct demonstration of interhemispheric inhibition of the human motor cortex produced by transcranial magnetic stimulation. *Exp Brain Res* 1999; 124: 520–524
 68. Werhahn KJ, Mortensen J, Van Boven RW, Zeuner KE, Cohen LG: Enhanced tactile spatial acuity and cortical processing during acute hand deafferentation. *Nat Neurosci* 2002; 5: 936–938
 69. Bjorkman A, Rosen B, Westén DV, Larsson EM, Lundborg G: Acute improvement of contralateral hand function after deafferentation. *Neuroreport* 2004; 15: 1861–1865
 70. Boroojerdi B, Phipps M, Kopylev L, Wharton CM, Cohen LG, Grafman J: Enhancing analogic reasoning with rTMS over the left prefrontal cortex. *Neurology* 2001; 56: 526–528
 71. Butefisch CM, Khurana V, Kopylev L, Cohen LG: Enhancing encoding of a motor memory in the primary motor cortex by cortical stimulation. *J Neurophysiol* 2004; 91: 2110–2116
 72. Bavelier D, Neville HJ: Cross-modal plasticity: where and how? *Nat Rev Neurosci* 2002; 3: 443–452
 73. Neville H, Bavelier D: Human brain plasticity: evidence from sensory deprivation and altered language experience. *Prog Brain Res* 2002; 138: 177–188
 74. Roder B, Stock O, Bien S, Neville H, Rosler F: Speech processing activates visual cortex in congenitally blind humans. *Eur J Neurosci* 2002; 16: 930–936
 75. Weeks R, Horwitz B, Aziz-Sultan A, Tian B, Wessinger CM, Cohen LG, Hallett M, Rauschecker JP: A positron emission tomographic study of auditory localization in the congenitally blind. *J Neurosci* 2000; 20: 2664–2672
 76. Lessard N, Pare M, Lepore F, Lassonde M: Early-blind human subjects localize sound sources better than sighted subjects. *Nature* 1998; 395: 278–280
 77. Bavelier D, Brozinsky C, Tomann A, Mitchell T, Neville H, Liu G: Impact of early deafness and early exposure to sign language on the cerebral organization for motion processing. *J Neurosci* 2001; 21: 8931–8942
 78. Pascual-Leone A, Torres F: Plasticity of the sensorimotor cortex representation of the reading finger in Braille readers. *Brain* 1993; 116: 39–52
 79. Pascual-Leone A, Wassermann EM, Sadato N, Hallett M: The role of reading activity on the modulation of motor cortical outputs to the reading hand in Braille readers. *Ann Neurol* 1995; 38: 910–915
 80. Goldreich D, Kanics IM: Tactile acuity is enhanced in blindness. *J Neurosci* 2003; 23: 3439–3445
 81. Valaki CE, Maestu F, Simos PG, Zhang W, Fernandez A, Amo CM, Ortiz TM, Papanicolaou AC: Cortical organization for receptive language functions in Chinese, English, and Spanish: a cross-linguistic MEG study. *Neuropsychologia* 2004; 42: 967–979
 82. Schlaug G: The brain of musicians. A model for functional and structural adaptation. *Ann NY Acad Sci* 2001; 930: 281–299
 83. Pascual-Leone A: The brain that plays music and is changed by it. *Ann NY Acad Sci* 2001; 930: 315–329
 84. Munte TF, Altenmüller E, Jancke L: The musician's brain as a model of neuroplasticity. *Nat Rev Neurosci* 2002; 3: 473–478
 85. Schlaug G, Jancke L, Huang Y, Staiger JF, Steinmetz H: Increased corpus callosum size in musicians. *Neuropsychologia* 1995; 33: 1047–1055
 86. Schmithorst VJ, Wilke M: Differences in white matter architecture between musicians and non-musicians: a diffusion tensor imaging study. *Neurosci Lett* 2002; 321: 57–60
 87. Elbert T, Pantev C, Wienbruch C, Rockstroh B, Taub E: Increased cortical representation of the fingers of the left hand in string players. *Science* 1995; 270: 305–307
 88. Hashimoto I, Suzuki A, Kimura T, Iguchi Y, Tanosaki M, Takino R, Haruta Y, Taira M: Is there training-dependent reorganization of digit representations in area 3b of string players? *Clin Neurophysiol* 2004; 115: 435–447
 89. Pantev C, Oostenveld R, Engelien A, Ross B, Roberts LE, Hoke M: Increased auditory cortical representation in musicians. *Nature* 1998; 392: 811–814
 90. Pantev C, Roberts LE, Schulz M, Engelien A, Ross B: Timbre-specific enhancement of auditory cortical representations in musicians. *Neuroreport* 2001; 12: 169–174
 91. Ohnishi T, Matsuda H, Asada T, Aruga M, Hirakata M, Nishikawa M, Katoh A, Imabayashi E: Functional anatomy of musical perception in musicians. *Cereb Cortex* 2001; 11: 754–760
 92. Pascual-Leone A, Grafman J, Hallett M: Modulation of cortical motor output maps during development of implicit and explicit knowledge. *Science* 1994; 263: 1287–1289
 93. Petersen SE, van Mier H, Fiez JA, Raichle ME: The effects of practice on the functional anatomy of task performance. *Proc Natl Acad Sci USA* 1998; 95: 853–860
 94. Pascual-Leone A, Nguyet D, Cohen LG, Brasil-Neto JP, Cammarota A, Hallett M: Modulation of muscle responses evoked by transcranial magnetic stimulation during the acquisition of new fine motor skills. *J Neurophysiol* 1995; 74: 1037–1045
 95. Maguire EA, Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RS, Frith CD: Navigation-related structural change in the hippocampus of taxi drivers. *Proc Natl Acad Sci USA* 2000; 97: 4398–4403
 96. Maguire EA, Spiers HJ, Good CD, Hartley T, Frackowiak RS, Burgess N: Navigation expertise and the human hippocampus: a structural brain imaging analysis. *Hippocampus* 2003; 13: 250–259
 97. Carter RJ, Hunt MJ, Morton AJ: Environmental stimulation increases survival in mice transgenic for exon 1 of the Huntington's disease gene. *Mov Disord* 2000; 15: 925–937
 98. van Dellen A, Blakemore C, Deacon R, York D, Hannan AJ: Delaying the onset of Huntington's in mice. *Nature* 2000; 404: 721–722
 99. Hockley E, Cordery PM, Woodman B, Mahal A, van Dellen A, Blakemore C, Lewis CM, Hannan AJ, Bates GP: Environmental enrichment slows disease progression in R6/2 Huntington's disease mice. *Ann Neurol* 2002; 51: 235–242
 100. Glass M, Dragunow M, Faull RL: The pattern of neurodegeneration in Huntington's disease: a comparative study of cannabinoid, dopamine, adenosine and GABA(A) receptor

- alterations in the human basal ganglia in Huntington's disease. *Neuroscience* 2000; 97: 505–519
101. Glass M, van Dellen A, Blakemore C, Hannan AJ, Faull RL: Delayed onset of Huntington's disease in mice in an enriched environment correlates with delayed loss of cannabinoid CB1 receptors. *Neuroscience* 2004; 123: 207–212
 102. Georgiou N, Bradshaw JL, Chiu E, Tudor A, O'Gorman L, Phillips JG: Differential clinical and motor control function in a pair of monozygotic twins with Huntington's disease. *Mov Disord* 1999; 14: 320–325
 103. Anca MH, Gazit E, Loewenthal R, Ostrovsky O, Frydman M, Giladi N: Different phenotypic expression in monozygotic twins with Huntington disease. *Am J Med Genet* 2004; 124A: 89–91
 104. Schuele S, Lederman RJ: Long-term outcome of focal dystonia in string instrumentalists. *Mov Disord* 2004; 19: 43–48
 105. Byl NN, Merzenich MM, Jenkins WM: A primate genesis model of focal dystonia and repetitive strain injury: I. Learning-induced dedifferentiation of the representation of the hand in the primary somatosensory cortex in adult monkeys. *Neurology* 1996; 47: 508–520
 106. Blake DT, Byl NN, Cheung S, Bedenbaugh P, Nagarajan S, Lamb M, Merzenich M: Sensory representation abnormalities that parallel focal hand dystonia in a primate model. *Somatosens Mot Res* 2002; 19: 347–357
 107. Stinear CM, Byblow WD: Impaired modulation of intracortical inhibition in focal hand dystonia. *Cereb Cortex* 2004; 14: 555–561
 108. McKenzie AL, Nagarajan SS, Roberts TP, Merzenich MM, Byl NN: Somatosensory representation of the digits and clinical performance in patients with focal hand dystonia. *Am J Phys Med Rehabil* 2003; 82: 737–749
 109. Butterworth S, Francis S, Kelly E, McGlone F, Bowtell R, Sawle GV: Abnormal cortical sensory activation in dystonia: an fMRI study. *Mov Disord* 2003; 18: 673–682
 110. Lerner A, Shill H, Hanakawa T, Bushara K, Goldfine A, Hallett M: Regional cerebral blood flow correlates of the severity of writer's cramp symptoms. *Neuroimage* 2004; 21: 904–913
 111. Levy LM, Hallett M: Impaired brain GABA in focal dystonia. *Ann Neurol* 2002; 51: 93–101
 112. Zeuner KE, Bara-Jimenez W, Noguchi PS, Goldstein SR, Dambrosia JM, Hallett M: Sensory training for patients with focal hand dystonia. *Ann Neurol* 2002; 51: 593–598
 113. Byl NN, Nagarajan S, McKenzie AL: Effect of sensory discrimination training on structure and function in patients with focal hand dystonia: a case series. *Arch Phys Med Rehabil* 2003; 84: 1505–1514
 114. Zeuner KE, Hallett M: Sensory training as treatment for focal hand dystonia: a 1-year follow-up. *Mov Disord* 2003; 18: 1044–1047
 115. Candia V, Wienbruch C, Elbert T, Rockstroh B, Ray W: Effective behavioral treatment of focal hand dystonia in musicians alters somatosensory cortical organization. *Proc Natl Acad Sci USA* 2003; 100: 7942–7946
 116. Garraux G, Bauer A, Hanakawa T, Wu T, Kansaku K, Hallett M: Changes in brain anatomy in focal hand dystonia. *Ann Neurol* 2004; 55: 736–739
 117. Frucht SJ, Fahn S, Greene PE, O'Brien C, Gelb M, Truong DD, Welsh J, Factor S, Ford B: The natural history of embouchure dystonia. *Mov Disord* 2001; 16: 899–906
 118. Hirata Y, Schulz M, Altenmüller E, Elbert T, Pantev C: Sensory mapping of lip representation in brass musicians with embouchure dystonia. *Neuroreport* 2004; 15: 815–818
 119. Mühlhnickel W, Elbert T, Taub E, Flor H: Reorganization of auditory cortex in tinnitus. *Proc Natl Acad Sci USA* 1998; 95: 10340–10343
 120. Merzenich MM, Jenkins WM: Reorganization of cortical representations of the hand following alterations of skin inputs induced by nerve injury, skin island transfers, and experience. *J Hand Ther* 1993; 6: 89–104
 121. Merzenich MM, Nelson RJ, Stryker MP, Cynader MS, Schoppmann A, Zook JM: Somatosensory cortical map changes following digit amputation in adult monkeys. *J Comp Neurol* 1984; 224: 591–605
 122. Pons TP, Garraghty PE, Ommaya AK, Kaas JH, Taub E, Mishkin M: Massive cortical reorganization after sensory deafferentation in adult macaques. *Science* 1991; 252: 1857–1860
 123. Pascual-Leone A, Peris M, Tormos JM, Pascual AP, Catala MD: Reorganization of human cortical motor output maps following traumatic forearm amputation. *Neuroreport* 1996; 7: 2068–2070
 124. Faggin BM, Nguyen KT, Nicoletis MA: Immediate and simultaneous sensory reorganization at cortical and subcortical levels of the somatosensory system. *Proc Natl Acad Sci USA* 1997; 94: 9428–9433
 125. Florence SL, Kaas JH: Large-scale reorganization at multiple levels of the somatosensory pathway follows therapeutic amputation of the hand in monkeys. *J Neurosci* 1995; 15: 8083–8095
 126. Wu CW, Kaas JH: The effects of long-standing limb loss on anatomical reorganization of the somatosensory afferents in the brainstem and spinal cord. *Somatosens Mot Res* 2002; 19: 153–163
 127. Wu CW, Kaas JH: Reorganization in primary motor cortex of primates with long-standing therapeutic amputations. *J Neurosci* 1999; 19: 7679–7697
 128. Grusser SM, Mühlhnickel W, Schaefer M, Villringer K, Christmann C, Koeppe C, Flor H: Remote activation of referred phantom sensation and cortical reorganization in human upper extremity amputees. *Exp Brain Res* 2004; 154: 97–102
 129. Flor H, Elbert T, Knecht S, Wienbruch C, Pantev C, Birbaumer N, Larbig W, Taub E: Phantom-limb pain as a perceptual correlate of cortical reorganization following arm amputation. *Nature* 1995; 375: 482–484
 130. Flor H: Phantom-limb pain: characteristics, causes, and treatment. *Lancet Neurol* 2002; 1: 182–189
 131. Topper R, Foltys H, Meister IG, Sparing R, Boroojerdi B: Repetitive transcranial magnetic stimulation of the parietal cortex transiently ameliorates phantom limb pain-like syndrome. *Clin Neurophysiol* 2003; 114: 1521–1130
 132. Lundborg G.: Richard P. Bunge memorial lecture. Nerve injury and repair – a challenge to the plastic brain. *J Peripher Nerv Syst* 2003; 8: 209–226
 133. Rosen B, Lundborg G: Sensory re-education after nerve repair: aspects of timing. *Handchir Mikrochir Plast Chir* 2004; 36: 8–12
 134. Brodal A: Self-observations and neuro-anatomical considerations after a stroke. *Brain* 1973; 96: 675–694
 135. Witte OW: Lesion-induced plasticity as a potential mechanism for recovery and rehabilitative training. *Curr Opin Neurol* 1998; 11: 655–662
 136. Nudo RJ: Recovery after damage to motor cortical areas. *Curr Opin Neurobiol* 1999; 9: 740–747
 137. Johansson BB: Brain plasticity and stroke rehabilitation. The Willis lecture. *Stroke* 2000; 31: 223–230
 138. Hallett M: Plasticity of the human motor cortex and recovery from stroke. *Brain Res Brain Res Rev* 2001; 36: 169–174
 139. Chen R, Cohen LG, Hallett M: Nervous system reorganization following injury. *Neuroscience* 2002; 111: 761–773
 140. Kolb B: Overview of cortical plasticity and recovery from brain injury. *Phys Med Rehabil Clin N Am* 2003; 14(1 Suppl): S7–25
 141. Keyvani K, Schallert T: Plasticity-associated molecular and structural events in the injured brain. *J Neuropathol Exp Neurol* 2002; 61: 831–840

142. Carmichael ST: Plasticity of cortical projections after stroke. *Neuroscientist* 2003; 9: 64–75
143. Jenkins WM, Merzenich MM: Reorganization of neocortical representations after brain injury: a neurophysiological model of the bases of recovery from stroke. *Prog Brain Res* 1987; 71: 249–266
144. Nudo RJ, Wise BM, SiFuentes F, Milliken GW: Neural substrates for the effects of rehabilitative training on motor recovery after ischemic infarct. *Science* 1996; 272: 1791–1794
145. Xerri C, Merzenich MM, Peterson BE, Jenkins W: Plasticity of primary somatosensory cortex paralleling sensorimotor skill recovery from stroke in adult monkeys. *J Neurophysiol* 1998; 79: 2119–2148
146. Kleim JA, Barbay S, Cooper NR, Hogg TM, Reidel CN, Rempel MS, Nudo RJ: Motor learning-dependent synaptogenesis is localized to functionally reorganized motor cortex. *Neurobiol Learn Mem* 2002; 77: 63–77
147. Frost SB, Barbay S, Friel KM, Plautz EJ, Nudo RJ: Reorganization of remote cortical regions after ischemic brain injury: a potential substrate for stroke recovery. *J Neurophysiol* 2003; 89: 3205–3214
148. Will B, Kelche C: Environmental approaches to recovery of function from brain damage: a review of animal studies (1981 to 1991). *Adv Exp Med Biol* 1992; 325: 79–103
149. Will B, Galani R, Kelche C, Rosenzweig MR: Recovery from brain injury in animals: relative efficacy of environmental enrichment, physical exercise or formal training (1990–2002). *Prog Neurobiol* 2004; 72: 167–182
150. Johansson BB: Environmental influence on recovery after brain lesions – experimental and clinical data. *J Rehabil Med* 2003; 41: 11–16
151. Johansson BB: Functional and cellular effects of environmental enrichment after experimental brain infarcts. *Rest Neurol Neurosci* 2004; 22: 163–174
152. Ohlsson AL, Johansson BB: Environment influences functional outcome of cerebral infarction in rats. *Stroke* 1995; 26: 644–649
153. Johansson BB, Ohlsson AL: Environment, social interaction, and physical activity as determinants of functional outcome after cerebral infarction in the rat. *Exp Neurol* 1996; 139: 322–327
154. Zhao LR, Mattsson B, Johansson BB: Environmental influence on brain-derived neurotrophic factor messenger RNA expression after middle cerebral artery occlusion in spontaneously hypertensive rats. *Neuroscience* 2000; 97: 177–184
155. Dahlqvist P, Ronnback A, Risedal A, Nergårdh R, Johansson IM, Seckl JR, Johansson BB, Olsson T: Effects of postischemic environment on transcription factor and serotonin receptor expression after permanent focal cortical ischemia in rats. *Neuroscience* 2003; 119: 643–652
156. Komitova M, Perfilieva E, Mattsson B, Eriksson PS, Johansson BB: Effects of cortical ischemia and postischemic environmental enrichment on hippocampal cell genesis and differentiation in the adult rat. *J Cereb Blood Flow Metab* 2002; 22: 852–860
157. Biernaskie J, Corbett D: Enriched rehabilitative training promotes improved forelimb motor function and enhanced dendritic growth after focal ischemic injury. *J Neurosci* 2001; 21: 5272–5280
158. Puurunen K, Jolkkonen J, Sirvio J, Haapalinna A, Sivenius J: Selegiline combined with enriched-environment housing attenuates spatial learning deficits following focal cerebral ischemia in rats. *Exp Neurol* 2001; 167: 348–355
159. Mattsson B, Sorensen JC, Zimmer J, Johansson BB: Neural grafting to experimental neocortical infarcts improves behavioral outcome and reduces thalamic atrophy in rats housed in enriched but not in standard environments. *Stroke* 1997; 28: 1225–1231
160. Zeng J, Mattsson B, Schulz MK, Johansson BB, Sorensen JC: Expression of zinc-positive cells and terminals in fetal neocortical homografts to adult rat depends on lesion type and rearing conditions. *Exp Neurol* 2000; 164: 176–183
161. Dobrossy MD, Dunnett SB: The influence of environment and experience on neural grafts. *Nat Rev Neurosci* 2001; 2: 871–879
162. Ward NS, Brown MM, Thompson AJ, Frackowiak RS: The influence of time after stroke on brain activations during a motor task. *Ann Neurol* 2004; 55: 829–834
163. Chollet F, DiPiero V, Wise RJ, Brooks DJ, Dolan RJ, Frackowiak RS: The functional anatomy of motor recovery after stroke in humans: a study with positron emission tomography. *Ann Neurol* 1991; 29: 63–71
164. Weiller C, Ramsay SC, Wise RJ, Friston KJ, Frackowiak RS: Individual patterns of functional reorganization in the human cerebral cortex after capsular infarction. *Ann Neurol* 1993; 33: 181–189
165. Carey JR, Kimberley TJ, Lewis SM, Auerbach EJ, Dorsey L, Rundquist P, Ugurbil K: Analysis of fMRI and finger tracking training in subjects with chronic stroke. *Brain* 2002; 125: 773–788
166. Ward NS, Brown MM, Thompson AJ, Frackowiak RS: Neural correlates of outcome after stroke: a cross-sectional fMRI study. *Brain* 2003; 126: 1430–1448
167. Ward NS, Brown MM, Thompson AJ, Frackowiak RS: Neural correlates of motor recovery after stroke: a longitudinal fMRI study. *Brain* 2003; 126: 2476–2496
168. Johansen-Berg H, Dawes H, Guy C, Smith SM, Wade DT, Matthews PM: Correlation between motor improvements and altered fMRI activity after rehabilitative therapy. *Brain* 2002; 125: 2731–2742
169. Johansen-Berg H, Rushworth MF, Bogdanovic MD, Kischka U, Wimalaratna S, Matthews PM: The role of ipsilateral premotor cortex in hand movement after stroke. *Proc Natl Acad Sci USA* 2002; 99: 14518–14523
170. Werhahn KJ, Conforto AB, Kadom N, Hallett M, Cohen LG: Contribution of the ipsilateral motor cortex to recovery after chronic stroke. *Ann Neurol* 2003; 54: 464–472
171. Fridman EA, Hanakawa T, Chung M, Hummel F, Leiguarda RC, Cohen LG: Reorganization of the human ipsilesional premotor cortex after stroke. *Brain* 2004; 127: 747–758
172. Fujii Y, Nakada T: Cortical reorganization in patients with subcortical hemiparesis: neural mechanisms of functional recovery and prognostic implication. *J Neurosurg* 2003; 98: 64–73
173. Murase N, Duque J, Mazzocchio R, Cohen LG: Influence of interhemispheric interactions on motor function in chronic stroke. *Ann Neurol* 2004; 55: 400–409
174. Taub E, Uswatte G, Pidikiti R: Constraint-Induced Movement Therapy: a new family of techniques with broad application to physical rehabilitation – a clinical review. *J Rehabil Res Dev* 1999; 36: 237–251
175. Taub E, Uswatte G, Elbert T: New treatments in neuro-rehabilitation founded on basic research. *Nat Rev Neurosci* 2002; 3: 228–236
176. Tower SS: Pyramidal lesions in the monkey. *Brain* 1940; 63: 36–90
177. Liepert J, Bauder H, Wolfgang HR, Miltner WH, Taub E, Weiller C: Treatment-induced cortical reorganization after stroke in humans. *Stroke* 2000; 31: 1210–1216
178. Wittenberg GF, Chen R, Ishii K, Bushara KO, Eckloff S, Croarkin E, Taub E, Gerber LH, Hallett M, Cohen LG: Constraint-induced therapy in stroke: magnetic-stimulation motor maps and cerebral activation. *Neurorehabil Neural Repair* 2003; 17: 48–57
179. Wolf SL, Blanton S, Baer H, Breshears J, Butler AJ: Repetitive

- task practice: a critical review of constraint-induced movement therapy in stroke. *Neurologist* 2002; 8: 325–338
180. Winstein CJ, Miller JP, Blanton S, Taub E, Uswatte G, Morris D, Nichols D, Wolf S: Methods for a multisite randomized trial to investigate the effect of constraint-induced movement therapy in improving upper extremity function among adults recovering from a cerebrovascular stroke. *Neurorehabil Neural Repair* 2003; 17: 137–152
 181. Lum P, Reinkensmeyer D, Mahoney R, Rymer WZ, Burgar C: Robotic devices for movement therapy after stroke: current status and challenges to clinical acceptance. *Top Stroke Rehabil* 2002; 8: 40–53
 182. Lum PS, Burgar CG, Shor PC, Majmundar M, Van der Loos M: Robot-assisted movement training compared with conventional therapy techniques for the rehabilitation of upper-limb motor function after stroke. *Arch Phys Med Rehabil* 2002; 83: 952–959
 183. Ferraro M, Palazzolo JJ, Krol J, Krebs HI, Hogan N, Volpe BT: Robot-aided sensorimotor arm training improves outcome in patients with chronic stroke. *Neurology* 2003; 61: 1604–1607
 184. Fasoli SE, Krebs HI, Stein J, Frontera WR, Hogan N: Effects of robotic therapy on motor impairment and recovery in chronic stroke. *Arch Phys Med Rehabil* 2003; 84: 477–482
 185. Hesse S, Schmidt H, Werner C, Bardeleben A: Upper and lower extremity robotic devices for rehabilitation and for studying motor control. *Curr Opin Neurol* 2003; 16: 705–710
 186. Altschuler EL, Wisdom SB, Stone L, Foster C, Galasko D, Llewellyn DM, Ramachandran VS: Rehabilitation of hemiparesis after stroke with a mirror. *Lancet* 1999; 353: 2035–2036
 187. Sathian K, Greenspan AI, Wolf SL: Doing it with mirrors: a case study of a novel approach to neurorehabilitation. *Neurorehabil Neural Repair* 2000; 14: 73–76
 188. Rizzolatti G, Luppino G: The cortical motor system. *Neuron* 2001; 31: 889–901
 189. Rizzolatti G, Fogassi L, Gallese V: Neurophysiological mechanisms underlying the understanding and imitation of action. *Nat Rev Neurosci* 2001; 2: 661–670
 190. Buccino G, Binkofski F, Riggio L: The mirror neuron system and action recognition. *Brain Lang* 2004; 89: 370–376
 191. Merians AS, Jack D, Boian R, Tremaine M, Burdea GC, Adamovich SV, Recce M, Poizner H: Virtual reality-augmented rehabilitation for patients following stroke. *Phys Ther* 2002; 82: 898–915
 192. Page SJ, Levine P, Sisto SA, Johnston MV: Mental practice combined with physical practice for upper-limb motor deficit in subacute stroke. *Phys Ther* 2001; 81: 1455–1462
 193. Deiber MP, Ibanez V, Honda M, Sadato N, Raman R, Hallett M: Cerebral processes related to visuomotor imagery and generation of simple finger movements studied with positron emission tomography. *Neuroimage* 1998; 7: 73–85
 194. Malouin F, Belleville S, Richards CL, Desrosiers J, Doyon J: Working memory and mental practice outcomes after stroke. *Arch Phys Med Rehabil* 2004; 85: 177–183
 195. Muellbacher W, Richards C, Ziemann U, Wittenberg G, Wetz D, Boroojerdi B, Cohen L, Hallett M: Improving hand function in chronic stroke. *Arch Neurol* 2002; 59: 1278–1282
 196. Sharp FR, Liu J, Bernabeu R: Neurogenesis following brain ischemia. *Brain Res Dev Brain Res* 2002; 134: 23–30
 197. Parent JM: Injury-induced neurogenesis in the adult mammalian brain. *Neuroscientist* 2003; 9: 261–272
 198. Kokaia Z, Lindvall O: Neurogenesis after ischaemic brain insults. *Curr Opin Neurobiol* 2003; 13: 127–132
 199. Lie DC, Song H, Colamarino SA, Ming GL, Gage FH: Neurogenesis in the adult brain: new strategies for central nervous system diseases. *Annu Rev Pharmacol Toxicol* 2004; 44: 399–421
 200. Magavi SS, Leavitt BR, Macklis JD: Induction of neurogenesis in the neocortex of adult mice. *Nature* 2000; 405: 951–955
 201. Arlotta P, Magavi SS, Macklis JD: Induction of adult neurogenesis: molecular manipulation of neural precursors in situ. *Ann NY Acad Sci* 2003; 991: 229–236
 202. Nakatomi H, Kuriu T, Okabe S, Yamamoto S, Hatano O, Kawahara N, Tamura A, Kirino T, Nakafuku M: Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors. *Cell* 2002; 110: 429–441
 203. Arvidsson A, Collin T, Kirik D, Kokaia Z, Lindvall O: Neuronal replacement from endogenous precursors in the adult brain after stroke. *Nat Med* 2002; 8: 963–970
 204. Parent JM, Vexler ZS, Gong C, Derugin N, Ferriero DM: Rat forebrain neurogenesis and striatal neuron replacement after focal stroke. *Ann Neurol* 2002; 52: 802–813
 205. Coq JO, Xerri C: Sensorimotor experience modulates age-dependent alterations of the forepaw representation in the rat primary somatosensory cortex. *Neuroscience* 2001; 104: 705–715
 206. Frick KM, Fernandez SM: Enrichment enhances spatial memory and increases synaptophysin levels in aged female mice. *Neurobiol Aging* 2003; 24: 615–626
 207. Kempermann G, Gast D, Gage FH: Neuroplasticity in old age: sustained fivefold induction of hippocampal neurogenesis by long-term environmental enrichment. *Ann Neurol* 2002; 52: 135–143
 208. Bartley AJ, Jones DW, Weinberger DR: Genetic variability of human brain size and cortical gyral patterns. *Brain* 1997; 120: 257–269
 209. Lohmann G, von Cramon DY, Steinmetz H: Sulcal variability of twins. *Cereb Cortex* 1999; 9: 754–763
 210. White T, Andreasen NC, Nopoulos P: Brain volumes and surface morphology in monozygotic twins. *Cereb Cortex* 2002; 12: 486–493
 211. Sullivan EV, Pfefferbaum A, Swan GE, Carmelli D: Heritability of hippocampal size in elderly twin men: equivalent influence from genes and environment. *Hippocampus* 2001; 11: 754–762
 212. Jarvenpää T, Laakso MP, Rossi R, Koskenvuo M, Kaprio J, Raiha I, Kurki T, Laine M, Frisoni GB, Rinne JO: Hippocampal MRI volumetry in cognitively discordant monozygotic twin pairs. *J Neurol Neurosurg Psychiatry* 2004; 75: 116–120
 213. Calautti C, Serrati C, Baron JC: Effects of age on brain activation during auditory-cued thumb-to-index opposition: A positron emission tomography study. *Stroke* 2001; 32: 139–146
 214. Mattay VS, Fera F, Tessitore A, Hariri AR, Das S, Callicott JH, Weinberger DR: Neurophysiological correlates of age-related changes in human motor function. *Neurology* 2002; 58: 630–635
 215. Ward NS, Frackowiak RS: Age-related changes in the neural correlates of motor performance. *Brain* 2003; 126: 873–888
 216. Morcom AM, Good CD, Frackowiak RS, Rugg MD: Age effects on the neural correlates of successful memory encoding. *Brain* 2003; 126: 213–229
 217. Cabeza R, Daselaar SM, Dolcos F, Prince SE, Budde M, Nyberg L: Task-independent and task-specific age effects on brain activity during working memory, visual attention and episodic retrieval. *Cereb Cortex* 2004; 14: 364–375
 218. Rabbitt P, Diggle P, Smith D, Holland F, McInnes L: Identifying and separating the effects of practice and of cognitive ageing during a large longitudinal study of elderly community residents. *Neuropsychologia* 2001; 39: 532–543
 219. Wilson RS, Bennett DA, Bienias JL, Mendes de Leon CF, Morris MC, Evans DA: Cognitive activity and cognitive decline in a biracial community population. *Neurology* 2003; 61: 812–816

220. Wilson RS, Bennett DA, Bienias JL, Aggarwal NT, Mendes De Leon CF, Morris MC, Schneider JA, Evans DA: Cognitive activity and incident AD in a population-based sample of older persons. *Neurology* 2002; 59: 1910–1914
221. Wilson RS, Mendes De Leon CF, Barnes LL, Schneider JA, Bienias JL, Evans DA, Bennett DA: Participation in cognitively stimulating activities and risk of incident Alzheimer disease. *JAMA* 2002; 287: 742–748
222. Ranganathan VK, Siemionow V, Sahgal V, Liu JZ, Yue GH: Skilled finger movement exercise improves hand function. *J Gerontol A Biol Sci Med Sci* 2001; 56: M518–522
223. Valenzuela MJ, Jones M, Wen W, Rae C, Graham S, Shnier R, Sachdev P: Memory training alters hippocampal neurochemistry in healthy elderly. *Neuroreport* 2003; 14: 1333–1337
224. Uotinen V, Suutama T, Ruoppila I: Age identification in the framework of successful aging. A study of older Finnish people. *Int J Aging Hum Dev* 2003; 56: 173–195
225. Wahlin A, Maitland SB, Backman L, Dixon RA: Interrelations between subjective health and episodic memory change in Swedish and Canadian samples of older adults. *Int J Aging Hum Dev* 2003; 57: 21–35
226. Petrovic P, Kalso E, Petersson KM, Ingvar M: Placebo and opioid analgesia – imaging a shared neuronal network. *Science* 2002; 295: 1737–1740
227. Mayberg HS, Silva JA, Brannan SK, Tekell JL, Mahurin RK, McGinnis S, Jerabek PA: The functional neuroanatomy of the placebo effect. *Am J Psychiatry* 2002; 159: 728–737
228. de la Fuente-Fernandez R, Stoessl AJ: The placebo effect in Parkinson's disease. *Trends Neurosci* 2002; 25: 302–306
229. Pritikin R: Marcel Duchamp, the artist, and the social expectations of aging. *Gerontologist* 1990; 30: 636–639
230. Simonton DK: Creativity. Cognitive, personal, developmental, and social aspects. *Am Psychol* 2000; 55: 151–158
231. Hendricks J: Creativity over the life course – a call for a relational perspective. *Int J Aging Hum Dev* 1999; 48: 85–111
232. Abra J: Changes in creativity with age: data, explanations, and further predictions. *Int J Aging Hum Dev* 1989; 28: 105–126
233. Simonton DK: The swan-song phenomenon: last-works effects for 172 classical composers. *Psychol Aging* 1989; 4: 42–47
234. Longevity of symphony conductors. *Stat Bull Metropol Life Insur Co* 1980; 61: 2–4
235. Krampe RT, Ericsson KA: Maintaining excellence: deliberate practice and elite performance in young and older pianists. *J Exp Psychol Gen* 1996; 125: 331–359
236. Meinz EJ: Experience-based attenuation of age-related differences in music cognition tasks. *Psychol Aging* 2000; 15: 297–312
237. Cockerham WC, Hattori H, Yamori Y: The social gradient in life expectancy: the contrary case of Okinawa in Japan. *Soc Sci Med* 2000; 51: 115–122
238. Yamori Y, Miura A, Taira K: Implications from and for food cultures for cardiovascular diseases: Japanese food, particularly Okinawan diets. *Asia Pac J Clin Nutr* 2001; 10: 144–145