REVIEW

Coagulation and fibrinolysis in human acute lung injury – New therapeutic targets?

Lorraine B. Ware, Julie A. Bastarache and Ling Wang

*Division of Allergy, Pulmonary, and Critical Care, Department of Medicine
Vanderbilt University School of Medicine, Nashville, TN, USA*

(Received for publication on May 3, 2005)

(Accepted for publication on July 21, 2005)

Abstract. Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) are common, life-threatening causes of acute respiratory failure that arise from a variety of local and systemic insults. The need for new specific therapies has led a number of investigators to examine the role of altered coagulation and fibrinolysis in the pathogenesis of ALI/ARDS. This review summarizes our current understanding of coagulation and fibrinolysis in human ALI/ARDS with an emphasis on pathways that could be potential therapeutic targets including the tissue factor pathway, the protein C pathway and modulation of fibrinolysis via plasminogen activator inhibitor-1. The available data suggest that clinical ALI and ARDS are characterized by profound alterations in both systemic and intra-alveolar coagulation and fibrinolysis. Fibrin deposition in the airspaces and lung microvasculature likely results from both activation of the coagulation cascade and impaired fibrinolysis, triggered by inflammation. Modulation of fibrin deposition in the lung through targeting activation and modulation of coagulation as well as fibrinolysis may be an important therapeutic target in clinical ALI/ARDS that deserves further exploration. (Keio J Med 54 (3): 142–149, September 2005)

Key words: fibrin deposition, thrombosis, activated protein C, tissue factor, plasminogen activator inhibitor

Introduction

Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) are common, life-threatening causes of acute respiratory failure that arise from a variety of local and systemic insults.¹ Despite some improvement in mortality with a lung-protective ventilator strategy,² both morbidity³ and mortality remain high.⁴ The need for new specific therapies has led a number of investigators to examine the role of altered coagulation and fibrinolysis in the pathogenesis of ALI/ARDS. Although the presence of intra-alveolar fibrin lining the denuded alveolar surface has long been recognized as a pathological hallmark of ALI/ARDS,⁵ the mechanisms that govern fibrin deposition in the acutely injured lung are poorly understood. This review will summarize our current understanding of coagulation and fibrinolysis in human ALI/ARDS with an emphasis on pathways that could be potential therapeutic targets. These pathways are summarized in Fig. 1 and include the tissue

factor pathway, the protein C pathway and modulation of fibrinolysis via plasminogen activator inhibitor-1 (PAI-1).

Tissue Factor and Tissue Factor Pathway Inhibitor

Initiation of coagulation through the extrinsic pathway is likely the predominant mechanism of activation of coagulation in the lung in ALI/ARDS. Tissue factor (TF) is a potent initiator of the extrinsic coagulation cascade. TF is a membrane bound protein that when complexed to factor VIIa converts factor X to Xa leading to the downstream generation of thrombin and conversion of fibrinogen to insoluble fibrin (Fig. 1). In the normal human lung, both alveolar macrophages and alveolar epithelial cells stain positive for TF⁶ and in culture, both alveolar macrophages and alveolar epithelial cells have TF activity.⁷ Initiation of coagulation through the TF pathway is modulated by the balance between TF and tissue factor pathway inhibitor (TFPI),

Fig. 1 Summary of coagulation and fibrinolytic pathways that may be therapeutic targets in clinical ALI/ARDS. A. Activation of coagulation by Tissue Factor (TF). TF initiates the extrinsic pathway of coagulation by acting as a cofactor for the activation of factor VII to VIIa. The factor VIIa-TF complex activates factor X to Xa which catalyzes the conversion of prothrombin to thrombin, leading to fibrin deposition. Tissue factor pathway inhibitor (TFPI) can bind to the TF:VIIa:Xa complex thus inhibiting downstream thrombin formation and fibrin deposition. **B. The protein C pathway.** The protein C pathway is a major endogenous inhibitor of coagulation. Once generated, thrombin binds to the cell surface receptor thrombomodulin (TM), activating the zymogen protein C to the activated protein C (APC). APC (along with protein S) inhibits Xa and thrombin formation through the inactivation of factors V and VIII (not shown). **C. Fibrinolysis.** Fibrinolysis is initiated by urokinase (uPA) and tissue plasminogen activator (tPA) both of which cleave plasminogen from plasminogen. Plasmin degrades fibrin into fibrin degradation products (FDP). Plasminogen activator inhibitor-1 (PAI-1) depresses fibrinolysis by inhibiting uPA and tPA. PAI-1 can be neutralized by APC. The net result of upregulated coagulation and downregulated fibrinolysis in the acutely injured lung is fibrin deposition.

an endogenous inhibitor of TF. TFPI is secreted by the vascular endothelium and circulates in the plasma and bound to the surface of platelets.⁸ TFPI can prevent the activation of X to Xa through binding to the TF:VIIa:Xa complex thereby inhibiting downstream formation of thrombin and deposition of fibrin. Thus, the balance between TF and TFPI levels in the lung may be an important determinant of intra-alveolar fibrin deposition.

An important role for TF in initiating intra-alveolar coagulation in clinical ALI/ARDS was first reported by Idell.⁹ In serial bronchoalveolar lavage (BAL) samples from patients with ARDS, procoagulant activity was upregulated by three days after the onset of ARDS and resolved by seven days.⁹ TF activity was primarily responsible for the procoagulant activity in the BAL since almost all of the procoagulant activity was blocked by an antibody to TF. TF-dependent procoagulant activity in the alveolar compartment has also been reported in patients at risk for ALI/ARDS.¹⁰ Upregulation of TF-dependent procoagulant activity is not completely specific for ALI/ARDS, however. Gunther *et al.* found increases in TF-dependent procoagulant activity in the BAL of patients with pneumonia even in the absence of respiratory failure or ARDS.¹¹ Upregulation of the TF pathway also occurs systemically in clinical ALI/ARDS. Circulating levels of TF were increased in patients with

ARDS but not in patients at risk for ARDS or without ARDS.¹²

Levels of TFPI, the endogenous TF inhibitor, have also been measured in the plasma and BAL of patients with ALI/ARDS. Sabharwal *et al.*¹³ reported a modest increase in levels of TFPI compared to normals in the plasma of patients at risk (1.3-fold) and those with established ARDS (1.8-fold) while Gando *et al.*¹⁴ found no difference in TFPI levels between control, at risk patients or patients with ARDS. By contrast, substantial increases in TFPI were seen in the BAL in those at risk (7-fold) and with established ARDS (20-fold) compared to normals.¹³ TF levels were not measured in that study. Thus, changes in TFPI levels appear to be more pronounced in the alveolar compartment in ALI/ARDS than in the plasma. The effect of the balance of TF and TFPI on overall procoagulant activity has not been studied in ALI/ARDS but has been reported in other lung diseases. The procoagulant activity of BAL was increased in patients with bacterial or *Pneumocystis carinii* pneumonia or interstitial lung disease compared to controls but TFPI levels were similar to controls¹⁵ suggesting that increases in TF activity were not balanced by comparable increases in TFPI. In patients with idiopathic pulmonary fibrosis, both TF and TFPI levels were increased in the BAL and correlated with severity of disease; however, the procoagulant activity

of the BAL was increased compared to controls, again demonstrating that the increases in TF were not balanced by increases in TFPI.¹⁶

The fundamental role of TF in initiation of coagulation combined with the systemic and intra-pulmonary abnormalities of TF and TFPI in clinical ALI/ARDS makes the TF pathway an attractive target for therapeutic intervention. Blockade of TF activity either prior to or in the early stages of ALI with TFPI is protective in several animal models of sepsis and ALI.^{17,18} However, human studies to date have been disappointing. A recent phase III trial of recombinant TFPI in sepsis, the most common cause of ALI/ARDS, had no effect on mortality¹⁹ despite a prior encouraging phase II study.²⁰ A site inactivated form of factor VIIa (factor VIIai) that binds to and inhibits TF activity has been developed as an anticoagulant. In a baboon model of sepsis factor VIIai was protective.^{21–23} A clinical trial in patients with ALI/ARDS is currently underway.

Protein C and Thrombomodulin

The protein C pathway is an important endogenous regulator of coagulation and fibrinolysis. Synthesized by the liver, protein C is a vitamin K-dependent glycoprotein that circulates as an inactive zymogen.²⁴ It is transformed to its active form on the cell surface by the TM-thrombin complex. Another cell surface protein, the endothelial cell protein C receptor (EPCR) can further enhance protein C activation by binding to the TM-thrombin complex.²⁵ Activated protein C proteolytically destroys coagulation factors Va and VIIIa, suppressing further thrombin formation (Fig. 1).^{26,27} Activated protein C has anti-inflammatory and anti-apoptotic effects in addition to its anti-coagulant properties that include suppression of production of pro-inflammatory cytokines, inhibition of leukocyte attachment to the endothelium,^{28,29} and inhibition of p53 mediated apoptosis.³⁰ Activated protein C also can neutralize PAI-1, thereby promoting fibrinolysis.³¹ Thus, modulation of the protein C pathway could have key regulatory effects on coagulation, fibrinolysis and inflammation in ALI/ARDS.

The majority of studies of the protein C pathway have focused on the endothelium as the primary site for activation of protein C. However, recent studies suggest that the lung epithelium can actively modulate the protein C pathway. Cultured human airway epithelial cells express protein C, EPCR and thrombomodulin and protein C was activated by these cells in the presence of thrombin.³² Exposure to inflammatory cytokines downregulated the activation of protein C.³² Alveolar epithelial cells also may modulate the protein C pathway. We reported that both a human alveolar epithelial cell line (A549) and primary isolates of hu-

man alveolar epithelial type II cells release thrombomodulin in response to stimulation by a mixture of pro-inflammatory cytokines or hydrogen peroxide (Fig. 2).³³ Thus, the alveolar epithelium may modulate intra-alveolar coagulation and inflammation through the protein C pathway similar to the well-described role of the vascular endothelium in modulating this pathway.

In humans, most of the clinical studies of the protein C pathway have been in patients with sepsis. In 70 patients with severe sepsis³⁴ plasma protein C levels were markedly reduced compared to normal levels and lower levels were associated with higher mortality, ventilator dependency and a higher prevalence of ARDS. Patients with severe sepsis also vary in their ability to generate activated protein C.³⁵ In a study of 32 patients with severe sepsis, activated protein C levels were higher in survivors although in that study there was no difference in protein C levels between survivors and nonsurvivors. Thrombomodulin levels in the plasma are elevated in patients with sepsis, and high levels correlate with the development of multiple organ failure.^{36,37} Levels of soluble EPCR are also high in patients with sepsis.³⁸ However, there was no correlation between the soluble EPCR levels and clinical outcomes including multiple organ failure score or survival.³⁸

There have been relatively few studies of the protein C pathway in patients with ALI/ARDS. In a single center study, plasma protein C levels were lower in ALI/ARDS compared with normal controls even in the absence of sepsis.³³ Levels of protein C in the alveolar compartment were even lower than plasma levels and lower levels of both plasma and edema fluid protein C were associated with worse clinical outcomes.³³ Like patients with sepsis, patients with ALI/ARDS have high circulating levels of thrombomodulin. In addition, thrombomodulin levels in the alveolar compartment in ALI/ARDS were 2-fold higher than simultaneous plasma samples, suggesting an intra-alveolar source (Fig. 3). High thrombomodulin levels in the plasma or alveolar compartment were associated with worse clinical outcomes.³³ Thus, abnormalities of the protein C pathway in ALI/ARDS are similar to abnormalities in sepsis and are strongly associated with adverse outcomes.

The observation that protein C and activated protein C levels are low in patients with severe sepsis led to a phase III randomized, multicenter clinical trial of recombinant human activated protein C (drotrecogin- α activated) in patients with severe sepsis and at least one organ failure. In this landmark trial, administration of recombinant human activated protein C significantly improved survival compared to placebo. The absolute risk reduction in mortality at 28 days was 6.1%.³⁹ This study was not designed to specifically evaluate the role

Fig. 2 Alveolar epithelial cells release soluble thrombomodulin in response to injurious stimuli. Confluent monolayers of human alveolar epithelial type II-like cells (A549 cells) were exposed for 6 hours to varying concentrations of H₂O₂ or cytomix (equal amounts of TNF- α , IL-1 β , IFN- γ), and conditioned media was collected for assay of thrombomodulin by ELISA. *p < 0.01 compared with 0.01% H₂O₂ or control. **p < 0.01 compared with 10 ng/ml cytomix or control. Reproduced with permission from *American Journal of Physiology Lung Cellular and Molecular Physiology*.³³

of activated protein C in ALI/ARDS. However, a retrospective analysis showed that in the group of patients with respiratory system dysfunction, many of whom likely had ALI/ARDS, the time to resolution of respiratory dysfunction was significantly shorter in the treated group compared to the placebo group.⁴⁰ A multicenter phase II trial of recombinant activated protein C in ALI and ARDS is currently underway. Interestingly, in a human model of pulmonary inflammation produced by instillation of endotoxin into the distal airspaces, intravenous administration of recombinant activated protein C produce sustained elevations in BAL levels of activated protein C⁴¹ that were accompanied by reduced leukocyte accumulation in the airspaces and reductions in neutrophil chemotaxis, independent of pulmonary cytokine or chemokine release.⁴² These findings suggest that in addition to its anticoagulant effects, systemically administered activated protein C may have potent anti-inflammatory effects in the distal airspaces that are mediated through modulation of neutrophil function.

Plasminogen Activators and Inhibitors

The balance between activation of coagulation and activation of fibrinolysis is likely an important determinant of the amount and duration of fibrin deposition in the injured lung. Plasminogen activator (PA) and plasminogen activator inhibitor-1 (PAI-1) regulate fibrinol-

Fig. 3 Boxplot summary of thrombomodulin levels in normal plasma and pulmonary edema fluid from 45 patients with ALI or ARDS. Box encompasses 25th to 75th percentile, error bars encompass 10th to 90th percentile, and horizontal bar shows the median. *p = 0.001 compared with normal plasma or ALI/ARDS plasma. Reproduced with permission from *American Journal of Physiology Lung Cellular and Molecular Physiology*.³³

ysis, the dissolution of fibrin clots, through modulation of the conversion of plasminogen to plasmin, a major fibrinolytic enzyme (Fig. 1). There are two forms of plasminogen activator (PA), urokinase-type plasminogen activator (uPA) and tissue-type plasminogen activator (tPA). uPA is a cell surface protein that is responsible for activating fibrinolysis at the tissue level⁴³ while tPA is a soluble protein that activates intravascular fibrinolysis.⁴⁴ Plasminogen activator inhibitor-1 (PAI-1) is a major endogenous inhibitor of both uPA and tPA.

There are multiple cellular sources of PA and PAI-1 that may be relevant to human ALI/ARDS. Unstimulated alveolar macrophages are pro-fibrinolytic. Primary isolates of human alveolar macrophages have PA activity and degrade a fibrin matrix in the presence of plasminogen.⁴⁵ By contrast, stimulated alveolar macrophages are anti-fibrinolytic. Human alveolar macrophages exposed to endotoxin inhibit fibrinolysis through an increase in PAI-1 activity.^{46,47} Increased PAI-1 activity has also been observed in macrophages isolated from patients with lung disease. For example, alveolar macrophages from patients with idiopathic pulmonary fibrosis (IPF) have increased PAI-1 staining compared to normal controls⁴⁸ and alveolar macrophages from

patients with ARDS have increased PAI-1 mRNA as demonstrated by *in situ* hybridization.⁴⁷ Overall, these findings suggest that when the lung is exposed to injurious stimuli there is a decrease in the fibrinolytic activity of alveolar macrophages through decreased PA activity and increased PAI-1 activity.

Like macrophages, human lung microvascular endothelial cells can secrete both PA⁴⁹ and PAI-1.⁵⁰ Pro-inflammatory stimuli increase PA secretion several-fold^{49,50} but the net effect on fibrinolysis is unclear. In one study of human lung microvascular endothelial cells isolated from patients with ARDS, the cells from ARDS patients constitutively expressed more PAI-1 than controls and had a lower fibrinolytic potential as measured by the PA/PAI-1 ratio.⁵¹ Although the data are not completely concordant, the studies from ARDS patients suggest that similar to alveolar macrophages, the pulmonary microvascular endothelium in its basal state favors fibrinolysis. Injurious stimuli shifts the balance away from fibrinolysis.

The lung epithelium can also express both plasminogen activators and inhibitors and is thus capable of regulating intra-alveolar fibrinolysis. However, there are relatively few studies of human lung epithelial cells. Primary isolates of rat alveolar epithelial cells express both PA and PAI-1; levels increase with time in culture as the cells differentiate from a type II cell phenotype to a type I phenotype.^{52,53} Exposure of the cells to lipopolysaccharide or TNF- α , increased PAI-1 and uPA suggesting that inflammatory mediators modulate the fibrinolytic capacity of alveolar epithelial cells. A human alveolar epithelial type II cell line (A549) had increases in uPA mRNA, protein, and activity following exposure to IL-1 β and TNF- α , but unlike rat alveolar epithelial cells, there was no change in PAI-1 expression.⁵⁴ By contrast, freshly isolated alveolar type I cells do express PAI-1 mRNA.⁵⁵ Thus, PAI-1 expression may be a phenotype of alveolar type I cells and not type II cells, similar to the findings with rat alveolar epithelial cells. Taken together, the *in vitro* data suggests that the lung epithelium is probably capable of modulating intra-alveolar fibrinolysis through expression of PA, and PAI-1.

There are only a few studies that address fibrinolysis in patients with ALI/ARDS. In a study comparing normal controls to four groups of patients (patients with pneumonia who were spontaneously breathing, patients with pneumonia requiring mechanical ventilation, patients with ARDS without pneumonia and patients with ARDS with pneumonia) there was reduced fibrinolytic capacity and an increase in uPA in all groups compared to control as well as an increase in PAI-1 activity in all groups except those with pneumonia who were spontaneously breathing.¹¹ Prabhakaran *et al.* measured PAI-1 antigen levels in plasma and simultaneous undiluted

Fig. 4 Plasma and edema fluid PAI-1 antigen in patients with acute lung injury (ALI) and hydrostatic pulmonary edema (HYDRO). PAI-1 values were measured in simultaneously drawn plasma (open boxes) and edema fluid (filled boxes) by ELISA from patients with clinical ALI or hydrostatic edema. Data are plotted in box plot format (median, 25–75%) on a log scale and compared using Mann-Whitney's *U*-test. P values are as indicated. Reproduced with permission from *American Journal of Physiology Lung Cellular and Molecular Physiology*.⁵⁶

pulmonary edema fluid from patients with ALI/ARDS and compared them to control patients with pulmonary edema due to hydrostatic causes.⁵⁶ Levels of PAI-1 were higher in both the edema fluid and plasma of patients with ALI/ARDS compared to the control patients (Fig. 4) and correlated with mortality in patients with ALI/ARDS. In patients with ALI/ARDS, PAI-1 levels were substantially higher in the edema fluid than the plasma suggesting an intra-alveolar source of PAI-1. In other studies that only examined plasma levels, patients with ARDS and those at risk for ARDS had higher circulating levels of PAI-1 than controls.^{57,58} However, plasma levels did not correlate with lung injury score, pulmonary microvascular permeability, or clinical outcome.⁵⁸ Taken together, these findings suggest that alterations in the intra-alveolar fibrinolytic environment may have more impact on outcomes in ALI/ARDS than alterations in systemic fibrinolytic pathways. Although there are no studies to date in patients with ALI/ARDS, polymorphisms in the promoter region of PAI-1 have been studied in two populations of patients at risk for ALI/ARDS, patients with meningococcal septicemia and patients with severe trauma at risk for ALI/ARDS.⁵⁹ Deletions in the promoter region lead to impaired fibrinolytic ability and are associated with disease severity and outcome in meningococcal disease^{60–62} and with susceptibility to sepsis and multi-organ failure in patients with severe trauma.⁶³

To summarize, there is substantial evidence that the fibrinolytic system is profoundly altered in patients with ALI/ARDS, both systemically and in the alveolar compartment. Upregulation of PAI-1, the major inhibitor of fibrinolysis, appears to play a primary role in the shift from pro-fibrinolytic to anti-fibrinolytic phenotypes in a variety of cell types including endothelium, lung epithelium and alveolar macrophages. A variety of strategies are being explored to develop inhibitors of PAI-1⁶⁴ that might be of therapeutic use in ALI/ARDS or other diseases associated with high levels of PAI-1 such as cardiovascular disease. However, no inhibitor of PAI-1 is currently clinically available.

Conclusions

Clinical ALI and ARDS are characterized by profound alterations in both systemic and intra-alveolar coagulation and fibrinolysis. Fibrin deposition in the airspaces and the lung microvasculature likely results from inflammation-induced activation of the coagulation cascade and impairment of fibrinolysis. Activation of coagulation with resultant fibrin deposition also has proinflammatory consequences, serving to further amplify the inflammatory cascade. Current evidence from human studies suggests that in the acutely injured lung there is activation of tissue factor, downregulation of protein C activation, and increased production of PAI-1. Together these abnormalities shift the intra-alveolar environment from anticoagulant and profibrinolytic to procoagulant and antifibrinolytic. Modulation of fibrin deposition in the lung through targeting coagulation and fibrinolysis may be an important therapeutic target in clinical ALI/ARDS. Because the lung epithelium appears to be an important site for modulation of intra-alveolar fibrin deposition, it represents a unique therapeutic target. Targeting the lung epithelium with inhaled therapies to modulate intra-alveolar fibrin deposition might avoid the deleterious side effects of systemic therapies and deserves further study.

References

1. Ware LB, Matthay MA: Medical progress: The acute respiratory distress syndrome. *N Engl J Med* 2000; 342: 1334–1349
2. The Acute Respiratory Distress Syndrome Network: Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *New Engl J Med* 2000; 342: 1301–1308
3. Herridge MS, Cheung AM, Tansey CM, Matte-Martyn A, Diaz-Granados N, Fatma A-S, Cooper AB, Guest CB, Mazer CD, Mehta S, Stewart TE, Barr A, Cook D, Slutsky AS, for the Canadian Critical Care Trials Group: One-year outcomes in survivors of the acute respiratory distress syndrome. *N Engl J Med* 2003; 348: 683–693
4. Brun-Buisson C, Minelli C, Bertolini G, Brazzi L, Pimentel J, Lewandowski K, Bion J, Romand J-A, Villar J, Thorsteinsson A, Damas P, Armaganidis A, Lemaire F, for the ALIVE Study Group: Epidemiology and outcome of acute lung injury in European intensive care units. *Intensive Care Med* 2004; 30: 51–61
5. Bachofen M, Weibel ER: Structural alterations of lung parenchyma in the adult respiratory distress syndrome. *Clin Chest Med* 1982; 3: 35–56
6. Drake TA, Morrissey JH, Edgington TS: Selective cellular expression of tissue factor in human tissues. Implications for disorders of hemostasis and thrombosis. *Am J Pathol* 1989; 134: 1087–1097
7. Prudhomme JB, Geiser T, Matthay MA, Ware LB: The alveolar epithelium can initiate the extrinsic coagulation cascade through expression of tissue factor on the cell surface. *Am J Respir Crit Care Med* 2004; 169: A414
8. McVey JH: Tissue factor pathway. *Baillieres Best Pract Res Clin Haematol* 1999; 12: 361–372
9. Idell S, Koenig K, Fair D, Martin T, McLarty J, Maunder R: Serial abnormalities of fibrin turnover in evolving adult respiratory distress syndrome. *Am J Physiol* 1991; 261: L240–L248
10. Fuchs-Buder T, de Moerloose P, Ricou B, Reber G, Vifian C, Nicod L, Romand J, Suter P: Time course of procoagulant activity and D dimer in bronchoalveolar fluid of patients at risk for or with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1996; 153: 163–167
11. Gunther A, Mosavi P, Heinemann S, Ruppert C, Muth H, Markart P, Grimminger F, Walrath D, Temmesfeld-Wolbrück B, Seeger W: Alveolar fibrin formation caused by enhanced procoagulant and depressed fibrinolytic capacities in severe pneumonia. Comparison with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2000; 161: 454–462
12. Gando S, Nanzaki S, Morimoto, Kobayashi S, Kemmotsu O: Systemic activation of tissue-factor dependent coagulation pathway in evolving acute respiratory distress syndrome in patients with trauma and sepsis. *J Trauma* 1999; 47: 719–723
13. Sabharwal AK, Bajaj SP, Ameri S, Tricomi SM, Hyers TM, Dahms TE, Taylor FB, Bajaj MS: Tissue factor pathway inhibitor and von Willebrand factor antigen levels in adult respiratory distress syndrome and in a primate model of sepsis. *Am J Respir Crit Care Med* 1995; 151: 758–767
14. Gando S, Kameue T, Matsuda N, Hayakawa M, Morimoto Y, Ishitani T, Kemmotsu O: Imbalances between the levels of tissue factor and tissue factor pathway inhibitor in ARDS patients. *Thromb Res* 2003; 109: 119–124
15. de Moerloose P, de Benedetti E, Nicod L, Vifian C, Reber G: Procoagulant activity in bronchoalveolar fluids: no relationship with tissue factor pathway inhibitor activity. *Thromb Res* 1992; 65: 507–518
16. Fujii M, Hayakawa H, Urano T, Sato A, Chida K, Nakamura H, Takada A: Relevance of tissue factor and tissue factor pathway inhibitor for hypercoagulable state in the lungs of patients with idiopathic pulmonary fibrosis. *Thromb Res* 2000; 99: 111–117
17. Creasey AA, Chang AC, Feigen L, Wun TC, Taylor FBJ, Hinshaw LB: Tissue factor pathway inhibitor reduces mortality from *Escherichia coli* septic shock. *J Clin Invest* 1993; 91: 2850–2856
18. Enkhbaatar P, Okajima K, Murakami K, Uchiba M, Okabe H, Okabe K, Yamaguchi Y: Recombinant tissue factor pathway inhibitor reduces lipopolysaccharide-induced pulmonary vascular injury by inhibiting leukocyte activation. *Am J Respir Crit Care Med* 2000; 162: 1752–1759
19. Abraham E, Reinhart K, Opal S, Demeyer I, Doig C, Rodriguez AL, Beale R, Svoboda P, Laterre P, Simon S, Light B, Spapen H, Stone J, Seibert A, Peckelsen C, De Deyne C, Postier R, Pettilä V, Artigas A, Percell S, Shu V, Zwingelstein C, Tobias J, Poole L, Stolzenbach J, Creasey A, OPTIMIST Trial Study Group: Efficacy and safety of tifacogin (recombinant tissue fac-

- tor pathway inhibitor) in severe sepsis: a randomized controlled trial. *JAMA* 2003; 290: 238–247
20. Abraham E, Reinhart K, Svoboda P, Seibert A, Olthoff D, Dal Nogare A, Postier R, Hempelmann G, Butler T, Martin E, Zwengelstein C, Percell S, Shu V, Leighton A, Creasey AA: Assessment of the safety of recombinant tissue factor pathway inhibitor in patients with severe sepsis: a multicenter, randomized, placebo-controlled, single-blind, dose escalation study. *Crit Care Med* 2001; 29: 2081–2089
 21. Welty-Wolf KE, Carraway MS, Miller DL, Ortel TL, Ezban M, Ghio AJ, Idell S, Piantadosi CA: Coagulation blockade prevents sepsis-induced respiratory and renal failure in baboons. *Am J Respir Crit Care Med* 2001; 164: 1988–1996
 22. Miller DL, Welty-Wolf K, Carraway MS, Ezban M, Ghio A, Suliman H, Piantadosi CA: Extrinsic coagulation blockade attenuates lung injury and proinflammatory cytokine release after intratracheal lipopolysaccharide. *Am J Respir Cell Mol Biol* 2002; 26: 650–658
 23. Carraway MS, Welty-Wolf KE, Miller DL, Ortel TL, Idell S, Ghio AJ, Petersen LC, Piantadosi CA: Blockade of tissue factor: treatment for organ injury in established sepsis. *Am J Respir Crit Care Med* 2003; 167: 1200–1209
 24. Esmon CT, Stenflo J, Suttie JW: A new vitamin K-dependent protein. A phospholipid-binding zymogen of a serine esterase. *J Biol Chem* 1976; 251: 3052–3056
 25. Stearns-Kurosawa DJ, Kurosawa S, Mollica JS, Ferrell GL, Esmon CT: The endothelial cell protein C receptor augments protein C activation by the thrombin-thrombomodulin complex. *Proc Natl Acad Sci USA* 1996; 93: 10212–10216
 26. Esmon CT: The anticoagulant and anti-inflammatory roles of the protein C anticoagulant pathway. *J Autoimmun* 2000; 15: 113–116
 27. Esmon CT: The protein C pathway. *Chest* 2003; 124: 26S–32S
 28. Esmon CT: Role of coagulation inhibitors in inflammation. *Thromb Haemost* 2001; 86: 51–56
 29. Shimizu S, Gabazza EC, Taguchi O, Yasui H, Taguchi Y, Hayashi T, Ido M, Shimizu T, Nakagaki T, Kobayashi H, Fukudome K, Tsuneyoshi N, D'Alessandro-Gabazza C, Izumizaki M, Iwase M, Homma I, Adachi Y, Suzuki K: Activated protein C inhibits the expression of platelet-derived growth factor in the lung. *Am J Respir Crit Care Med* 2003; 167: 1416–1426
 30. Cheng T, Liu D, Griffin JH, Fernandez JA, Castellino F, Rosen ED, Fukudome K, Zlokovic BV: Activated protein C blocks p53-mediated apoptosis in ischemic human brain endothelium and is neuroprotective. *Nat Med* 2003; 9: 338–342
 31. Sakata Y, Loskutoff DJ, Gladson CL, Hekman CM, Griffin JH: Mechanism of protein C-dependent clot lysis: role of plasminogen activator inhibitor. *Blood* 1986; 68: 1218–1223
 32. Hataji O, Taguchi O, Gabazza EC, Yuda H, Fujimoto H, Suzuki K, Adachi Y: Activation of protein C pathway in the airways. *Lung* 2002; 180: 47–59
 33. Ware LB, Fang X, Matthay MA: Protein C and thrombomodulin in human acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 2003; 285: L514–L521
 34. Yan SB, Helterbrand JD, Hartman DL, Wright TJ, Bernard GR: Low levels of protein C are associated with poor outcome in severe sepsis. *Chest* 2001; 120: 915–922
 35. Liaw PC, Esmon Ct, Kahnamoui K, Schmidt s, Kahnamoui S, Ferrell G, Beaudin S, Julian JA, Weitz JI, Crowther M, Loeb M, Cook D: Patients with severe sepsis vary markedly in their ability to generate activated protein C. *Blood* 2004; 104: 3958–3964
 36. Iba T, Yagi Y, Kidokoro A, Fukunaga M, Fukunaga T: Increased plasma levels of soluble thrombomodulin in patients with sepsis and organ failure. *Surg Today* 1995; 25: 585–590
 37. Krafte-Jacobs B, Brilli R: Increased circulating thrombomodulin in children with septic shock. *Crit Care Med* 1998; 26: 933–938
 38. Kurosawa S, Stearns-Kurosawa DJ, Carson CW, D'Angelo A, Della Valle P, Esmon CT: Plasma levels of endothelial protein C receptor are elevated in patients with sepsis and systemic lupus erythematosus: lack of correlation with thrombomodulin suggests involvement of different pathological processes. *Blood* 1998; 91: 725–727
 39. Bernard G, Vincent J-L, Laterre P-F, La Rosa S, Dhainaut J-F, Lopez-Rodriguez A, Steingrub J, Garber G, Helterbrand J, Ely E: Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 2001; 344: 699–709
 40. Dhainaut J-F, Laterre P-F, Janes JM, Bernard GR, Artigas A, Bakker J, Riess H, Basson BR, Charpentier J, Utterback BG, Vincent JL, Recombinant Human Activated Protein C Worldwide Evaluation in Sepsis (PROWESS) Study Group: Drotrecogin alfa (activated) in the treatment of severe sepsis patients with multiple-organ dysfunction: data from the PROWESS trial. *Intensive Care Med* 2003; 29: 894–903
 41. van der Poll T, Levi M, Nick JA, Abraham E: Activated protein C inhibits local coagulation after intrapulmonary delivery of endotoxin in humans. *Am J Respir Crit Care Med* 2005; in press
 42. Nick JA, Coldren CD, Geraci MW, Poch KR, Fouty BW, O'Brien J, Gruber M, Zarini S, Murphy R, Kuhn K, Richter D, Kast K, Abraham E: Recombinant human activated protein C reduces human endotoxin-induced pulmonary inflammation via inhibition of neutrophil chemotaxis. *Blood* 2004; 104: 3878–3885
 43. Mondino A, Blasi F: uPA and uPAR in fibrinolysis, immunity and pathology. *Trends Immunol* 2004; 25: 450–455
 44. Yepes M, Lawrence DA: New functions for an old enzyme: nonhemostatic roles for tissue-type plasminogen activator in the central nervous system. *Exp Biol Med* 2004; 229: 1097–1104
 45. Chapman HA, Stone OL, Vavrin Z: Degradation of fibrin and elastin by intact human alveolar macrophages *in vitro*. Characterization of a plasminogen activator and its role in matrix degradation. *J Clin Invest* 1984; 73: 806–815
 46. Chapman HA, Stone OL: A fibrinolytic inhibitor of human alveolar macrophages. Induction with endotoxin. *Am Rev Respir Dis* 1985; 132: 569–575
 47. Chapman HA, Yang XL, Sailor LZ, Sugarbaker DJ: Developmental expression of plasminogen activator inhibitor type I by human alveolar macrophages. Possible role in lung injury. *J Immunol* 1990; 145: 3398–3405
 48. Kotani I, Sato A, Hayakawa H, Urano T, Takada Y, Takada A: Increased procoagulant and antifibrinolytic activities in the lungs with idiopathic pulmonary fibrosis. *Thromb Res* 1995; 77: 493–504
 49. Takahashi K, Uwabe Y, Sawasaki Y, Kiguchi T, Nakamura H, Kashiwabara K, Yagyu H, Matsuoka T: Increased secretion of urokinase-type plasminogen activator by human lung microvascular endothelial cells. *Am J Physiol* 1998; 275: L47–L54
 50. Muth H, Maus U, Wygrecka M, Lohmeyer J, Grimminger F, Seeger W, Gunther A: Pro- and antifibrinolytic properties of human pulmonary microvascular versus artery endothelial cells: impact of endotoxin and tumor necrosis factor- α . *Crit Care Med* 2004; 32: 217–226
 51. Grau GE, de Moerloose P, Bulla O, Lou J, Lei Z, Reber G, Mili N, Ricou B, Morel DR, Suter PM: Haemostatic properties of human pulmonary and cerebral microvascular endothelial cells. *Thromb Haemost* 1997; 77: 585–590
 52. Gross TJ, Simon RH, Kelly CJ, Sitrin RG: Rat alveolar epithelial cells concomitantly express plasminogen activator inhibitor-1 and urokinase. *Am J Physiol Lung Cell Mol Physiol* 1991; 260: L286–L295
 53. Gross TJ, Simon RH, Sitrin RG: Expression of urokinase-type plasminogen activator by rat pulmonary alveolar epithelial cells. *Am J Respir Cell Mol Biol* 1990; 3: 449–456
 54. Marshall BC, Xu Q-P, Rao NV, Brown BR, Hoidal JR: Pulmo-

- nary epithelial cell urokinase-type plasminogen activator. *J Biol Chem* 1992; 267: 11462–11469
55. Qiao R, Zhou B, Liebler JM, Li C, Crandall ED, Borok Z: Identification of three genes of known function expressed by alveolar epithelial type I cells. *Am J Respir Cell Mol Biol* 2003; 29: 98–105
56. Prabhakaran P, Ware L, White K, Cross M, Matthay M, Olman M: Elevated levels of plasminogen activator inhibitor-1 in pulmonary edema fluid are associated with mortality in acute lung injury. *Am J Physiol Lung Cell Mol Physiol* 2003; 285: L20–L28
57. Moalli R, Doyle JM, Tahhan HR, Hasan FM, Braman SS, Saldeen T: Fibrinolysis in critically ill patients. *Am Rev Respir Dis* 1989; 140: 287–293
58. Groeneveld AB, Kindt I, Raijmakers PG, Hack CE, Thijs LG: Systemic coagulation and fibrinolysis in patients with or at risk for the adult respiratory distress syndrome. *Thromb Haemost* 1997; 78: 1444–1449
59. Texereau J, Pene F, Chiche JD, Rousseau C, Mira JP: Importance of hemostatic gene polymorphisms for susceptibility to and outcome of severe sepsis. *Crit Care Med* 2004; 32: S313–S319
60. Hermans PW, Hibberd ML, Booy R, Daramola O, Hazelzet JA, de Groot R, Levin M: 4G/5G promoter polymorphism in the plasminogen-activator-inhibitor-1 gene and outcome of meningococcal disease: Meningococcal Research Group. *Lancet* 1999; 354: 556–560
61. Westendorp RG, Hottenga JJ, Slagboom PE: Variation in plasminogen-activator-inhibitor-1 gene and risk of meningococcal septic shock. *Lancet* 1999; 354: 561–563
62. Haralambous E, Hibberd ML, Hermans PW, Ninis N, Nadel S, Levin M: Role of functional plasminogen-activator-inhibitor-1 4G/5G promoter polymorphism in susceptibility, severity, and outcome of meningococcal disease in Caucasian children. *Crit Care Med* 2003; 31: 2788–2793
63. Menges T, Hermans PW, Little SG, Langefeld T, Boning O, Engel J, Sluijter M, de Groot R, Hempelmann G: Plasminogen-activator-inhibitor-1 4G/5G promoter polymorphism and prognosis of severely injured patients. *Lancet* 2001; 357: 1096–1097
64. Gils A, Declerck PJ: The structural basis for the pathophysiological relevance of PAI-1 in cardiovascular diseases and the development of potential PAI-1 inhibitors. *Thromb Haemost* 2004; 91: 425–437