q -COMPLETENESS WITH CORNERS OF UNBRANCHED RIEMANN DOMAINS

George-Ionuț IONIȚĂ

(Received 12 February 2014)

Abstract. We prove that if we consider $p : Y \rightarrow X$ to be an unbranched Riemann domain between two complex spaces with isolated singularities, X is Stein and p is locally q -complete with corners, then Y is q -complete with corners.

1. Introduction

The ‘local Steinness problem’ or the ‘Levi problem on singular spaces’ is one of the most important and difficult problems in several complex variables. A survey concerning the Levi problem on Stein spaces is given in [5] and [22]. For complex spaces with isolated singularities this problem was solved by Andreotti and Narasimhan in [2]. Namely they proved that if X is a Stein space with isolated singularities and $Y \subset X$ is a locally Stein open subset of X , then Y is Stein.

This result was generalized by Coltoiu and Diederich in [6], as they showed that if $p : Y \rightarrow X$ is an unbranched Riemann domain between two complex spaces with isolated singularities such that X is Stein and p is a Stein morphism, then Y is also Stein.

The notions of q -convex and q -complete complex spaces were introduced by Andreotti and Grauert in [1]. They proved finiteness and vanishing theorems for the cohomology of q -convex and q -complete spaces with values in a coherent analytic sheaf.

The purpose of this paper is to prove the following result which generalizes [6] (p being locally 1-complete with corners corresponds to p being a Stein morphism).

THEOREM 1. *Let X and Y be complex spaces with isolated singularities and $p : Y \rightarrow X$ an unbranched Riemann domain. Assume that X is Stein and that p is locally q -complete with corners, i.e. each point $x \in X$ has a neighbourhood $V = V(x)$ such that $p^{-1}(V)$ is q -complete with corners. Then Y is q -complete with corners.*

If p is the inclusion map, then the above theorem was proved by Vâjăitu in [23]. Also if X and Y are smooth, Vâjăitu showed that if X is r -complete with corners and p is locally q -complete with corners, then Y is $(q + r - 1)$ -complete with corners (see [24]).

2010 Mathematics Subject Classification: Primary 32F10, 32C55; Secondary 32C15.

Keywords: q -complete with corners space; unbranched Riemann domain.

2. Preliminaries

In the following all complex spaces are assumed to be reduced and with countable topology. If X is a complex space, then by $T_x X$ we denote the holomorphic tangent space of X at a point $x \in X$. As usual, set $TX := \bigcup_{x \in X} T_x X$. If we are referring to $\mathbb{C}^n$, then the tangent space of $\mathbb{C}^n$ at any point $x \in \mathbb{C}^n$ is just $\mathbb{C}^n$.

Let X be a complex space and U an open subset of X . A (local) chart of X is a holomorphic embedding $i : U \hookrightarrow U'$, where U' is an open subset of $\mathbb{C}^n$ (for some integer $n \geq 1$) such that $i(U)$ is an analytic subset of U' and U and $i(U)$ are biholomorphic (via i).

Now, following Andreotti and Grauert [1], we define the notion of a q -convex function on a complex space.

Let $f : D \rightarrow \mathbb{R}$ be a smooth real function defined on some open subset $D \subset \mathbb{C}^n$. If $z_0 \in D$, then we denote by $L(f, z_0)$ the Levi form of f at z_0 , namely

$$L(f, z_0)(\xi, \eta) = \sum_{i,j=1}^n \frac{\partial^2 f}{\partial z_i \partial \bar{z}_j}(z_0) \xi_i \bar{\eta}_j,$$

where $\xi, \eta \in \mathbb{C}^n$. If $\xi = \eta$, then we simply denote the Levi form of f at z_0 by $L(f, z_0)\xi$. A function f is called q -convex if its Levi form has at least $n - q + 1$ positive (> 0) eigenvalues at any point of D .

Using local charts the above notion can be generalized to arbitrary complex spaces.

Definition 1. Let X be a complex space. A function $f : X \rightarrow \mathbb{R}$ is called q -convex at a point $x \in X$ if there exists a local chart $i : U \hookrightarrow U'$ of X , $x \in U$ and a smooth real q -convex function $f' : U' \rightarrow \mathbb{R}$ such that $f' \circ i = f|_U$.

The function f is said to be q -convex on a subset $W \subset X$ if it is q -convex at every point of W .

An upper semi-continuous function $f : X \rightarrow \mathbb{R}$ is said to be an *exhaustion* function on X if the sublevel sets $\{x \in X : f(x) < c\}$ are relatively compact for any $c \in \mathbb{R}$.

Definition 2. A complex space X is said to be q -convex, if there exists a compact subset K of X and a smooth exhaustion function $f : X \rightarrow \mathbb{R}$, which is q -convex on $X \setminus K$. If we can choose $K = \emptyset$, then X is said to be q -complete.

By [9] and [18], the function $f \in C^0(X, \mathbb{R})$ is called q -convex with corners if for every point $x \in X$ there is an open neighbourhood U of x and finitely many q -convex functions $f_1, \dots, f_k$ on U such that

$$f|_U = \max(f_1, \dots, f_k).$$

We denote by $F_q(X)$ the set of all q -convex functions with corners on X .

Definition 3. A complex space X is said to be q -convex with corners, if there exists a compact subset K of X and an exhaustion function $f : X \rightarrow \mathbb{R}$, which is q -convex with corners on $X \setminus K$. If we can choose $K = \emptyset$, then X is said to be q -complete with corners.

Definition 4. Let $D \subset \mathbb{C}^n$ be an open subset and $f : D \rightarrow [-\infty, \infty)$ an upper semicontinuous function. We say that f is subpluriharmonic if for every relatively compact subset $G \subset\subset D$ and for every pluriharmonic function h defined on a neighbourhood of $\overline{G}$

(i.e. h is locally the real part of a holomorphic function) such that $f|_{\partial G} \leq h|_{\partial G}$ we have also $f \leq h$ on $\overline{G}$.

Definition 5. Let X be a complex space, $f : X \rightarrow [-\infty, \infty)$ an upper semicontinuous function and q a positive integer. We say that f is:

- (1) q -plurisubharmonic if for every open set $G \subset \mathbb{C}^q$ and every holomorphic map $g : G \rightarrow X$, the function $f \circ g$ is subpluriharmonic;
- (2) strongly q -plurisubharmonic if for every $\theta \in C_0^\infty(X, \mathbb{R})$ there exists $\epsilon > 0$ such that the function $f + \epsilon\theta$ is q -plurisubharmonic on X .

For a complex space X , let $P_q(X)$ be the set of all q -plurisubharmonic functions on X and $SP_q(X)$ the set of strictly q -plurisubharmonic functions.

In the literature (see [11, 12, 15, 16, 17, 21]) the concepts of subpluriharmonicity and (strongly) q -plurisubharmonicity are defined in various ways. For example, in [15], a function defined on an open subset $D \subset \mathbb{C}^n$ and with values in $[-\infty, \infty)$ is called q -plurisubharmonic ($1 \leq q \leq n$) in D if it is upper semicontinuous and if it is subpluriharmonic on the intersection of every q -dimensional complex plane with D . Using local embeddings we can generalize this definition to arbitrary complex spaces. Fujita proved in [12] that for X smooth the above notions coincide. This happens also in the singular case for $q = 1$ (see [10]). For $q > 1$, but only for continuous functions, this result was announced by Popa-Fischer in [20] and proved in [21].

The next approximation result was proved by Bungart [3] for open subsets of some $\mathbb{C}^n$, but Matsumoto [17] remarked that it is also true for complex manifolds.

THEOREM 2. *Let X be a complex manifold and $f : X \rightarrow \mathbb{R}$ a continuous strongly q -plurisubharmonic function. Then for an arbitrary continuous function $\delta : X \rightarrow (0, \infty)$ there exists a function $\tilde{f} \in F_q(X)$ such that $|\tilde{f} - f| < \delta$.*

In [7] Colţoiu and Mihalache proved the following result.

THEOREM 3. *Let X be a 1-convex complex space. Then X carries a strongly plurisubharmonic exhaustion function $\Phi : X \rightarrow [-\infty, \infty)$. Moreover, Φ can be chosen $-\infty$ exactly on the exceptional set S of X and real analytic outside S .*

We give now a criterion for q -completeness with corners of Văjăitu [24].

PROPOSITION 1. *Let X be a complex space and $\Phi \in F_q(X)$ such that for every $c \in \mathbb{R}$ the set $X_c := \{\Phi < c\}$ is q -complete with corners. Then X is q -complete with corners.*

From [4] we quote the following result.

LEMMA 1. *Consider X to be a complex space, $A \subset X$ an analytic subset and $f \in F_q(A)$. Then for every $\eta \in C_0^0(A, \mathbb{R})$, $\eta > 0$ there is an open neighbourhood U of A in X and $\tilde{f} \in F_q(U)$ such that $|\tilde{f}|_A - f| \leq \eta$.*

The following result was proved by Peternell in [19].

LEMMA 2. *Let X be a complex space and $A \subset X$ an analytic subset. Then there exists $h \in C^\infty(X, \mathbb{R})$, $h \geq 0$ such that:*

- (a) $\{h = 0\} = A$;

- (b) for every $x \in X$ there exists an open neighbourhood U of x and a smooth function $\sigma : U \rightarrow \mathbb{R}$ such that $\log(h|_{U \setminus A}) + \sigma|_{U \setminus A}$ is plurisubharmonic.

The function $\log h$ is locally equal to the sum of a plurisubharmonic function and a smooth function. Such a function is called *almost plurisubharmonic* or *quasi-plurisubharmonic* (see [8]).

3. The proof

First we assume that $\text{Sing}(X)$ is a finite set and that $p(Y)$ is relatively compact in X . Let us even assume that $\text{Sing}(X) = \{x_0\}$. Otherwise the proof is almost the same.

There are two possibilities.

Case 1: $x_0 \notin p(Y)$. Consider $\pi : \tilde{X} \rightarrow X$ to be a resolution of the singularity x_0 . Thus, $\tilde{X}$ is a 1-convex manifold. From Theorem 3 we get a strongly plurisubharmonic exhaustion function $\varphi : \tilde{X} \rightarrow [-\infty, \infty)$ which can be chosen $-\infty$ exactly on the exceptional set of $\tilde{X}$.

Since $x_0 \notin p(Y)$, we consider $p_1 : Y \rightarrow \tilde{X}$ to be the Riemann domain such that $p = \pi \circ p_1$. Because p is locally q -complete with corners, we have that p_1 is also locally q -complete with corners. We denote by $U \subset \subset \tilde{X}$ a strongly pseudoconvex neighbourhood of B (see [13]). This neighbourhood can be chosen such that $p_1^{-1}(U)$ is q -complete with corners. Also on $\tilde{X}$ we have a smooth plurisubharmonic function $\alpha : \tilde{X} \rightarrow \mathbb{R}_+$ such that α is 0 on U and α is ≥ 0 and strongly plurisubharmonic on $\tilde{X} \setminus \overline{U}$. We choose strongly pseudoconvex neighbourhoods V and V' of B with $B \subset U \subset \subset V \subset \subset V'$ and such that $p_1^{-1}(V')$ is q -complete with corners. Also we may assume that $\varphi \geq 0$ outside U . Since $p_1^{-1}(V')$ is q -complete with corners, we denote by $h_1 : p_1^{-1}(V') \rightarrow \mathbb{R}_+$ a smooth q -convex with corners exhaustion function.

In what follows, some arguments are similar to those in [6]. However, for reader's convenience, we repeat them here.

Since $K := \overline{p_1(Y)} \subset \tilde{X}$ there exists a finite number of open balls $\{U_i\}$, such that $Y_i := p_1^{-1}(U_i)$ is q -complete with corners. Thus, $-\log \delta_i$ is a q -plurisubharmonic function (see [24, Proposition 7, p. 513]), where δ_i represents the boundary distance measured in the euclidian metric for the Riemann domain $Y_i \rightarrow U_i \subset \subset \mathbb{C}^n$. Consider now concentric balls $V_i \subset \subset U_i$ such that K is still covered by V_i . By Lemma 3 of Matsumoto [16] (or Peternell [18]) the quotients δ_i/δ_j are bounded on $p_1^{-1}(V_i \cap V_j)$. Therefore, the differences $\log \delta_j - \log \delta_i$ are also bounded.

For each i we can suitably choose a function $\theta_i \in C_0^\infty(V_i)$, $\theta_i \geq 0$ such that the function

$$l(y) := \max_{p_1(y) \in V_i} (-\log \delta_i(y) + \theta_i(p_1(y)))$$

is continuous on Y . Moreover, for a sufficiently large constant $A > 0$, the function

$$q := A \cdot \varphi \circ p_1 + l$$

is strongly q -plurisubharmonic on Y and has the following property:

$$p_1(\{q < c\}) \subset \subset \tilde{X} \quad \text{for every } c \in \mathbb{R}.$$

For the details one should consult Matsumoto [16, pp. 107–108].

The idea is to use Lemmas 1 and 2 for $A = \text{Sing}(Y)$. Since Y is with isolated singularities we have that $\dim A = 0$; hence, A is q -complete with corners. Lemma 1 gives us an open neighbourhood U of A and a function $g \in F_q(U)$.

Using Lemma 2, there is a smooth function $h : Y \rightarrow [0, \infty)$ such that $\{h = 0\} = A$ and $\log h$ is almost plurisubharmonic. Now select $\chi \in C^\infty(\mathbb{R}, \mathbb{R})$ strictly increasing and convex such that $q' := \chi \circ q + \log h \in \text{SP}_q(Y \setminus A)$ and $q' > 1 + g$ on ∂U . By Theorem 2 there is $f \in F_q(Y \setminus A)$ such that $|f - q'| < 1$ on $Y \setminus A$. Finally we can define a function $\Phi : Y \rightarrow \mathbb{R}$ such that $\Phi \in F_q(Y)$:

$$\Phi := \begin{cases} f & \text{on } Y \setminus U, \\ \max(f, g) & \text{on } U \setminus A, \\ g & \text{near } A. \end{cases}$$

In order to prove that Y is q -complete with corners, we will apply Proposition 1. We have to show that for every $c \in \mathbb{R}$ the set $\{\Phi < c\}$ is q -complete with corners, i.e. to define for every $c \in \mathbb{R}$ an exhaustion function $\eta_c : \{\Phi < c\} \rightarrow \mathbb{R}$ that is q -convex with corners.

Let us denote by g a fixed Riemannian metric on $\tilde{X}$ and by g^* its pull-back to Y . We define for each $\epsilon > 0$ the set

$$Y_\epsilon := \{y \in Y : \delta(y) > \epsilon\},$$

where δ is the induced boundary distance on the Riemann domain $p_1 : Y \rightarrow \tilde{X}$.

Using the regularization method of Hörmander [14, pp. 141–142], Coltoiu and Diederich [6] constructed a C^2 -function $\phi = \phi_\epsilon : Y_\epsilon \rightarrow \mathbb{R}_+$ such that ϕ is a Lipschitz vertical exhaustion function on Y_ϵ and the Levi form of ϕ is bounded from below.

We define the function $\mu := \phi \cdot \tilde{\alpha} : \{\Phi < c\} \rightarrow \mathbb{R}$, where $\tilde{\alpha} = \alpha \circ p_1$. In order for μ to be well defined we choose a small enough $\epsilon = \epsilon_c > 0$ such that $\{\Phi < c\} \setminus p_1^{-1}(U) \subset \{\delta > \epsilon\}$.

From the formula

$$L(\mu) = \tilde{\alpha}L(\phi) + \phi L(\tilde{\alpha}) + 2\text{Re}(\partial\phi)(\bar{\partial}\tilde{\alpha})$$

we get that $L(\mu)$ is bounded from below on $\{\Phi < c\}$, μ is a vertical exhaustion function outside $p_1^{-1}(\bar{U})$ and it is identically 0 on $p_1^{-1}(\bar{U})$. Now choose a strongly pseudoconvex neighbourhood $U' \subset\subset U$ of B such that there exists a smooth 1-convex function $\psi : \tilde{X} \rightarrow \mathbb{R}_+$ which enjoys the same properties as the function α . So, for a sufficiently large constant C , the function

$$\mu + C \cdot \psi \circ p_1 : \{\Phi < c\} \rightarrow \mathbb{R}$$

is 1-convex on all $\{\Phi < c\}$, relatively exhausting outside $p_1^{-1}(\bar{U})$ and identically 0 on $p_1^{-1}(\bar{U})$. Now we select a smooth rapidly increasing strictly convex function $\chi : [0, \infty) \rightarrow [0, \infty)$, $\chi(0) = 0$ such that

$$\chi \circ (\mu + C \cdot \psi \circ p_1) > h_1 \quad \text{on } \{\Phi < c\} \cap p_1^{-1}(\partial V).$$

Remember that the goal is to construct a continuous q -convex with corners exhaustion function on $\{\Phi < c\}$. Now consider the maximum between h_1 and $\chi \circ (\mu + C \cdot \psi \circ p_1)$ over $\{\Phi < c\} \cap p_1^{-1}(V)$. Next we extend this function by $\chi \circ (\mu + C \cdot \psi \circ p_1)$ and we obtain a q -convex with corners function on $\{\Phi < c\}$; let us denote it by λ_c . Therefore, the function $\eta_c := \lambda_c + 1/(c - \Phi)$ is a continuous q -convex with corners exhaustion function on $\{\Phi < c\}$. Using Proposition 1 we get that Y is q -complete with corners.

Case 2: $x_0 \in p(Y)$. Let $\pi : \tilde{X} \rightarrow X$ be the local desingularization at x_0 . Now consider the fibre product of $p : Y \rightarrow X$ and π , that is the set $\tilde{Y} = \{(y, \tilde{x}) \in Y \times \tilde{X} : p(y) = \pi(\tilde{x})\}$. We have obtained two projection maps: one on Y which will be denoted by π_1 and another one on $\tilde{X}$ which will be denoted by p_1 . As before let $B := \pi^{-1}(\{x_0\})$ be the exceptional set of $\tilde{X}$. We have that π_1 is a proper modification of Y at the discrete set $p^{-1}(\{x_0\}) = \{a_n\}_n$ and $p_1 : \tilde{Y} \rightarrow \tilde{X}$ is a Riemann domain over $\tilde{X}$. Now the proof is almost the same as in case 1: p_1 is locally q -complete with corners, thus there exists a strongly pseudoconvex neighbourhood $U \subset \subset \tilde{X}$ of B such that $p_1^{-1}(U)$ is a proper modification at a discrete subset of a q -complete with corners space. We get a strongly plurisubharmonic exhaustion function $\tilde{\varphi} : \tilde{Y} \rightarrow \mathbb{R}$ which can be chosen $-\infty$ exactly on the exceptional set $\tilde{B}$ of $\tilde{Y}$. Finally, for each $c \in \mathbb{R}$, we have on the open set $\{\tilde{\varphi} < c\} \subset \tilde{Y}$ a continuous real-valued q -convex with corners exhaustion function. Using the fact that π_1 is biholomorphic outside $\tilde{B}$, we can define $\Phi = \tilde{\varphi} \circ \pi_1^{-1}$ on $Y \setminus p^{-1}(\{x_0\})$ and $\Phi = -\infty$ on $p^{-1}(\{x_0\})$. From Proposition 1 we get that Y is q -complete with corners.

Now we only have to prove that it is enough to assume $\text{Sing}(X)$ to be a finite set, i.e. knowing that $\text{Sing}(X)$ is finite we should infer that Y is q -complete with corners. Since X is Stein, let us denote by ψ a continuous strongly plurisubharmonic exhaustion function on X . We write $X = \bigcup_c X_c$, where $X_c = \{\psi < c\}$. Since X has isolated singularities we have that $\text{Sing}(X_c)$ is a finite set. We put $Y_c = p^{-1}(X_c)$ and we get that each Y_c is q -complete with corners. We have that $\Phi := \psi \circ p$ is 1-convex on Y . Since $Y_c = \{\psi \circ p < c\}$ and using Proposition 1 we have that Y is q -complete with corners.

Using the same argument as before, we can assume that $p(Y) \subset \subset X$.

Acknowledgements. I am very grateful to Professor Mihnea Colţoiu for suggesting this problem to me and for his helpful advice. The author was supported by CNCS grant PN-II-ID-PCE-2011-3-0269.

REFERENCES

- [1] A. Andreotti and H. Grauert. Théorèmes de finitude pour la cohomologie des espaces complexes. *Bull. Soc. Math. France* **90** (1962), 193–259.
- [2] A. Andreotti and R. Narasimhan. Oka's Heftungslemma and the Levi problem for complex spaces. *Trans. Amer. Math. Soc.* **111** (1964), 345–366.
- [3] L. Bungart. Piecewise smooth approximations to q -plurisubharmonic functions. *Pacific J. Math.* **142** (1990), 227–244.
- [4] M. Colţoiu. n -concavity of n -dimensional complex spaces. *Math. Z.* **210** (1992), 203–206.
- [5] M. Colţoiu. The Levi problem on Stein spaces with singularities. A survey. *Rend. Mat. (Roma)* **29** (2009), 341–353.
- [6] M. Colţoiu and K. Diederich. The Levi problem for Riemann domains over Stein spaces with singularities. *Math. Ann.* **338** (2007), 283–289.
- [7] M. Colţoiu and N. Mihalache. Strongly plurisubharmonic exhaustion functions on 1-convex spaces. *Math. Ann.* **270** (1985), 63–68.
- [8] J.-P. Demailly. Cohomology of q -convex spaces in top degrees. *Math. Z.* **204** (1990), 283–296.
- [9] K. Diederich and J. E. Fornaess. Smoothing q -convex functions in the singular case. *Math. Ann.* **273** (1986), 665–671.
- [10] J. E. Fornaess and R. Narasimhan. The Levi problem on complex spaces with singularities. *Math. Ann.* **248** (1980), 47–72.
- [11] O. Fujita. Domaines pseudoconvexes d'ordre général et fonctions pseudoconvexes d'ordre général. *J. Math. Kyoto Univ.* **30** (1990), 637–649.

- [12] O. Fujita. On the equivalence of the q -plurisubharmonic functions and the pseudoconvex functions of general order. Ann. Rep. Graduate School Human Culture. Nara Women's Univ. **7** (1992), 77–81.
- [13] H. Grauert. Über Modifikationen und exzeptionelle analytische Mengen. Math. Ann. **146** (1962), 331–368.
- [14] L. Hörmander. An Introduction to Complex Analysis in Several Variables. North-Holland, Amsterdam, 1990.
- [15] L. R. Hunt and J. J. Murray. q -plurisubharmonic functions and a generalized Dirichlet problem. Michigan Math. J. **25** (1978), 299–316.
- [16] K. Matsumoto. Pseudoconvex Riemann domains of general order over Stein manifolds. Kyushu J. Math. **44** (1990), 95–107.
- [17] K. Matsumoto. Boundary distance functions and q -convexity of pseudoconvex domains of general order in Kähler manifolds. J. Math. Soc. Japan **48** (1996), 85–107.
- [18] M. Peternell. Continuous q -convex exhaustion functions. Invent. Math. **85** (1986), 249–262.
- [19] M. Peternell. Algebraische Varietäten und q -vollständige komplexe Räume. Math. Z. **200** (1989), 547–581.
- [20] A. Popa-Fischer. Sur un théorème de Fornaess et Narasimhan. C. R. Acad. Sci. Paris Sér. I Math. **329** (1999), 11–14.
- [21] A. Popa-Fischer. A generalization to the q -convex case of a theorem of Fornaess and Narasimhan. Michigan Math. J. **50** (2002), 483–492.
- [22] Y.-T. Siu. Pseudoconvexity and the problem of Levi. Bull. Amer. Math. Soc. **840** (1978), 481–512.
- [23] V. Văjăitu. Locally q -complete open sets in Stein spaces with isolated singularities. Kyushu J. Math. **51** (1997), 355–368.
- [24] V. Văjăitu. Pseudoconvex domains over q -complete manifolds. Ann. Sc. Norm. Sup. Pisa Cl. Sci. (5) **29** (2000), 503–530.

George-Ionuț Ioniță

Department of Mathematics and Computer Science

'Politehnica' University of Bucharest

313 Splaiul Independenței

Bucharest 060042, Romania

and

Simion Stoilow Institute of Mathematics of the Romanian Academy

Research group of the project ID-3-0269

P.O. Box 1-764

Bucharest 014700, Romania

(E-mail: georgeionutionita@yahoo.com)