Review

Matrix-Assisted Laser Desorption/Ionization Mass Spectrometry: Mechanistic Studies and Methods for Improving the Structural Identification of Carbohydrates

Yin-Hung Lai^{*1} and Yi-Sheng Wang²

¹Department of Chemistry, Stanford University, Stanford, California 94305, USA

²Genomics Research Center, Academia Sinica, Taipei 115, Taiwan, ROC

Although matrix-assisted laser desorption/ionization (MALDI) mass spectrometry is one of the most widely used soft ionization methods for biomolecules, the lack of detailed understanding of ionization mechanisms restricts its application in the analysis of carbohydrates. Structural identification of carbohydrates achieved by MALDI mass spectrometry helps us to gain insights into biological functions and pathogenesis of disease. In this review, we highlight mechanistic details of MALDI, including both ionization and desorption. Strategies to improve the ion yield of carbohydrates are also reviewed. Furthermore, commonly used fragmentation methods to identify the structure are discussed.

Copyright © 2017 Yin-Hung Lai and Yi-Sheng Wang. This is an open access article distributed under the terms of Creative Commons Attribution License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Please cite this article as: Mass Spectrom (Tokyo) 2017; 6(3): S0072

Keywords: matrix-assisted laser desorption/ionization, mechanism, carbohydrate, ion yield, structural identification

(Received July 31, 2017; Accepted August 3, 2017)

INTRODUCTION

Matrix-assisted laser desorption/ionization (MALDI) is one of the most important soft ionization methods for biological mass spectrometry.^{1,2} Several unique aspects of MALDI, such as the production of primarily singly-charged species and the high tolerance against contaminants, make MALDI a promising tool to explore unknown structures in a complex system.^{3,4} It has become one of the most well-established approaches in proteomics owing to its high sensitivity, low sample consumption, and minimal sample handling. With the growing interest in biological roles of carbohydrates, the structural identification of carbohydrates has also been attempted by MALDI mass spectrometry. However, compared to proteins, carbohydrates have lower ionization efficiency in MALDI, making carbohydrate analysis difficult. In fact, the lack of knowledge about the reaction mechanism in MALDI has restricted its applicability in analytical chemistry.

Carbohydrates (also called glycans) play important roles in mediating intercellular and intracellular functions such as protein folding, signal transduction, and cell-cell interaction.⁵⁻⁷ In particular, glycans become attached to proteins during glycosylation, which is one of the most ubiquitous protein post-translational modification (PTM). Aberrant

glycosylation has been found to indicate the development and progression of cancer.⁸ Therefore, expression profiles of glycans are utilized as biomarkers for diagnosis and prognosis of cancers as well as therapeutic evaluation.⁹⁻¹³

The “holy grail” of glycoanalysis is to be able to characterize the oligosaccharide heterogeneity at each glycosylation site on a protein or proteoglycan.¹⁴ The structural characterization of biologically significant glycoconjugates also presents opportunities for the synthesis of specific glycans to provide new therapeutics, such as carbohydrate-based vaccines for cancer.¹⁵ However, owing to their highly branched structures, thermally labile properties, and lower ionization efficiency, improvement of ion abundance as well as the structural determination of carbohydrate identification and sequencing with MS are still challenging. The study of ionization mechanisms is therefore crucial to the facilitation of efficient analysis of carbohydrates.

In this review, we focus first on the mechanisms of MALDI, including ionization and desorption processes. The discussion of underlying ionization processes of carbohydrates and methods for improving structural analysis of carbohydrates are reviewed, with special emphasis in the analysis of glycans in glycopeptides and glycoproteins. For further advances in carbohydrate analysis, please also refer to the recent review articles and books.¹⁶⁻²¹

* Correspondence to: Yin-Hung Lai, Department of Chemistry, Stanford University, Stanford, California 94305, USA, e-mail: lai@stanford.edu

MALDI MECHANISMS

Matrix-assisted laser desorption/ionization comprises photochemical (ionization and fragmentation) and mechanical (desorption) events. In MALDI, analytes are incorporated into organic matrices in a solid or viscous phase (ionic liquid matrices). A pulsed laser irradiates the sample to facilitate the ionization/desorption reactions. During laser excitation, matrices take up and redistribute the laser energy for desorption and ionization of analytes. In this section, we first review critical factors determining efficiencies of ionization and desorption. Models proposed for ionization and desorption processes from the perspectives of both experimental observation and theory are discussed. After the review of general MALDI mechanisms, the specific aspects of ionization mechanisms on carbohydrates and methods that improve ion abundances of carbohydrates are reviewed.

Ion production in MALDI

Owing to the high complexity of the MALDI process, a single ionization model that could elucidate phenomenon universally to every MALDI experiment is highly unlikely. Ionization environment varies among different matrix systems as well as excitation conditions.²²⁾ For investigating mechanisms in MALDI, the ion-to-neutral ratio (ionization efficiency) is normally discussed.^{23–26)} Typically, ionization efficiency differs across types of analytes. The ionization efficiency for molecules with higher proton affinity than matrices, such as peptides and proteins, is 10^{-3} – 10^{-4} .^{27–32)} For analytes with lower proton affinity, such as carbohydrates, the ionization efficiency drops by a few orders of magnitude to 10^{-7} – 10^{-8} .^{23,33,34)} A better understanding of mechanisms leads to improvements in the ionization efficiency and increases its effectiveness toward more fields. Over the last three decades, fundamental ionization mechanisms have been systematically studied by manipulating one or more parameters among different matrix systems.^{22,35)} Effects of the incorporation of analytes into matrix solid,^{36–40)} matrix-to-analyte ratio,^{41,42)} solvent constituents,^{43–47)} laser excitation conditions,^{22,48,49)} and the crystal morphology of samples^{47,50–55)} on ion production were investigated.

A prerequisite of successful MALDI processes is absorption of laser energy by matrices. The optimal MALDI-MS sensitivity is generally achieved when the utilized laser wavelength corresponds to the high optical absorption band of the solid-phase matrices.⁵⁶⁾ After photo-absorption, a two-step ionization process suggests that ionization of matrices is as an initial ionization step, followed by ion-molecule reactions to produce charged analyte species.^{35,57)} The second step is less controversial, and ion yields of analytes can be estimated by relative thermodynamic properties between matrices and analytes.^{58–61)} However, how matrices are initially ionized remains open to debate. Two major arguments of postulated initial ionizations include 1) photoionization in the gas phase (Eq. (1)),^{35,62–65)} and 2) thermal energy promoted ionizations in the condensed phase (Eq. (2)).^{34,66–74)}

in which M and M* represent the matrix in the electronic ground and excited states, respectively. In Eq. (1), the posi-

Fig. 1. Initial ionization channels, including photoionization and thermally induced ionization, as well as radiative and radiationless relaxation processes after the absorption of laser photons by matrices. Reprinted with permission from ref. 22. Copyright (2012) American Chemical Society.

tive matrix radical ion and photoelectrons are generated by multi-photon ionization *via* energy pooling/annihilation processes. It should be noted that, although the impact of photoelectrons on the generation of MALDI ion and trigger of fragmentation has been extensively discussed,^{75,76)} the major source of electrons is still in debate.^{75,77)} In Eq. (2), absorbed photon energy is converted to thermal energy *via* ultrafast non-radiative relaxation. In this equation, an increase in the system temperature facilitates both ionization and desorption. Proton disproportionation of matrices is thermally induced to create protonated/deprotonated matrices. Our group proposed a quantitative thermal model and successfully predicted the laser energy dependence of the ion yield.^{68,69)} Both thermal and chemical equilibria can presumably be established by collisions on the solid surface. In our thermal model, ion abundance depended on the concentration on the surface and desorption rate constant.

Contrary to the two equations, the lucky survivor model suggests that matrices and analytes are pre-charged in solids.^{78,79)} In this model, the laser energy overcomes ion-pair interactions to free analyte ions from their respective counter ions. It should be emphasized that a single ionization mechanism that is applied universally to elucidate all MALDI observations is highly unlikely owing to different photochemical properties of matrices as well as excitation conditions. Contributions of different ionization channels need to be considered even for a single matrix system in Fig. 1.²²⁾

Desorption (phase transition)

Desorption in MALDI is the phase transition of samples from the condensed phase to the gas phase. It is a prerequisite for molecules to be detected by MALDI mass spectrometry. Desorption processes, including thermal desorption occurring at low and moderate laser energy density to phase explosion happening at high laser energy density, were studied quantitatively on both experimental and theoretical bases.^{22,80)} Results of these studies suggest that factors affecting the desorption efficiency can be classified into two categories: excitation parameters^{36,81–86)} and intrinsic properties of samples.^{51,87–92)} Generally, desorption efficiency increases when MALDI crystal size decreases⁵¹⁾ and the system temperature increases.^{87–92)}

On the basis of factors influencing desorption efficiency, mathematical equations as well as computational methods, such as molecular dynamics (MD) simulations, have been

Fig. 2. Snapshots from MD simulations of laser ablation of a molecular solid illustrating different mechanisms of material ejection: (a) desorption of monomers; (b) phase explosion of the overheated material; (c) hydrodynamic sputtering due to the fast melting and motion of liquid in the surface region; (d) photomechanical spallation of the surface layer caused by the relaxation of laser-induced thermoelastic stresses. The laser pulse durations are 150 ps (a, b) and 15 ps (c, d), fluences are 34 J/m^2 (a), 61 J/m^2 (b), 40 J/m^2 (c), and 31 J/m^2 (d). The laser penetration depth is 50 nm in all simulations. The irradiation parameters correspond to the regime of thermal confinement in (a) and (b) and to the regime of stress confinement in (c) and (d). Reprinted with permission from ref. 95. Copyright (2003) American Chemical Society.

used to quantitatively describe the desorption process.^{86,93,94} As shown in Fig. 2, Zhigilei *et al.* utilized MD simulations to describe desorption of MALDI after laser irradiation.⁹⁵ However, most of the aforementioned desorption models simplified the impact of ionization, which deviates from the fact that MALDI relies on the complex interplay of both desorption and ionization. By utilizing transition state theory to derive desorption rate, our group developed a comprehensive model that both considered desorption and ionization to predict ion abundance as a function of laser fluence,⁶⁹

$$I_{\text{ion}} = A \frac{k(T_0 + rF)}{h} \left(1 - e^{-(hv/k(T_0 + rF))} \right) e^{-(E_a/k(T_0 + rF))} [\text{ion}]_s \quad (3)$$

in which, A , k , h , v , E_a , represent the pre-exponential factor, Boltzmann constant, Planck constant, vibrational frequency of the intermolecular bond, and activation energy for desorption, respectively. $[\text{ion}]_s$ stands for the concentration of surface ions. T_0 is the initial system temperature, and r is the conversion factor between system temperature and laser energy density. Linear relations between temperature and laser energy density were obtained within typical MALDI conditions.^{69,86,91}

System temperature is a particularly important factor because it not only affects desorption efficiency but also

ionization/fragmentation processes. The predicted system temperature varies with respect to experimental conditions, such as the type of matrices, energy densities, and irradiation wavelengths. Many efforts have been made to evaluate system temperatures of MALDI, such as predictions by theoretical models,^{62,69,70,86,96} MD simulations,³² the measurement of infrared emission of the system based on the theory of black body radiation,^{22,91} and the use of thermometer molecules as probes of internal energy.^{87,88,90,92} Although evaluations by different methods have certain inconsistencies, a consensus of transient MALDI temperature might reach $\sim 1000 \text{ K}$, which is detrimental to thermally labile carbohydrates.

Mechanistic studies of carbohydrate ionization in MALDI

Unlike proteins or nucleic acids that are protonated or deprotonated, carbohydrates are predominately ionized by forming adducts with alkali metal ions (mainly sodium and potassium ions) in MALDI.^{97,98} Several groups have attempted both experimentally and theoretically to understand ionization mechanisms of carbohydrates in MALDI. Mohr *et al.* found that the binding affinities of proton and alkali metal ions to carbohydrates followed the order: $\text{H}^+ \ll \text{Li}^+ < \text{Na}^+ < \text{K}^+ < \text{Cs}^+$.⁹⁹ Since carbohydrates are incorporated into the matrix solid, the relative binding competition between carbohydrates and matrices needs to be considered. The binding affinity of protons to carbohydrates is lower than that to commonly used matrices such as 2,5-dihydroxybenzoic acid (DHB) and 2,4,6-trihydroxyacetophenol (THAP).^{30,100,101} In contrast to the proton affinity, the sodium affinity of carbohydrates is higher than that of matrices.^{98,101,102} Thus, it is energetically favorable for carbohydrates to form the adduct with alkali metal ions rather than with protons. A reaction model proposed by Lee *et al.* suggests that the increase in thermal energy can effectively induce the production of alkali ion adducts of carbohydrates.¹⁰³ They suggested alkali metal ions were generated from the dissolution of salts at high-temperature MALDI condition. Those alkali metal ions attached to nearby matrices and analytes during desorption. However, mechanisms about how carbohydrates are alkalized are still unclear.¹⁰⁴

Ion yields of carbohydrates are related to multiple parameters, such as reaction enthalpies for forming adducts with alkali metal ions, volatility, and structural stabilities.^{101,105} Owing to the thermally labile nature of carbohydrates, Chen *et al.* reported that low ion yield might be the result of rates of fragmentation and/or desodiation (loss of sodium ions back to its neutral form) being faster than sodiation in the high-temperature ionization environment.¹⁰¹ Chen *et al.* also suggested that protonated carbohydrates were easy to decompose through the dehydration reaction. Cancilla *et al.* reported that the fragmentation behavior of carbohydrates was also related to their degree of branching.¹⁰⁶ A branched glycan can establish multiple bindings to an alkali metal ion, resulting in lesser fragments. Computational methods, including MD simulation, semi-empirical and *ab initio* calculations, were utilized by Fukui *et al.* to investigate the relationship between the structure and reactivity (fragmentation) of sodiated oligosaccharides.¹⁰⁵ Fukui *et al.* found that the coordination of metal ions to oligosaccharides affected the propensity for fragmentation. Based on the calculation

Table 1. Methods to improve ion abundances of carbohydrates in MALDI mass spectrometry.

Strategy	Methods	Ref.
Mediating high-temperature ionization environment	Soft ionization by ionic liquid matrices (ILMs)	
	–1,1,3,3-Tetramethylguanidium (TMG) salt of <i>p</i> -coumaric acid (G_3CA)	107
	–3-Aminoquinoline (3-AQ) based ILM	108, 109
	–2,4,6-Trihydroxyacetophene (THAP) based ILM	110
	Ice as a soft matrix	111
	Temperature-controlled sample plate allowing a frozen sample inside the vacuum	112
Doping ionization agent	Trylayer sample preparation method with thermal energy-dissipating diamond nanoparticles	113
	Ammonium salts	114, 115
	Introducing permanent charges with succinimidylxycarbonylmethyl <i>tris</i> (2,4,6-trimethoxy-phenyl) phosphonium bromide (TMPP-AcOSu)	116
	Nanoparticles containing ionization agents	117–123
Chemical derivatization	Permethylation	124, 127, 135
	Amidation	126, 128, 132
	Esterification	133, 134
	Derivatization with hydrazines	125, 129

Fig. 3. Positive-ion mass spectra of 100 fmol fetuin GPI, 100 fmol NA4 glycan, and 10 fmol β -casein 1–25 using (a, d, g) 3-AQ/CA, (b, e, h) 3-AQ/CHCA, and (c, f, i) 2,5-DHB. Reprinted with permission from ref. 109. Copyright (2014) American Chemical Society.

result, increasing the number of oxygen atoms interacting with the sodium ion stabilized the carbohydrate structure (less fragmentation) and increased the ion yield.

Based on the understanding of mechanisms of MALDI and the nature of carbohydrates, three strategies are generally adopted to improve ion yield of carbohydrates: 1) thermalizing high-temperature ionization environments to reduce dealkalization and fragmentation; 2) doping ionization agents (permanent charges) to enhance ion yield; 3) changing the nature of glycans by derivatizations. We highlighted some of selected advancements in Table 1.

The selection of matrices is a crucial step for the analysis of carbohydrates because every matrix produces unique ionization environment after the laser irradiation.²²⁾ Problems associated with decomposition of thermally labile moieties and low ionization efficiency could be overcome by a suitable selection of the matrix. As shown in Fig. 3, “cool” ionic liquid matrices: 1,1,3,3-tetramethylguanidium (TMG) salt of *p*-coumaric acid (G_3CA),¹⁰⁷⁾ and 3-aminoquinoline/*p*-coumaric acid (3-AQ/CA)^{108,109)} were developed by Tanaka and coworkers to yield a less locally heated ionization environment for enhancing the sensitivity and suppressing the losses of sialic acid moieties. Applying a similar concept, THAP-based ionic liquid matrices¹¹⁰⁾ were also reported for automated identification and quantitative analysis of na-

tive oligosaccharides. Witt *et al.* utilized ice as an extremely soft matrix to prevent carbohydrates from decomposition and enhance the capability of MALDI mass spectrometry for the structural characterization of glycans.¹¹¹⁾ The use of ice as a matrix is advantageous because it provides an ideal “near-physiological” condition. Liang *et al.* modified the ion source of commercial MALDI time-of-flight mass spectrometer to enable the preparation of a frozen carbohydrate/DHB sample at 100 K. Enhancement of sensitivity (20–30 folds) and reduction of fragmentation were achieved by this method.¹¹²⁾ Our group developed a trilayer sample preparation method, which changes sample morphology by consecutively depositing matrices, diamond nanoparticles, and carbohydrates. Owing to the high thermal conductivity, diamond nanoparticles thermalized the MALDI temperature. Diamond nanoparticles also adjusted the incorporation of matrices and carbohydrates. As shown in Fig. 4, we selectively enhanced the sensitivity of underivatized carbohydrates by more than an order of magnitude from protein/carbohydrate mixtures.¹¹³⁾ This sample preparation method is potentially suitable for diagnosing carbohydrates in real samples because carbohydrates are often accompanied by proteins in real biological samples.

In the second category, Yamagaki *et al.* improved ionization efficiency of neutral glycans by adding ammonium

Fig. 4. Mass spectra of 5 pmol each of dextran and ACTH fragment 18–39 obtained from various samples, including (a) dried-droplet, (b) thin-layer, and (c) trilayer samples. The numbers indicate the degrees of polymerization (n) of sodiated dextran. Reprinted with permission from ref. 113. Copyright (2013) American Chemical Society.

salts in MALDI samples to form chloride-anionized molecules.¹¹⁴ In a similar fashion, doping salts containing anions, such as NO_3^- and Cl^- , was conducted by Domann *et al.* to enhance the ionization efficiency of neutral N-linked glycans.¹¹⁵ Domann *et al.* reported the best matrix for ionizing glycans was mixing THAP with ammonium nitrate (1 M) to form $[\text{M}+\text{NO}_3]^+$. Gao *et al.* reported a derivatization method that enhanced 20- to 50-fold ion abundance of N-linked glycans by introducing permanent charges using nonspecific proteolysis with pronase E and co-derivatization with succinimidylsuccinylmethyl tris(2,4,6-trimethoxyphenyl)-phosphonium bromide (TMPP-AcOSu) and methylamidation.¹¹⁶ Nanoparticles (NPs), such as gold, silver, titanium dioxide, iron oxide (Fe_3O_4), zinc oxide, and manganese oxide, have been utilized for the analysis of carbohydrates because they enhance ionization efficiency by releasing ionization agents.^{117–121} After absorbing laser energy, NPs also convert photon energy to thermal energy to facilitate material desorption. Charge and energy transfer from NPs to analytes triggers ionization/fragmentation and desorption of carbohydrates.^{117,122} Furthermore, functionalization of NPs with conventional organic matrices have been reported to serve as new matrix materials to preserve the intact glycan as well as the generation of extensive cleavages in MALDI.¹²³ More details about methods on the structural identification of carbohydrates are reviewed in the following section.

Chemical derivatization has been accepted as a well-established option for the analysis of carbohydrates in MALDI¹²⁴ because chemical derivatization protects glycans from losing acetyl groups or sialic acid moieties to increase carbohydrate ionization efficiency.^{125,126} Hakomori first demonstrated permethylation of complex carbohydrates.¹²⁷ Derivatization was also reported to be beneficial to compensate differences in ionization efficiency. A novel amidation

using acetohydrazide was demonstrated by Toyoda *et al.* to eliminate the discrepancy of ionization efficiency between α 2,3- and α 2,6-linked sialic acids-containing oligosaccharides.¹²⁸ Derivatization also improves chromatographic analysis by reducing the polarity of glycans. Various methods for derivatization have been developed, such as uses of hydrazines¹²⁹ and other labeling techniques.¹³⁰

Although derivatization provides advantages, conventional derivatization typically requires harsh conditions (strong base) and long reaction times. Most derivatization methods require clean-up procedures for the removal of chemicals, resulting in a loss of samples, which especially leads to a serious problem for trace analysis.¹³¹ Thus, derivatization methods have been improved to meet the demand of low sample loss and high efficiency.¹³² Miura *et al.* reported a solid-phase methyl esterification of sialic acid residues of glycans for rapid and quantitative glycoform profiling by MALDI-TOF MS.¹³³ As shown in Fig. 5, the workflow of their on-bead/on gold nanoparticle derivatization is potentially suitable for the large-scale glycomics study. Using a combination of carboxylic acid activators in ethanol, Reiding *et al.* achieved nearly complete ethyl esterification of α 2,6-linked sialic acids and lactonization of α 2,3-linked variants in a short time and mild conditions.¹³⁴ A solid-phase miniaturized approach was demonstrated by Kang *et al.* for quantitative permethylation of oligosaccharides in less than a minute.¹³⁵ By mixing analytes with methyl iodide in dimethyl sulfoxide solution, this method avoided excessive clean-up and yielded picomole-level sensitivity.

FRAGMENTATION METHODS FOR THE STRUCTURAL ANALYSIS OF CARBOHYDRATES

Structural information of carbohydrates, including sequence from glycosidic cleavages (C–O bonds) and branching and linkages from cross-ring cleavages (C–C bonds), is available by utilizing multiple fragmentation techniques in tandem mass spectrometry. Fragmentation techniques are essential methods to deposit energy into a targeted ion to cause reproducible bond cleavages. Cleavages yield diagnostic and interpretable fragment ions that reveal structural or sequence information about the molecule of interest. Commonly used fragmentation techniques in MALDI mass spectrometry include laser-induced promptly in-source decay (ISD),¹³⁶ post-source decay (PSD, a unimolecular dissociation of metastable ions),¹³⁷ and collision-induced dissociation (CID).^{138–140}

ISD ions are promptly produced by thermal activation and/or chemical reactions during desorption/ionization processes in the source region, such as hydrogen radical mediated¹⁴¹ and/or free electron mediated reactions.^{142,143} More energy input¹⁴⁴ and higher energy-coupling efficiency between matrices and analytes⁸⁹ enhance the ISD process. Because ISD ions receive the full acceleration energy in the ion source, they can be well resolved in the mass spectra. On the other hand, PSD ions are typically generated on the time scale of micro- to milliseconds in the field-free region. Owing to having the same velocity as their precursor ions, PSD ions cannot be resolved in linear TOF instruments.¹³⁷ In order to resolve PSD ions, which are usually buried into the background in linear TOF instruments, it is necessary

Fig. 5. Workflow for “on-bead methyl esterification” of sialic acids for immobilization and recovery of oligosaccharides. Reprinted with permission from ref. 133, John Wiley & Sons, Inc. (2007).

to analyze such ions by sweeping reflector potentials in reflectron TOF instruments. CID ions are generated through collisions of precursor ions with neutral gas molecules.¹⁴⁵⁾ During collision, redistribution of the kinetic energy of neutral gas molecules to the internal energy of precursor ions leads to fragmentation.¹⁴⁵⁾ A systematic nomenclature developed by Domon and Costello has become the common language for the community to specify fragmentation patterns of carbohydrates.¹⁴⁶⁾ Our goal for this section is to review some case studies of those fragmentation techniques in the structural analysis of carbohydrates. Readers are encouraged to refer to some of the more comprehensive reviews, which elucidate mechanisms of fragmentation techniques.^{89,137,141–143,147–151)}

Complementary fragmentation methods were utilized to gain the structural information of carbohydrates. Suzuki *et al.* and Jovanović *et al.* conducted comprehensive structural identifications of neutral oligosaccharides in the negative ion mode utilizing ISD, PSD, and CID.^{152,153)} Both groups reported that more information on linkage and branching, which were unobtainable using positive ion mode, could be acquired in the negative ion mode. Derivatization of carboxyl groups by methylamine using the condensing reagent (7-azabenzotriazol-1-yloxy)-tripyrrolidinophosphonium hexafluoro-phosphate (PyAOP) was applied by Nishikaze *et al.* to conduct glycoprofiling.¹⁵⁴⁾ They found that positive-ion CID of derivatized glycopeptides mainly provided information on peptide sequence and

glycan composition, whereas negative-ion CID provided in-depth structural information on glycan moieties. PSD and CID were combined by Lattova *et al.* to profile derivatized N-linked glycans from ovalbumin.¹⁵⁵⁾ Compared to MALDI-PSD, they found that more structural information from cross-ring cleavages (a- and x-type fragments) could be obtained using CID in tandem mass spectrometry. Laser-induced ISD combined with high-energy collision-induced dissociation (heCID) was reported by Wührer *et al.* to be a capable method for targeted structural identification of glycans.¹⁵⁰⁾ Another design that involved two consecutive time-of-flight (TOF/TOF) mass analyzers combined with a collision cell and a timed ion selector was developed by Suckau and coworkers.¹⁵⁶⁾ MALDI TOF/TOF was reported to yield more cross-ring cleavages that allow us to access the information of linkage positions.^{157,158)}

Akin to an ISD process, functionalized nanoparticles release charges and redistribute energy that facilitate fragmentation after absorbing laser energy.¹²²⁾ Our group synthesized DHB-functionalized mercury telluride nanoparticles (HgTe@DHB NPs) as a novel matrix, which induced both glycosidic and cross-ring cleavages in the solid phase. This method allowed us to do pseudo-MS/MS in a single step by regulating amounts of HgTe@DHB. Moreover, it preserved intact underivatized glycans. We have shown proof-of-concept for the novel matrix that permits the structural characterization of glycans.

Dissociation could also be initiated by absorption of

CONCLUSION AND PERSPECTIVES

REFERENCES

- Page 7 of 12

- 14) N. H. Packer, N. Karlsson. in *Glycomics: Methods and Protocols*, 1 Ed., Humana Press, 2009, pp. XIV, 390.
- 15) S. J. Danishefsky, Y. K. Shue, M. N. Chang, C. H. Wong. Development of globo-H cancer vaccine. *Acc. Chem. Res.* 48: 643–652, 2015.
- 16) S. J. North, P. G. Hitchen, S. M. Haslam, A. Dell. Mass spectrometry in the analysis of N-linked and O-linked glycans. *Curr. Opin. Struct. Biol.* 19: 498–506, 2009.
- 17) M. J. Kailemia, L. R. Ruhaak, C. B. Lebrilla, I. J. Amster. Oligosaccharide analysis by mass spectrometry: A review of recent developments. *Anal. Chem.* 86: 196–212, 2014.
- 18) W. R. Alley Jr., B. F. Mann, M. V. Novotny. High-sensitivity analytical approaches for the structural characterization of glycoproteins. *Chem. Rev.* 113: 2668–2732, 2013.
- 19) J. J. Kohler, S. M. Patrie. in *Mass Spectrometry of Glycoproteins Methods and Protocols*, 2013, p. 339.
- 20) N. H. Packer, N. G. Karlsson. in *Glycomics Methods and Protocols*, Humana Press, 2009.
- 21) D. Šagi, J. Peter-Katalinić. MALDI-MS of Glycans. MALDI MS. Wiley-VCH Verlag GmbH & Co. KGaA, 2007, pp. 181–214.
- 22) Y.-H. Lai, C.-C. Wang, C. W. Chen, B.-H. Liu, S. H. Lin, Y. T. Lee, Y.-S. Wang. Analysis of initial reactions of MALDI based on chemical properties of matrixes and excitation condition. *J. Phys. Chem. B* 116: 9635–9643, 2012.
- 23) W. Ens, Y. Mao, F. Mayer, K. G. Standing. Properties of matrix-assisted laser desorption measurements with a time-to-digital converter. *Rapid Commun. Mass Spectrom.* 5: 117–123, 1991.
- 24) C. D. Mowry, M. V. Johnston. Simultaneous detection of ions and neutrals produced by matrix-assisted laser desorption. *Rapid Commun. Mass Spectrom.* 7: 569–575, 1993.
- 25) A. P. Quist, T. Huth-Fehre, B. U. R. Sundqvist, A. Vertes. Total yield measurements in matrix-assisted laser desorption using a quartz crystal microbalance. *Rapid Commun. Mass Spectrom.* 8: 149–154, 1994.
- 26) A. A. Puretzky, D. B. Geohegan. Gas-phase diagnostics and LIF-imaging of 3-hydroxypicolinic acid MALDI-matrix plumes. *Chem. Phys. Lett.* 286: 425–432, 1998.
- 27) A. Moser, K. Range, D. M. York. Accurate proton affinity and gas-phase basicity values for molecules important in biocatalysis. *J. Phys. Chem. B* 114: 13911–13921, 2010.
- 28) A. G. Harrison. The gas-phase basicities and proton affinities of amino acids and peptides. *Mass Spectrom. Rev.* 16: 201–217, 1997.
- 29) P. D. Schnier, D. S. Gross, E. R. Williams. Electrostatic forces and dielectric polarizability of multiply protonated gas-phase cytochrome-*c* ions probed by ion/molecule chemistry. *J. Am. Chem. Soc.* 117: 6747–6757, 1995.
- 30) K. A. Jebber, K. Zhang, C. J. Cassidy, A. Chung-Phillips. *Ab initio* and experimental studies on the protonation of glucose in the gas phase. *J. Am. Chem. Soc.* 118: 10515–10524, 1996.
- 31) R. Knochenmuss. Ion formation mechanisms in UV-MALDI. *Analyst (Lond.)* 131: 966–986, 2006.
- 32) R. Knochenmuss, L. V. Zhigilei. Molecular dynamics simulations of MALDI: Laser fluence and pulse width dependence of plume characteristics and consequences for matrix and analyte ionization. *J. Mass Spectrom.* 45: 333–346, 2010.
- 33) M.-T. Tsai, S. Lee, I. C. Lu, K. Y. Chu, C.-W. Liang, C. H. Lee, Y. T. Lee, C.-K. Ni. Ion-to-neutral ratio of 2,5-dihydroxybenzoic acid in matrix-assisted laser desorption/ionization. *Rapid Commun. Mass Spectrom.* 27: 955–963, 2013.
- 34) Y. J. Bae, Y. S. Shin, J. H. Moon, M. S. Kim. Degree of ionization in MALDI of peptides: Thermal explanation for the gas-phase ion formation. *J. Am. Soc. Mass Spectrom.* 23: 1326–1335, 2012.
- 35) H. Ehring, M. Karas, F. Hillenkamp. Role of photoionization and photochemistry in ionization processes of organic molecules and relevance for matrix-assisted laser desorption/ionization mass spectrometry. *J. Mass Spectrom.* 27: 472–480, 1992.
- 36) V. Horneffer, K. Dreisewerd, H. C. Lüdemann, F. Hillenkamp, M. Läge, K. Strupat. Is the incorporation of analytes into matrix crystals a prerequisite for matrix-assisted laser desorption/ionization mass spectrometry? A study of five positional isomers of dihydroxybenzoic acid. *Int. J. Mass Spectrom.* 185–187: 859–870, 1999.
- 37) M. Glückmann, A. Pfenninger, R. Krüger, M. Thierolf, M. Karas, V. Horneffer, F. Hillenkamp, K. Strupat. Mechanisms in MALDI analysis: Surface interaction or incorporation of analytes? *Int. J. Mass Spectrom.* 210–211: 121–132, 2001.
- 38) V. Horneffer, M. Glückmann, B. Krüger, M. Karas, K. Strupat, F. Hillenkamp. Matrix-analyte-interaction in MALDI-MS: Pellet and nano-electrospray preparations. *Int. J. Mass Spectrom.* 249–250: 426–432, 2006.
- 39) X. Lou, J. L. J. van Dongen, L.-G. Milroy, E. W. Meijer. Generation of gas-phase ions from charged clusters: An important ionization step causing suppression of matrix and analyte ions in matrix-assisted laser desorption/ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* 30: 2628–2634, 2016.
- 40) J. Xu, T. Fujino. Stabilization of matrix molecules by nonvolatile analyte plays an important role in matrix-assisted laser desorption/ionization. *Chem. Phys. Lett.* 609: 6–10, 2014.
- 41) J. Yao, J. R. Scott, M. K. Young, C. L. Wilkins. Importance of matrix: Analyte ratio for buffer tolerance using 2,5-dihydroxybenzoic acid as a matrix in matrix-assisted laser desorption/ionization Fourier transform mass spectrometry and matrix-assisted laser desorption/ionization time of flight. *J. Am. Soc. Mass Spectrom.* 9: 805–813, 1998.
- 42) S. T. Tsai, C. H. Chen, Y. T. Lee, Y. S. Wang. Desorption dynamics of neutral molecules in matrix-assisted laser desorption/ionization. *Mol. Phys.* 106: 239–247, 2008.
- 43) H. Hazama, S. Furukawa, K. Awazu. Effect of solvent on ionization efficiency in matrix-assisted laser desorption/ionization mass spectrometry of peptides. *Chem. Phys.* 419: 196–199, 2013.
- 44) S. D. Hanton, P. A. Cornelio Clark, K. G. Owens. Investigations of matrix-assisted laser desorption/ionization sample preparation by time-of-flight secondary ion mass spectrometry. *J. Am. Soc. Mass Spectrom.* 10: 104–111, 1999.
- 45) S. Shimma, Y. Takashima, J. Hashimoto, K. Yonemori, K. Tamura, A. Hamada. Alternative two-step matrix application method for imaging mass spectrometry to avoid tissue shrinkage and improve ionization efficiency. *J. Mass Spectrom.* 48: 1285–1290, 2013.
- 46) I. D. Figueroa, O. Torres, D. H. Russell. Effects of the water content in the sample preparation for MALDI on the mass spectra. *Anal. Chem.* 70: 4527–4533, 1998.
- 47) V. Horneffer, R. Reichelt, K. Strupat. Protein incorporation into MALDI-matrix crystals investigated by high resolution field emission scanning electron microscopy. *Int. J. Mass Spectrom.* 226: 117–131, 2003.
- 48) K. Dreisewerd, S. Berkenkamp, A. Leisner, A. Rohlfling, C. Menzel. Fundamentals of matrix-assisted laser desorption/ionization mass spectrometry with pulsed infrared lasers. *Int. J. Mass Spectrom.* 226: 189–209, 2003.
- 49) H. Qiao, V. Spicer, W. Ens. The effect of laser profile, fluence, and spot size on sensitivity in orthogonal-injection matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* 22: 2779–2790, 2008.
- 50) O. Vorm, P. Roepstorff, M. Mann. Improved resolution and very high sensitivity in MALDI TOF of matrix surfaces made by fast evaporation. *Anal. Chem.* 66: 3281–3287, 1994.
- 51) T. W. Jaskolla, M. Karas, U. Roth, K. Steinert, C. Menzel, K. Reihs. Comparison between vacuum sublimed matrices and conventional dried droplet preparation in MALDI-TOF mass spectrometry. *J. Am. Soc. Mass Spectrom.* 20: 1104–1114, 2009.
- 52) Y. Chen, A. Vertes. Pumping rate and surface morphology dependence of ionization processes in matrix-assisted laser desorption/ionization. *J. Phys. Chem. A* 107: 9754–9761, 2003.
- 53) A. Westman, T. Huth-Fehre, P. Demirev, B. U. R. Sundqvist.

- Sample morphology effects in matrix-assisted laser-desorption/ionization mass spectrometry of proteins. *J. Mass Spectrom.* 30: 206–211, 1995.
- 54) P. Önnérjörd, S. Ekström, J. Bergquist, J. Nilsson, T. Laurell, G. Marko-Varga. Homogeneous sample preparation for automated high throughput analysis with matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry. *Rapid Commun. Mass Spectrom.* 13: 315–322, 1999.
- 55) T. I. Williams, D. A. Saggese, R. J. Wilcox, J. D. Martin, D. C. Muddiman. Effect of matrix crystal structure on ion abundance of carbohydrates by matrix-assisted laser desorption/ionization Fourier transform ion cyclotron resonance mass spectrometry. *Rapid Commun. Mass Spectrom.* 21: 807–811, 2007.
- 56) M. Wiegelmann, J. Soltwisch, T. W. Jaskolla, K. Dreisewerd. Matching the laser wavelength to the absorption properties of matrices increases the ion yield in UV-MALDI mass spectrometry. *Anal. Bioanal. Chem.* 405: 6925–6932, 2013.
- 57) P. C. Liao, J. Allison. Ionization processes in matrix-assisted laser desorption/ionization mass spectrometry: Matrix-dependent formation of $[M+H]^+$ vs. $[M+Na]^+$ ions of small peptides and some mechanistic comments. *J. Mass Spectrom.* 30: 408–423, 1995.
- 58) M. Hatakeyama, M. Tachikawa. *Ab initio* quantum chemical study on the mechanism of exceptional behavior of lysine for ion yields in MALDI—Role of vibrational entropic contribution in thermally averaged proton affinities. *J. Mass Spectrom.* 46: 376–382, 2011.
- 59) K. Breuker, R. Knochenmuss, J. Zhang, A. Stortelder, R. Zenobi. Thermodynamic control of final ion distributions in MALDI: In-plume proton transfer reactions. *Int. J. Mass Spectrom.* 226: 211–222, 2003.
- 60) R. Knochenmuss, A. Stortelder, K. Breuker, R. Zenobi. Secondary ion–molecule reactions in matrix-assisted laser desorption/ionization. *J. Mass Spectrom.* 35: 1237–1245, 2000.
- 61) F. Hillenkamp, E. Wäfler, M. C. Jecklin, R. Zenobi. Positive and negative analyte ion yield in matrix-assisted laser desorption/ionization revisited. *Int. J. Mass Spectrom.* 285: 114–119, 2009.
- 62) R. Knochenmuss. A quantitative model of ultraviolet matrix-assisted laser desorption/ionization. *J. Mass Spectrom.* 37: 867–877, 2002.
- 63) R. Knochenmuss. MALDI mechanisms: Wavelength and matrix dependence of the coupled photophysical and chemical dynamics model. *Analyst (Lond.)* 139: 147–156, 2014.
- 64) L. Molin, R. Seraglia, Z. Czarnocki, J. K. Maurin, F. A. Pluciński, P. Traldi. On the primary ionization mechanism(s) in matrix-assisted laser desorption ionization. *J. Anal. Methods Chem.* 2012: 161865, 2012.
- 65) K. M. Kirmess, R. Knochenmuss, G. J. Blanchard, G. R. Kinsel. MALDI ionization mechanisms investigated by comparison of isomers of dihydroxybenzoic acid. *J. Mass Spectrom.* 51: 79–85, 2016.
- 66) S. F. Niu, W. Z. Zhang, B. T. Chait. Direct comparison of infrared and ultraviolet wavelength matrix-assisted laser desorption/ionization mass spectrometry of proteins. *J. Am. Soc. Mass Spectrom.* 9: 1–7, 1998.
- 67) X. J. Chen, J. A. Carroll, R. C. Beavis. Near-ultraviolet-induced matrix-assisted laser desorption/ionization as a function of wavelength. *J. Am. Soc. Mass Spectrom.* 9: 885–891, 1998.
- 68) Y.-H. Lai, B.-G. Chen, Y. T. Lee, Y.-S. Wang, S. H. Lin. Contribution of thermal energy to initial ion production in matrix-assisted laser desorption/ionization observed with 2,4,6-trihydroxyacetophenone. *Rapid Commun. Mass Spectrom.* 28: 1716–1722, 2014.
- 69) Y.-H. Lai, C.-C. Wang, S.-H. Lin, Y.-T. Lee, Y.-S. Wang. Solid-phase thermodynamic interpretation of ion desorption in matrix-assisted laser desorption/ionization. *J. Phys. Chem. B* 114: 13847–13852, 2010.
- 70) K. Y. Chu, S. Lee, M. T. Tsai, I. C. Lu, Y. A. Dyakov, Y. H. Lai, Y. T. Lee, C. K. Ni. Thermal proton transfer reactions in ultraviolet matrix-assisted laser desorption/ionization. *J. Am. Soc. Mass Spectrom.* 25: 310–318, 2014.
- 71) I. C. Lu, C. Lee, Y. T. Lee, C. K. Ni. Ionization mechanism of matrix-assisted laser desorption/ionization. In: Annual Review of Analytical Chemistry Vol. 8 (Ed.: R. G. Cooks, J. E. Pemberton), 2015, pp. 21–39.
- 72) J. H. Moon, S. Yoon, Y. J. Bae, M. S. Kim. Formation of gas-phase peptide ions and their dissociation in MALDI: Insights from kinetic and ion yield studies. *Mass Spectrom. Rev.* 34: 94–115, 2015.
- 73) S. H. Ahn, K. M. Park, Y. J. Bae, M. S. Kim. Quantitative reproducibility of mass spectra in matrix-assisted laser desorption ionization and unraveling of the mechanism for gas-phase peptide ion formation. *J. Mass Spectrom.* 48: 299–305, 2013.
- 74) Y. J. Bae, M. S. Kim. A thermal mechanism of ion formation in MALDI. In: Annual Review of Analytical Chemistry, Vol. 8 (Ed.: R. G. Cooks, J. E. Pemberton), 2015, pp. 41–60.
- 75) B. H. Liu, O. P. Charkin, N. Klemenko, C. W. Chen, Y. S. Wang. Initial ionization reaction in matrix-assisted laser desorption/ionization. *J. Phys. Chem. B* 114: 10853–10859, 2010.
- 76) N. L. Asfandiarov, S. A. Pshenichnyuk, A. I. Fokin, V. G. Lukin, V. S. Fal'ko. Electron capture negative ion mass spectra of some typical matrix-assisted laser desorption/ionization matrices. *Rapid Commun. Mass Spectrom.* 16: 1760–1765, 2002.
- 77) E. Alonso, R. Zenobi. Non-linear photoelectron effect contributes to the formation of negative matrix ions in UV-MALDI. *Phys. Chem. Chem. Phys.* 18: 19574–19587, 2016.
- 78) M. Karas, R. Krüger. Ion formation in MALDI: The cluster ionization mechanism. *Chem. Rev.* 103: 427–439, 2003.
- 79) T. W. Jaskolla, M. Karas. Compelling evidence for lucky survivor and gas phase protonation: The unified MALDI analyte protonation mechanism. *J. Am. Soc. Mass Spectrom.* 22: 976–988, 2011.
- 80) K. Dreisewerd. The desorption process in MALDI. *Chem. Rev.* 103: 395–425, 2003.
- 81) M. W. Little, J. Laboy, K. K. Murray. Wavelength dependence of soft infrared laser desorption and ionization. *J. Phys. Chem. C* 111: 1412–1416, 2007.
- 82) J. Soltwisch, T. W. Jaskolla, K. Dreisewerd. Color matters—Material ejection and ion yields in UV-MALDI mass spectrometry as a function of laser wavelength and laser fluence. *J. Am. Soc. Mass Spectrom.* 24: 1477–1488, 2013.
- 83) E. Nordhoff, A. Ingendoh, R. Cramer, A. Overberg, B. Stahl, M. Karas, F. Hillenkamp, P. F. Crain, B. Chait. Matrix-assisted laser desorption ionization mass-spectrometry of nucleic-acids with wavelengths in the ultraviolet and infrared. *Rapid Commun. Mass Spectrom.* 6: 771–776, 1992.
- 84) J. Soltwisch, T. W. Jaskolla, F. Hillenkamp, M. Karas, K. Dreisewerd. Ion yields in UV-MALDI mass spectrometry as a function of excitation laser wavelength and optical and physico-chemical properties of classical and halogen-substituted MALDI matrixes. *Anal. Chem.* 84: 6567–6576, 2012.
- 85) K. Dreisewerd, M. Schürenberg, M. Karas, F. Hillenkamp. Matrix-assisted laser desorption/ionization with nitrogen lasers of different pulse widths. *Int. J. Mass Spectrom. Ion Process.* 154: 171–178, 1996.
- 86) K. Dreisewerd, M. Schürenberg, M. Karas, F. Hillenkamp. Influence of the laser intensity and spot size on the desorption of molecules and ions in matrix-assisted laser desorption/ionization with a uniform beam profile. *Int. J. Mass Spectrom. Ion Process.* 141: 127–148, 1995.
- 87) C. D. Mowry, M. V. Johnston. Internal energy of neutral molecules ejected by matrix-assisted laser desorption. *J. Phys. Chem.* 98: 1904–1909, 1994.
- 88) G. H. Luo, I. Marginean, A. Vertes. Internal energy of ions generated by matrix-assisted laser desorption/ionization. *Anal. Chem.* 74: 6185–6190, 2002.

- 89) V. Gabelica, E. Schulz, M. Karas. Internal energy build-up in matrix-assisted laser desorption/ionization. *J. Mass Spectrom.* 39: 579–593, 2004.
- 90) Y. J. Bae, J. H. Moon, M. S. Kim. Expansion cooling in the matrix plume is under-recognized in MALDI mass spectrometry. *J. Am. Soc. Mass Spectrom.* 22: 1070–1078, 2011.
- 91) A. Koubenakis, V. Frankevich, J. Zhang, R. Zenobi. Time-resolved surface temperature measurement of MALDI matrices under pulsed UV laser irradiation. *J. Phys. Chem. A* 108: 2405–2410, 2004.
- 92) E. Schulz, M. Karas, F. Rosu, V. Gabelica. Influence of the matrix on analyte fragmentation in atmospheric pressure MALDI. *J. Am. Soc. Mass Spectrom.* 17: 1005–1013, 2006.
- 93) R. E. Johnson. Models for matrix-assisted desorption by a laser-pulse. *Int. J. Mass Spectrom.* 139: 25–38, 1994.
- 94) M. Schürenberg, K. Dreisewerd, S. Kamanabrou, F. Hillenkamp. Influence of the sample temperature on the desorption of matrix molecules and ions in matrix-assisted laser desorption ionization. *Int. J. Mass Spectrom. Ion Process.* 172: 89–94, 1998.
- 95) L. V. Zhigilei, E. Leveugle, B. J. Garrison, Y. G. Yingling, M. I. Zeifman. Computer simulations of laser ablation of molecular substrates. *Chem. Rev.* 103: 321–347, 2003.
- 96) R. Knochenmuss, L. V. Zhigilei. What determines MALDI ion yields? A molecular dynamics study of ion loss mechanisms. *Anal. Bioanal. Chem.* 402: 2511–2519, 2012.
- 97) B. Stahl, M. Steup, M. Karas, F. Hillenkamp. Analysis of neutral oligosaccharides by matrix-assisted laser desorption ionization mass spectrometry. *Anal. Chem.* 63: 1463–1466, 1991.
- 98) E. Botek, J. L. Debrun, B. Hakim, L. Morin-Allory. Attachment of alkali cations on β -D-glucopyranose: Matrix-assisted laser desorption/ionization time-of-flight studies and *ab initio* calculations. *Rapid Commun. Mass Spectrom.* 15: 273–276, 2001.
- 99) M. D. Mohr, K. O. Bornsen, H. M. Widmer. Matrix-assisted laser desorption/ionization mass spectrometry: Improved matrix for oligosaccharides. *Rapid Commun. Mass Spectrom.* 9: 809–814, 1995.
- 100) S. P. Mirza, N. P. Raju, M. Vairamani. Estimation of the proton affinity values of fifteen matrix-assisted laser desorption/ionization matrices under electrospray ionization conditions using the kinetic method. *J. Am. Soc. Mass Spectrom.* 15: 431–435, 2004.
- 101) J.-L. Chen, C. Lee, I.-C. Lu, C.-L. Chien, Y.-T. Lee, W.-P. Hu, C.-K. Ni. Theoretical investigation of low detection sensitivity for underivatized carbohydrates in ESI and MALDI. *J. Mass Spectrom.* 51: 1180–1186, 2016.
- 102) G. Ohanessian. Interaction of MALDI matrix molecules with Na^+ in the gas phase. *Int. J. Mass Spectrom.* 219: 577–592, 2002.
- 103) C. Lee, I. C. Lu, C. Hsu, H.-Y. Lin, S.-P. Liang, Y.-T. Lee, C.-K. Ni. Formation of metal-related ions in matrix-assisted laser desorption ionization. *J. Am. Soc. Mass Spectrom.* 27: 1491–1498, 2016.
- 104) A. Antonopoulos, P. Bonnet, E. Botek, J. L. Debrun, B. Hakim, B. Herbreteau, L. Morin-Allory. Study of the attachment of Na^+ on glucose and on some of its methylated derivatives. *Rapid Commun. Mass Spectrom.* 17: 122–125, 2003.
- 105) K. Fukui, A. Kameyama, Y. Mukai, K. Takahashi, N. Ikeda, Y. Akiyama, H. Narimatsu. A computational study of structure–reactivity relationships in Na-adduct oligosaccharides in collision-induced dissociation reactions. *Carbohydr. Res.* 341: 624–633, 2006.
- 106) M. T. Cancilla, S. G. Penn, J. A. Carroll, C. B. Lebrilla. Coordination of alkali metals to oligosaccharides dictates fragmentation behavior in matrix assisted laser desorption ionization Fourier transform mass spectrometry. *J. Am. Chem. Soc.* 118: 6736–6745, 1996.
- 107) Y. Fukuyama, S. Nakaya, Y. Yamazaki, K. Tanaka. Ionic liquid matrices optimized for MALDI-MS of sulfated/sialylated/neutral oligosaccharides and glycopeptides. *Anal. Chem.* 80: 2171–2179, 2008.
- 108) T. Nishikaze, K. Kaneshiro, S. Kawabata, K. Tanaka. Structural analysis of N-glycans by the glycan-labeling method using 3-aminoquinoline-based liquid matrix in negative-ion MALDI-MS. *Anal. Chem.* 84: 9453–9461, 2012.
- 109) Y. Fukuyama, N. Funakoshi, K. Takeyama, Y. Hioki, T. Nishikaze, K. Kaneshiro, S. Kawabata, S. Iwamoto, K. Tanaka. 3-Aminoquinoline/*p*-coumaric acid as a MALDI matrix for glycopeptides, carbohydrates, and phosphopeptides. *Anal. Chem.* 86: 1937–1942, 2014.
- 110) M. Kuang, Y. Zhang, P. Yang, H. Lu. Novel ionic matrices for enhanced ionization of oligosaccharides/glycopeptides during MALDI-MS analysis. *Acta Chim. Sin.* 71: 1007–1010, 2013.
- 111) L. Witt, A. Pirkel, F. Draude, J. Peter-Katalinic, K. Dreisewerd, M. Mormann. Water ice is a soft matrix for the structural characterization of glycosaminoglycans by infrared matrix-assisted laser desorption/ionization. *Anal. Chem.* 86: 6439–6446, 2014.
- 112) C.-W. Liang, P.-J. Chang, Y.-J. Lin, Y.-T. Lee, C.-K. Ni. High ion yields of carbohydrates from frozen solution by UV-MALDI. *Anal. Chem.* 84: 3493–3499, 2012.
- 113) C. L. Wu, C. C. Wang, Y. H. Lai, H. Lee, J. D. Lin, Y. T. Lee, Y. S. Wang. Selective enhancement of carbohydrate ion abundances by diamond nanoparticles for mass spectrometric analysis. *Anal. Chem.* 85: 3836–3841, 2013.
- 114) T. Yamagaki, H. Suzuki, K. Tachibana. Solid-phase fluorescence and ionization efficiency in negative-ion matrix-assisted laser desorption/ionization of neutral oligosaccharides: Interaction between beta-carboline matrix and ammonium salt. *J. Am. Soc. Mass Spectrom.* 18: 714–723, 2007.
- 115) P. Domann, D. I. R. Spencer, D. J. Harvey. Production and fragmentation of negative ions from neutral N-linked carbohydrates ionized by matrix-assisted laser desorption/ionization. *Rapid Commun. Mass Spectrom.* 26: 469–479, 2012.
- 116) W. Gao, G. Ou, X. Feng, B.-F. Liu, H. Zhang, X. Liu. Matrix-assisted laser desorption/ionization mass spectrometry analysis of glycans with co-derivatization of asparaginyl-oligosaccharides. *Anal. Chim. Acta* 896: 102–110, 2015.
- 117) R. P. Obena, M.-C. Tseng, I. Primadona, J. Hsiao, I. C. Li, R. Y. Capangpangan, H.-F. Lu, W.-S. Li, I. Chao, C.-C. Lin, Y.-J. Chen. UV-activated multilayer nanomatrix provides one-step tunable carbohydrate structural characterization in MALDI-MS. *Chem. Sci. (Camb.)* 6: 4790–4800, 2015.
- 118) Y. Gholipour, S. L. Giudicessi, H. Nonami, R. Erra-Balsells. Diamond, titanium dioxide, titanium silicon oxide, and barium strontium titanium oxide nanoparticles as matrices for direct matrix-assisted laser desorption/ionization mass spectrometry analysis of carbohydrates in plant tissues. *Anal. Chem.* 82: 5518–5526, 2010.
- 119) H.-J. Yang, A. Lee, M.-K. Lee, W. Kim, J. Kim. Detection of small neutral carbohydrates using various supporting materials in laser desorption/ionization mass spectrometry. *Bull. Korean Chem. Soc.* 31: 35–40, 2010.
- 120) Y. Sahashi, I. Osaka, S. Taira. Nutrition analysis by nanoparticle-assisted laser desorption/ionisation mass spectrometry. *Food Chem.* 123: 865–871, 2010.
- 121) C.-L. Su, W.-L. Tseng. Gold nanoparticles as assisted matrix for determining neutral small carbohydrates through laser desorption/ionization time-of-flight mass spectrometry. *Anal. Chem.* 79: 1626–1633, 2007.
- 122) I. Primadona, Y.-H. Lai, R. Y. Capangpangan, R. P. Obena, M.-C. Tseng, M.-F. Huang, H.-T. Chang, S.-T. Li, C.-Y. Wu, W.-T. Chien, C.-C. Lin, Y.-S. Wang, Y.-J. Chen. Functionalized HgTe nanoparticles promote laser-induced solid phase ionization/dissociation for comprehensive glycan sequencing. *Analyst (Lond.)* 141: 6093–6103, 2016.
- 123) H. N. Abdelhamid. Organic matrices, ionic liquids, and organic matrices@nanoparticles assisted laser desorption/ionization mass spectrometry. *Trends Analyt. Chem.* 89: 68–98, 2017.

- 124) I. Ciucanu, F. Kerek. A simple and rapid method for the permethylation of carbohydrates. *Carbohydr. Res.* 131: 209–217, 1984.
- 125) E. Lattova, H. Perreault. Labelling saccharides with phenylhydrazine for electrospray and matrix-assisted laser desorption/ionization mass spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* 793: 167–179, 2003.
- 126) S. Sekiya, Y. Wada, K. Tanaka. Derivatization for stabilizing sialic acids in MALDI-MS. *Anal. Chem.* 77: 4962–4968, 2005.
- 127) S. I. Hakomori. A rapid permethylation of glycolipid, and polysaccharide catalyzed by methylsulfinyl carbanion in dimethyl sulfoxide. *J. Biochem.* 55: 205–208, 1964.
- 128) M. Toyoda, H. Ito, Y. Matsuno, H. Narimatsu, A. Kameyama. Quantitative derivatization of sialic acids for the detection of sialoglycans by MALDI MS. *Anal. Chem.* 80: 5211–5218, 2008.
- 129) E. Lattova, H. Perreault. The usefulness of hydrazine derivatives for mass spectrometric analysis of carbohydrates. *Mass Spectrom. Rev.* 32: 366–385, 2013.
- 130) Y. Mechref, Y. Hu, J. L. Desantos-Garcia, A. Hussein, H. Tang. Quantitative glycomics strategies. *Mol. Cell. Proteomics* 12: 874–884, 2013.
- 131) Y. Wada. Glycan profiling: Label-free analysis of glycoproteins. In: *Mass Spectrometry of Glycoproteins Methods and Protocols* (Ed.: J. J. Kohler, S. M. Patrie), Humana Press, 2013, pp. 245–253.
- 132) W. J. Gao, H. H. Li, Y. Y. Liu, Y. H. Liu, X. J. Feng, B. F. Liu, X. Liu. Rapid and sensitive analysis of N-glycans by MALDI-MS using permanent charge derivatization and methylamidation. *Talanta* 161: 554–559, 2016.
- 133) Y. Miura, Y. Shinohara, J. Furukawa, N. Nagahori, S.-I. Nishimura. Rapid and simple solid-phase esterification of sialic acid residues for quantitative glycomics by mass spectrometry. *Chemistry* 13: 4797–4804, 2007.
- 134) K. R. Reidling, D. Blank, D. M. Kuijper, A. M. Deelder, M. Wührer. High-throughput profiling of protein N-glycosylation by MALDI-TOF-MS employing linkage-specific sialic acid esterification. *Anal. Chem.* 86: 5784–5793, 2014.
- 135) P. Kang, Y. Mechref, I. Klouckova, M. V. Novotny. Solid-phase permethylation of glycans for mass spectrometric analysis. *Rapid Commun. Mass Spectrom.* 19: 3421–3428, 2005.
- 136) R. S. Brown, J. J. Lennon. Sequence-specific fragmentation of matrix-assisted laser-desorbed protein peptide ions. *Anal. Chem.* 67: 3990–3999, 1995.
- 137) B. Spengler. Post-source decay analysis in matrix-assisted laser desorption/ionization mass spectrometry of biomolecules. *J. Mass Spectrom.* 32: 1019–1036, 1997.
- 138) K. Tseng, J. L. Hedrick, C. B. Lebrilla. Catalog-library approach for the rapid and sensitive structural elucidation of oligosaccharides. *Anal. Chem.* 71: 3747–3754, 1999.
- 139) S. König, J. A. Leary. Evidence for linkage position determination in cobalt coordinated pentasaccharides using ion trap mass spectrometry. *J. Am. Soc. Mass Spectrom.* 9: 1125–1134, 1998.
- 140) D. J. Harvey, R. H. Bateman, M. R. Green. High-energy collision-induced fragmentation of complex oligosaccharides ionized by matrix-assisted laser desorption/ionization mass spectrometry. *J. Mass Spectrom.* 32: 167–187, 1997.
- 141) D. Asakawa. Principles of hydrogen radical mediated peptide/protein fragmentation during matrix-assisted laser desorption/ionization mass spectrometry. *Mass Spectrom. Rev.* 35: 535–556, 2016.
- 142) R. A. Zubarev, N. L. Kelleher, F. W. McLafferty. Electron capture dissociation of multiply charged protein cations. A nonergodic process. *J. Am. Chem. Soc.* 120: 3265–3266, 1998.
- 143) D. Suckau, A. Resemann. T³-Sequencing: Targeted characterization of the N- and C-termini of undigested proteins by mass spectrometry. *Anal. Chem.* 75: 5817–5824, 2003.
- 144) H. Yang, Y. Yu, F. Song, S. Liu. Structural characterization of neutral oligosaccharides by laser-enhanced in-source decay of MALDI-FTICR MS. *J. Am. Soc. Mass Spectrom.* 22: 845–855, 2011.
- 145) L. Sleno, D. A. Volmer. Ion activation methods for tandem mass spectrometry. *J. Mass Spectrom.* 39: 1091–1112, 2004.
- 146) B. Domon, C. E. Costello. A systematic nomenclature for carbohydrate fragmentations in FAB-MS/MS spectra of glycoconjugates. *Glycoconj. J.* 5: 397–409, 1988.
- 147) D. Asakawa, N. Smargiasso, E. De Pauw. Identification and relative-quantification of glycans by matrix-assisted laser desorption/ionization in-source decay with hydrogen abstraction. *Anal. Chem.* 84: 7463–7468, 2012.
- 148) H. Yang, M. Li, Z. Li, S. Liu. Gas-phase fragmentation of oligosaccharides in MALDI laser-enhanced in-source decay induced by thermal hydrogen radicals. *Analyst (Lond.)* 137: 3624–3626, 2012.
- 149) N. Smargiasso, E. De Pauw. Optimization of matrix conditions for the control of MALDI in-source decay of permethylated glycans. *Anal. Chem.* 82: 9248–9253, 2010.
- 150) M. Wührer, A. M. Deelder. Matrix-assisted laser desorption/ionization in-source decay combined with tandem time-of-flight mass spectrometry of permethylated oligosaccharides: Targeted characterization of specific parts of the glycan structure. *Rapid Commun. Mass Spectrom.* 20: 943–951, 2006.
- 151) R. Knochenmuss, R. Zenobi. MALDI ionization: The role of in-plume processes. *Chem. Rev.* 103: 441–452, 2003.
- 152) H. Suzuki, T. Yamagaki, K. Tachibana. Fragmentation of neutral oligosaccharides in negative-ion MALDI mass spectrometry. *Trends Glycosci. Glycotechnol.* 18: 277–292, 2006.
- 153) M. Jovanović, J. Peter-Katalinić. Negative ion MALDI-TOF MS, ISD and PSD of neutral underivatized oligosaccharides without anionic dopant strategies, using 2,5-DHAP as a matrix. *J. Mass Spectrom.* 51: 111–122, 2016.
- 154) T. Nishikaze, S. Kawabata, K. Tanaka. In-depth structural characterization of N-linked glycopeptides using complete derivatization for carboxyl groups followed by positive- and negative-ion tandem mass spectrometry. *Anal. Chem.* 86: 5360–5369, 2014.
- 155) E. Lattova, H. N. Perreault, O. Krokhin. Matrix-assisted laser desorption/ionization tandem mass spectrometry and post-source decay fragmentation study of phenylhydrazones of N-linked oligosaccharides from ovalbumin. *J. Am. Soc. Mass Spectrom.* 15: 725–735, 2004.
- 156) D. Suckau, A. Resemann, M. Schuerenberg, P. Hufnagel, J. Franzen, A. Holle. A novel MALDI LIFT-TOF/TOF mass spectrometer for proteomics. *Anal. Bioanal. Chem.* 376: 952–965, 2003.
- 157) W. Morelle, M. C. Slomianny, H. Diemer, C. Schaeffer, A. van Dorselaer, J. C. Michalski. Fragmentation characteristics of permethylated oligosaccharides using a matrix-assisted laser desorption/ionization two-stage time-of-flight (TOF/TOF) tandem mass spectrometer. *Rapid Commun. Mass Spectrom.* 18: 2637–2649, 2004.
- 158) E. Spina, L. Sturiale, D. Romeo, G. Impallomeni, D. Garozzo, D. Waidelich, M. Glueckmann. New fragmentation mechanisms in matrix-assisted laser desorption/ionization time-of-flight/time-of-flight tandem mass spectrometry of carbohydrates. *Rapid Commun. Mass Spectrom.* 18: 392–398, 2004.
- 159) J. S. Brodbelt. Photodissociation mass spectrometry: New tools for characterization of biological molecules. *Chem. Soc. Rev.* 43: 2757–2783, 2014.
- 160) H. J. An, C. B. Lebrilla. Structure elucidation of native N- and O-linked glycans by tandem mass spectrometry (tutorial). *Mass Spectrom. Rev.* 30: 560–578, 2011.
- 161) D. Goldberg, M. Bern, B. S. Li, C. B. Lebrilla. Automatic determination of O-glycan structure from fragmentation spectra. *J. Proteome Res.* 5: 1429–1434, 2006.
- 162) B. J. Ko, J. S. Brodbelt. 193 nm ultraviolet photodissociation of deprotonated sialylated oligosaccharides. *Anal. Chem.* 83: 8192–8200, 2011.

- 163) T. Yamagaki, H. Suzuki, K. Tachibana. A comparative study of the fragmentation of neutral lactooligosaccharides in negative-ion mode by UV-MALDI-TOF and UV-MALDI ion-trap/TOF mass spectrometry. *J. Am. Soc. Mass Spectrom.* 17: 67–74, 2006.
- 164) T. Yamagaki, H. Suzuki, K. Tachibana. In-source and post-source decay in negative-ion matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of neutral oligosaccharides. *Anal. Chem.* 77: 1701–1707, 2005.