〔原著論文〕 Downregulation of Major Histocompatibility Complex Class I Expression in Polymorphonuclear Leukocytes by Triggering CD16 Antigen: Different Effects of Two Clones of anti-Fc γ RIII Monoclonal Antibody

Motoko Nishimura¹⁾, Yoshihide Ishikawa¹⁾, Shigeki Mitsunaga²⁾

¹⁾Department of Research, Tokyo Metropolitan Red Cross, Blood Center, Tokyo, Japan

²⁾Department of Research, The Japanese Red Cross, Central Blood Center, Tokyo, Japan

(Received January 17, 2002)

Summary

Polymorphonuclear leukocytes (PMNs) constitutively express major histocompatibility complex (MHC) class I molecules on their surface. In spite of the prominent roles of PMNs in host immune defense, the roles of MHC class I molecules in PMNs, such as presenting intracellularly processed antigens to CD8⁺ T cells, or preventing the cytotoxic effect of natural killer (NK) cells by their upregulated expression levels, have not been fully investigated thus far. PMNs express two low-affinity Fc γ receptors, Fc γ R II (CD32) and Fc γ R III (CD16). The intracellular signaling pathway of PMNs is activated by triggering or cross-linking CD16, resulted in the upregulation of MHC class I expression. We found that the expression levels of MHC class I molecules of PMNs could be downregulated by triggering CD16, but not by triggering CD32.

Keywords : MHC class I, PMN, Fc γ Rs, Anti-CD16 mAb, NA1 / NA2 antigens.

Introduction

Polymorphonuclear leukocytes (PMNs) play important roles in host immune defense through phagocytosis, clearance of immune complexes and production of reactive oxygen species (ROS), and induce phagocytic inflammatory responses against a variety of microbial products. Inflammatory responses involve adhesion molecule and chemokine receptor upregulation, and calcium release by cross-linking β 2 integrin, CD11b

and CD18. PMNs constitutively express major histocompatibility complex (MHC) class I molecules on their surface. MHC class I molecules present intracellularly processed nonself antigens to CD8⁺ T cells, such as microorgans, and tumor-derived peptides. A recent report indicated that MHC class II molecules observed on the surface of PMNs activated by lipopolysaccharide (LPS) or granulocyte-macrophage colony stimulating factor (GM-CSF) may play a role in

Corresponding author : **Motoko Nishimura, Ph.D.**
Department of Research, Tokyo Metropolitan Red Cross, Blood Center
4-1-31 Hiroo, Shibuya-ku, Tokyo 150-0012

Phone: 03-3406-1211
Fax: 03-3406-7892
E-mail:

presenting antigens to T cells (1, 2). On the other hand, the roles of MHC class I on the PMN cell surface have not been fully investigated thus far. PMNs constitutively express two low-affinity receptors for the Fc fragments of IgG (Fc γ R II and Fc γ R III) that couple humoral and cellular immune responses by triggering immune complexes (3, 4, 5). By triggering or cross-linking, Fc γ Rs easily activate the intracellular signal transduction pathway. Our study shows that downregulation of MHC class I molecules in PMNs may be triggered by Fc γ R III (CD16) in the presence of anti-CD16 monoclonal antibodies (mAbs), but not by Fc γ R II (CD32) in the presence of anti-CD32 mAb.

Materials and Methods

Cell culture

PMNs were prepared from the citrated anti-coagulant treated peripheral bloods samples from several healthy individuals using the conventional method with Ficoll density gradient centrifugation at 1,700 rpm for 20 minutes. The lower layer PMNs of red blood cells was harvested, followed by red blood cell lysis with NH_4Cl (6). The purity of the obtained PMNs was $98.8 \pm 3.2\%$ [mean $\pm$ standard deviation (SD)] in 10 randomly selected results from 123 separate PMN preparations, assessed by staining the cells with fluorescence isothiocyanate (FITC)-conjugated mouse anti-human neutrophil-specific mAb, namely, anti-CD16b mAb (clone 1D3: Immunotec, France), and by flow cytometry (FCM). The viability of the PMNs was $>98\%$ as assessed by trypan blue dye exclusion. PMNs were cultured in the RPMI 1640 medium supplemented with 10% heat-inactivated (56°C for 30 minutes) fetal calf serum (FCS: HyClone, UT, USA) in the presence or absence of 200-fold-diluted anti-CD16 mAbs, clone OKNK (IgM, Ortho Diagnostic Systems K. K., Japan) or clone 3G8 (IgG₁, BD Pharmingen, CA, USA) or anti-CD32 mAb (Immunotec, France) at 37°C for 3–48 hours at a cell concentration of 2×10^6 cells mL^{-1} .

Immunofluorescence analysis

Freshly prepared PMNs cultured for three, eight or 24

hours in the presence or absence of anti-CD16 mAbs, clone OKNK, clone 3G8 or anti-CD32 mAb, were harvested and stained with 40-fold-diluted phycoerythrin (PE)-conjugated mouse anti-human MHC class I (clone W6/32: Serotec, UK) at 4°C for 30 minutes. These cells were then washed twice with FCM buffer [phosphate-buffered saline (PBS) + 0.2% bovine serum albumin (BSA) + 0.1% NaN_3], followed by FCM using Cytron (Ortho Diagnostic Systems K. K., Japan). In the case of staining with PE-conjugated anti-CD45 (DAKO, Denmark), anti-CD32 or anti-CD14 mAbs (DAKO, Denmark), we used essentially the same method as that used in staining MHC class I molecules from freshly prepared PMNs or 24-hour-cultured PMNs in the presence of 200-fold-diluted anti-CD16 mAbs, OKNK or 3G8. In another experiment, freshly prepared PMNs were subjected to direct staining with 40-fold-diluted FITC-conjugated anti-CD16 mAbs, clone OKNK or clone 3G8, at 4°C for 30 minutes, and then washed twice with FCM buffer. Staining profiles were also analyzed using Cytron.

Competitive treatment of two clones of anti-CD16 mAb

Freshly prepared PMNs were primarily treated with 40-fold-diluted mAbs, clone OKNK or clone 3G8, at 4°C for 30 minutes. After washing twice with FCM buffer, 40-fold-diluted FITC-conjugated clone OKNK was added to clone 3G8-pretreated PMNs. In contrast, 40-fold-diluted FITC-conjugated clone 3G8 was added to clone OKNK-pretreated PMNs. After staining at 4°C for 30 minutes, PMNs were washed twice with FCM buffer and staining profiles were analyzed with Cytron. As negative control, FITC-conjugated mouse IgG (DAKO) was added to PMNs that were pretreated with clone OKNK or clone 3G8.

Assessment of cytotoxicity against PMNs by allogeneic peripheral blood mononuclear cells (PBMCs)

The assessment of PMN susceptibility to cytotoxic activity of PBMCs was performed according to the conventional ^{51}Cr -release assay. PMNs were cultured for 24 hours in the presence or absence of 200-fold-diluted clone OKNK at a cell concentration of $2 \times 10^6/\text{ml}$ in the

RPMI 1640 medium supplemented with 10% FCS. Thereafter, PMNs were labeled with ^{51}Cr at $50 \mu\text{Ci}/1 \times 10^6$ cells at 37°C for one hour. After the incubation period, PMNs were washed three times with the RPMI 1640 medium supplemented with 1% FCS and used as target cells. Effector allogeneic PBMCs were prepared from the blood of several healthy individuals by Ficoll gradient centrifugation, obtaining the mononuclear cell layer that was washed three times with the RPMI 1640 medium supplemented with 1% FCS. Target cells were inoculated in 96-well round-bottomed plates (1×10^6 cells/well, $100 \mu\text{l}$ of the RPMI 1640 medium supplemented with 10% FCS. Then effector cells were added at a target : effector ratios of 1:10, 1:20, 1:40 and 1:80 at $100 \mu\text{l}$ /well with containing the RPMI 1640 medium supplemented with 10% FCS. After centrifugation at 1,000 rpm for 10 minutes to facilitate contact between target and effector cells, plates were incubated at 37°C for six hours. Then centrifugation was performed at 2,000 rpm for 10 minutes; $100 \mu\text{l}$ of culture supernatants was harvested and radioactivities were counted with a γ counter. Specific percentage lysis was calculated using the following formula: $(\text{cpm of samples} - \text{cpm of spontaneous lysis} / \text{cpm of maximum lysis}) \times 100$. Spontaneous release was assessed in the wells of the target alone and maximum lysis was obtained from wells to which Triton X-100 (final 0.1% V/V) was added instead of effector cells.

Results

Unexpectedly, anti-CD16 mAb treatment induced the downregulation of MHC class I molecule expression, that is, a slight downregulation was induced in clone 3G8-treated PMNs, whereas complete downregulation was induced in clone OKNK-treated PMNs (Figure 1). This downregulation was already observed three hours after mAb treatment. The downregulation of expression of MHC class I molecules continued for 48 hours in clone OKNK-treated PMNs, but completely recovered at 72 hours (data not shown). In contrast, in clone 3G8 treatment, slight downregulation of MHC class I

molecules was observed at three hours, which gradually recovered at eight hours and completely recovered at 24 hours (Figure 1). On the other hand, downregulation was not observed in the presence of anti-CD32 mAb (Figure 2). However, the same staining profile was observed in the case of FITC-conjugated anti-CD16 mAbs in freshly prepared PMNs (Figure 3). A slightly competitive effect was observed in clone 3G8-pretreated PMNs for clone OKNK binding, but not in clone OKNK-pretreated PMNs for clone 3G8 binding (Figure 4). Moreover, the downregulation of cell surface antigens other than MHC class I, such as CD11a, CD14 and CD45 was not observed. For instance, almost the same expression levels of CD45 were observed in PMNs pretreated with clone OKNK and clone 3G8 for 24 hours (Figure 5). It is generally considered that cells that do not express MHC class I molecules are susceptible to cytotoxicity of NK cells in PBMCs. Thus, we performed ^{51}Cr -release assay using PMNs that were cultured in the presence or absence of clone OKNK for 24 hours. Notable killing was observed in clone OKNK treated PMNs, but not PMNs cultured in the medium alone (Figure 6). Unexpectedly, although slightly less than allogeneic PBMCs, autologous PBMCs also displayed eminent killing activity (Figure 6).

Discussion

The mechanism of the downregulation of MHC class I molecule expression has not yet been clarified. It is a completely reverse phenomenon to the upregulation of MHC class I molecules in tumor cells and virus-infected cells to escape from natural-killer (NK)-cell surveillance (7, 8). It is speculated that PMNs that are activated by triggering CD16 molecules respond within three hours and attain the state of being capable of playing roles in host immune defense. After performing several roles in host immune defense, PMNs may downregulate MHC class I molecule expression, resulting in susceptibility to the cytotoxic effect of NK cells, to restrain inflammatory reactions, in which PMNs play central roles through ROS production, phagocytosis, and clearance of immune

Figure 1.

FCM profiles of MHC class I molecule expression in freshly prepared PMNs (a), PMNs cultured in medium alone (b, e, h), PMNs cultured in the presence of clone OKNK (c, f, i) or clone 3G8 (d, g, j). Cells were harvested at the indicated time and stained with PE-conjugated anti-MHC class I mAb (—) or isotype-matched PE-conjugated mouse IgG (-----) as negative control. Representative results from five independent experiments are shown.

Figure 2.

FCM profiles of freshly prepared PMNs stained with FITC-conjugated anti-MHC class I mAb, freshly prepared PMNs (a), 24-hour cultured PMNs (b) and 24-hour cultured in the presence of anti-CD32 mAb (c), and staining profiles of isotype-matched FITC-conjugated mouse IgG (-----) as negative control. Representative results from three independent experiments are shown.

Figure 3.

FCM profiles of freshly prepared PMNs stained with FITC-conjugated anti-CD16 mAbs, clone OKNK (left) and clone 3G8 (right) (—). Staining profiles of isotype-matched FITC-conjugated mouse IgG (-----) as negative control. Representative results from three independent experiments are shown.

complexes.

In line with these notions, other clone of mAb against low-affinity receptor for IgG (CD32), except for using in this study also might cause downregulation of MHC class I expression. This is remained to be resolved.

The different effects of the two mAbs have not been

explained thus far. One possible reason for the different effects is that clone OKNK has a strong potency for activating the intracellular signal pathway followed by downregulation of MHC class I molecule expression, whereas clone 3G8 has a weak potency. Alternatively, from the observation that the basal level of

fluorescence intensity was higher in PMNs that were treated with clone OKNK than in those are treated with clone 3G8, it was speculated that clone OKNK has the effect of increasing autofluorescence, than that of clone 3G8.

Figure 4.

FCM profiles obtained by direct staining with FITC-conjugated clone OKNK of PMNs pretreated with clone 3G8 (left, thin line) or without clone 3G8 (left, thick line), or by direct staining with FITC-conjugated clone 3G8 of PMNs pretreated with clone OKNK (right, thin line) or without clone OKNK (right, thick line). Staining profiles of isotype-matched FITC-conjugated mouse IgG as negative control (-----). Representative results from three independent experiments is shown.

Figure 5.

FCM profiles of CD45 expression in freshly prepared PMNs (a), PMNs cultured in the medium alone in the absence of primary mAbs (b), PMNs cultured for 24 hours and pretreated with clone OKNK (c) or clone 3G8 (d). were harvested, washed and stained with PE-conjugated antiCD45 mAb (—). Staining profiles of isotype-matched PE-conjugated mouse IgG as negative control (-----). Representative results from three independent experiments are shown.

A slightly competitive effect was observed in clone 3G8-pretreated PMNs for clone OKNK binding, but not

seen in OKNK-pretreated PMNs for clone 3G8 binding. The binding ability may be more flexible in clone OKNK than in clone 3G8. Alternatively, the binding site for

Figure 6.

Cytotoxicity of allogeneic PBMCs (upper) and autologous PBMCs (lower) against ^{51}Cr labeled PMNs cultured in the presence or absence of clone OKNK for 24 hours at the indicated effector : target ratios. After six-hour incubation, radioactivities of culture supernatants were counted with a γ counter.

clone OKNK may be partially blocked by the pretreatment with clone 3G8, suggesting that the overlapping binding site has a rather stronger affinity for

clone 3G8 than clone OKNK. Another possibility is that clone 3G8 can be completely bound without the overlapping site.

Antibodies against neutrophil-specific antigens, such as NA1 and NA2 bind to Fc γ R III, are considered to cause transfusion-related acute lung injury (TRALI) (9, 10, 11), when these antibodies are present in blood transfusion products. We confirmed that PMNs from NA1 homozygous donors exhibited downregulation of MHC class I molecule expression following anti-NA1 mAb MG38 treatment (Nichrei Co., Ltd., Japan), but not following anti-NA2 mAb GRM1 treatment (Southern Biotechnology Associates, Inc. AL, USA) and vice versa (data not shown).

Unexpected result that PMNs downregulated MHC class I molecules were killed by not only allogeneic PBMCs but autologous PBMCs may be reasonable. Because a possible explanation is deduced from the result from the result that PMNs completely finished their several roles in inflammatory site could rapidly be cleared by scavenger cells in PBMCs, probably natural killer (NK) cells may contributed to relieve inflammatory reactions in vivo.

References

1. Fanger NA, Lin C, Guyre PM, *et al.*: Activation of human T cells by major histocompatibility complex class II expressing neutrophil: Proliferation in the presence of super antigen, but not tetanus toxoid. *Blood* **89**:4128-4135, 1997.
2. Radska M, Iking-Konert S, Andrassy K, *et al.*: Polymorphonuclear neutrophils as accessory cells for T-cell activation: major histocompatibility complex class II restricted antigen-dependent induction of T-cell proliferation. *Immunology* **101**:521-530, 2000.
3. Huizinga TWJ, Kleijer M, Tetteroo PA, *et al.*: Biallelic neutrophil Na-antigen system is associated with a polymorphism on the phospho-inositol-linked Fc γ receptor III (CD16). *Blood* **75**:213-217, 1990.
4. Huizinga TWJ, Kuijpers RW, Kleijer M, *et al.*: Maternal genomic neutrophil Fc γ RIII deficiency leading to neonatal isoimmune neutropenia. *Blood* **76**:1927-1932, 1990.
5. Stroncek DF, Skubiz KM, Plachta LB. *et al.*: Alloimmune neonatal neutropenia due to an antibody to the neutrophil Fc- γ receptor III with maternal deficiency of CD16 antigen. *Blood* **77**:1572-1580, 1991.
6. Chin-Yee I, Keeney M, Krueger L, *et al.*: Supernatant from stored red cells activates neutrophils. *Transfusion Medicine* **8**: 49-56, 1998.
7. Kaufman DS, Schoon RA, Leibson PJ.: MHC class I expression on tumor cell targets inhibits natural-killer cell-mediated cytotoxicity without interfering with target recognition. *Immunol* **150**:1429-1436, 1993.
8. Liao NS, Bix M, Zulstra M, *et al.*: MHC class I deficiency: susceptibility to natural killer (NK) cells and impaired NK activity. *Science* **253**:199, 1991.
9. Zupanska B, Chrynowska M, Konopla L.: Transfusion-related acute lung injury due to granulocyte-agglutinating antibody in a patient with paroxysmal nocturnal hemoglobinuria. *Transfusion* **39**:944-947, 1991.
10. Popovsky MA, Davenport RD.: Transfusion-related acute lung injury: femme fatale? *Transfusion*, **41**:312-315, 2001.
11. Leach M, Vora AJ, Jones DA. *et al.*: Transfusion-related acute lung injury (TRALI) following autologous stem cell transplant for relapsed acute myeloid leukemia: A case report and review of the literature. *Transfusion Medicine*, **8**:333-337, 1998.