Short Communication

Environmental Conditions Affecting N_2 Fixation by *Chromatium* sp. Bloom in Lake Kaiike

MICHIRO MATSUYAMA

Faculty of Fisheries, Nagasaki University

Bunkyo-machi 1-14, Nagasaki 852

(Received December 24, 1993—Accepted November 14, 1994)

Key words: *Chromatium* sp., N_2 fixation, Lake Kaiike

Two kinds of large-sized cell bacteria, *Chromatium* sp. and *Macromonas* sp., bloom at an upper boundary of the H_2S layer of Lake Kaiike throughout all seasons (bacterial plate). Maximum cell density of *Chromatium* sp. observed in Lake Kaiike was on the order of 10^5 – 10^6 cells $\cdot$ ml^{-1} , the highest density exhibited by a member of the Chromatiaceae family in natural unpolluted waters⁴⁾. Two requisites for bacterial photosynthesis, light and H_2S , are met only at the upper boundary of the H_2S layer to cause the bloom of *Chromatium* sp. Little information has been presented on environmental factors favoring *Macromonas* bloom.

Chromatium sp. is able to fix N_2 in the presence of light and H_2S , and the N_2 fixation is predominantly controlled by photosynthesis²⁾. Thus, the bacterium can grow well by using N_2 as the sole nitrogen source when light is provided at the saturation level (ca. 2,000 lux). However, *in situ* measurement of N_2 fixation by the C_2H_2 reduction method at the upper boundary of the H_2S layer of Lake Kaiike revealed very low N_2 fixation activity^{2,3)}, although the level of photosynthesis was measurable. A question arises what factor is responsible for the observed low rate of N_2 fixation. This study was conducted to find an answer to this question.

Figure 1 shows vertical profiles of the cell density of two bacterial species and environmental variables in Lake Kaiike on 15 December 1992. The methods of lake water sampling, chemical analyses and bacterial counting were the same as described in Matsuyama and Shirouzu¹⁾.

Chromatium sp. and *Macromonas* sp. appeared at 4–5 m depth layers, and reached to the maximum cell density at 6.0–6.5 m depth (6.7×10^5 and 6.2×10^6 cells $\cdot$ ml^{-1} , respectively), and below that layer their cell density decreased sharply. Below 3 m depth, dissolved O_2 decreased with depth and disappeared at 6.5 m depth. H_2S was detected at 5.5 m depth, reaching 1 mM at 10 m depth. Concentrations of NH_4^+ and PO_4^{3-} were very low in the water above 5 m depth ($< 1 \mu M$). Their concentrations increased gradually with depth, reaching 400 μM (NH_4^+) and 90 μM (PO_4^{3-}) at 10 m depth.

Figure 2 shows the N_2 fixation rate measured by the C_2H_2 reduction method^{2,3)} of the bacterial plate sample collected from 6 m depth on 15 December 1992. The water samples were filled in a 1-l sterile glass bottle and brought to the laboratory under a cool, dark condition. After inverting the samples several times to disperse the bacterial cells uniformly, each 100 ml of the samples was introduced into a series of 100-ml syringes without exposure to air. A 10 ml aliquot of 10% C_2H_2 in

Fig. 1. Vertical profiles of the blooms of *Chromatium* sp. (A) and *Macromonas* sp. (B), and some environmental variables in Lake Kaiike on 15 December 1992.

argon gas was then injected into each syringe, followed by the addition of various amounts of H₂S (as a neutralized Na₂S solution) and NH₄⁺ (as NH₄Cl). Within 48 h after collection, they were incubated at 25°C under a continuous light (300 lux) with an incandescent lamp. After 2 days and 5 days incubation, 0.5 ml of the gas phase in each syringe was assayed by C₂H₄ measurement using a semiconductor detector-gaschromatograph (Sensortec Inc. Ltd.).

Without H₂S addition, the bacterial plate sample showed limited formation of C₂H₄ (Fig. 2), indicating a limitation of the N₂ fixation by deficiency of H₂S (50 μM) (Fig. 1) and probably a more severe limitation by the presence of NH₄⁺ (30 μM) as mentioned later. H₂S addition markedly accelerated N₂ fixation while the simultaneous addition of NH₄⁺ depressed it.

Figure 3 shows the change in the C₂H₄ concentration in the pure culture of *Chromatium* sp. Prior to the culture experiment, the bacterium was grown in the inorganic medium of Pfennig⁵⁾, which contained 6.4 mM NH₄⁺ as a nitrogen source. The bacteria from the exponential growth phase were centrifuged, followed by dispersion in a fresh H₂S- and NH₄⁺-free medium and repeated centrifugation. These treatments were done to get rid of both H₂S and NH₄⁺ from the cell suspension. The bacterial cells were finally resuspended in H₂S- and NH₄⁺-free medium and dispersed into several 100-ml syringes. The bacterial suspension was then incubated under a temperature and light condition similar to those employed for the N₂ fixation measurement. During the incubation, a fixed amount of H₂S (0.3 mM) and various amounts of NH₄⁺ (0–220 μM) were added daily. The resul-

Fig. 2. N₂ fixation rate of the bacterial plate sample of Lake Kaiike added with different amounts of H₂S and NH₄⁺. The samples were collected from 6 m depth on 15 December 1992. Each treatment was performed in duplicate. Vertical bars represent the range of two replicates. Numbers above the columns indicate incubation time (days).

tant H₂S concentration in the suspension was not markedly different from the *in situ* concentration throughout the experiment.

In the absence of NH₄⁺, the N₂ fixation activity of the bacterium was considerably high (Fig. 3). But an addition of NH₄⁺ depressed the N₂ fixation, indicating that NH₄⁺ was inhibitory for the bacte-

rial N₂ fixation as reported on other N₂ fixing microorganisms⁶⁾. Even a daily addition of 30 μM NH₄⁺, one-tenth of that of H₂S, was enough to depress the bacterial N₂ fixation completely.

Increased N₂ fixation of the bacterial plate sample at 5 days incubation in which NH₄⁺ was externally added up to 20 μM (Fig. 2) indicates that an addition of 0.6 mM H₂S was sufficient enough to use up environmental NH₄⁺ up to 50 μM (initially contained NH₄⁺ plus externally added NH₄⁺), and remained H₂S was turned to the bacterial use for N₂ fixation. While, limited N₂ fixation of the sample added with 0.6 mM H₂S and 100 μM NH₄⁺ was probably due to remaining NH₄⁺ after H₂S consumption. The relative abundance of NH₄⁺ and H₂S in the bacterial surroundings seems to have a decisive effect on the N₂ fixation. For the occurrence of bacterial N₂ fixation, a significant amount of H₂S must be present after NH₄⁺ consumption.

In the water layers below the bacterial plate, H₂S and NH₄⁺ occurred at a molar ratio of ΔH₂S: ΔNH₄⁺ = 1 : 0.36 (Fig. 1), which was close to that reported in an anoxic water column of Lake Nitinat⁸⁾, suggesting these compounds seemed to be formed by sulfate reduction expressed by the following equation⁷⁾:

$(\text{CH}_2\text{O})_{106}(\text{NH}_3)_{16}\text{H}_3\text{PO}_4 + 53\text{SO}_4^{2-} = 106\text{CO}_2 + 53\text{S}^{2-} + 16\text{NH}_3 + 106\text{H}_2\text{O} + \text{H}_3\text{PO}_4$, where $(\text{CH}_2\text{O})_{106}(\text{NH}_3)_{16}\text{H}_3\text{PO}_4$ is the empirical formula of average organic matter to be decomposed. H₂S production which is required for N₂ fixation of *Chromatium* sp. is accompanied by liberation of NH₄⁺ as much as one-third of H₂S on a molar basis. Therefore, as sulfate reduction is the principal process of organic matter decomposition below the bacterial plate, *Chromatium* sp. would never depend on the N₂ fixation.

Fig. 3. N_2 fixation rate of the pure culture of *Chromatium* sp. under daily addition of H_2S (0.3 mM) and different amounts of NH_4^+ (0–220 μM). Numbers in the figure indicate added NH_4^+ concentrations. Bacterial density at the start of the culture was 1.8×10^6 cells $\cdot$ ml $^{-1}$.

References

- 1) Matsuyama, M. and E. Shirouzu. 1978. Importance of photosynthetic sulfur bacteria, *Chromatium* sp., as an organic matter producer in Lake Kaiike. *Jpn. J. Limnol.* **39**: 103–111.
- 2) Matsuyama, M. 1986. N_2 fixation of a large-phototrophic bacterium isolated from the bacterial plate of Lake Kaiike with some considerations on its *in situ* growth. *Jpn. J. Limnol.* **47**: 369–375.
- 3) Matsuyama, M. 1987. H_2 production by a large phototrophic bacterium isolated from the bacterial plate of Lake Kaiike. *Jpn. J. Limnol.* **48**: 133–136.
- 4) Matsuyama, M. 1988. Ecology of a large-celled phototrophic bacterium densely populating the upper boundary of H_2S water of Lake Kaiike. *Bull. Jpn. Soc. Microbiol.* **3**: 35–46. (in Japanese)
- 5) Pfennig, N. 1965. Anreicherungskulturen für rote und grüne Schwefelbakterien. *Zbl. Bakt.* **1**. Abt. Org. Suppl. **1**: 179–189, 503–505.
- 6) Postgate, J. Nitrogen Fixation. Edward Arnold, London, 1978.
- 7) Redfield, A.C., B.H. Ketchum and F.A. Richards. 1963. 2. The influence of organisms on the composition of sea-water. in "The Sea", vol. 2. ed. by M.N. Hill, E.D. Goldberg, C.O'D. Iselin and W.H. Munk. Interscience, New York and London, pp. 26–77.
- 8) Richards, F.A., J.D. Cline, W.W. Broenkow and L.P. Atkinson. 1965. Some consequences of the decomposition of organic matter in Lake Nitinat, an anoxic fjord. *Limnol. Oceanogr.* **10**: R185–R201.