Letter to the Editor

To the Editor:

To the Editors and the authors presenting “Cervical Unilateral Open-Door Laminectomy With Titanium Miniplates Through Newly Designed Hydroxyapatite Spacers” in *Neurologia medico-chirurgica* 2011 September issue.

The authors make a reasonable proposal to keep the muscles in the counter side intact during one-sided open-door laminectomy.

I have one question on how to form a ditch at the appropriate location (just medial to the facet) in one segmental laminoplasty, in which retraction of the muscle group is too difficult to provide adequate space for the operator to make the ditch at the appropriate location.

Norio AOYAGI, MD
Department of Neurosurgery
Sumida Chuou Hospital
Tokyo, Japan

Reply:

We thank Dr. Aoyagi for his comments. As he says, a gutter is difficult to make on the contralateral side without detaching the muscles in our method. Recently, we measured the distance from the midline to the adequate portion of the hinge on the preoperative CT scan. In actual surgery, the muscles on the contralateral side of the lamina are detached over the periosteum partially, and a 4-mm diamond burr with a shaft marked with Pyoktanin blue according to the preoperative measurement is inserted under the partially detached muscle. The diamond burr is then moved similarly to a windshield wiper to form the gutter.

Satoshi TANAKA, MD
Department of Neurosurgery
Shioya Hospital
International University of Health and Welfare
Yaita, Tochigi, Japan

Article under discussion

Neurol Med Chir (Tokyo) 51, 673–677, 2011

Cervical Unilateral Open-Door Laminoplasty With Titanium Miniplates Through Newly Designed Hydroxyapatite Spacers —Technical Note—

Satoshi TANAKA,¹ Takashi TASHIRO,¹ Akira GOMI,² and Hiroshi UJIE³

¹Department of Neurosurgery, Shioya Hospital, International University of
Health and Welfare, Yaita, Tochigi;

²Department of Pediatric Neurosurgery, Tochigi Children's Medical Center,
Jichi Medical University, Shimotsuke, Tochigi;

³Department of Neurosurgery, Tokyo Rosai Hospital, Tokyo

Abstract

A newly designed hydroxyapatite (HA) spacer for cervical laminoplasty is provided with a hole through which the titanium miniplate is passed. A new method of unilateral open-door laminoplasty with titanium miniscrew and miniplate through this HA spacer is described. Twenty-two patients with cervical stenotic disorders were treated by this procedure. Only the predominantly affected side of the laminae is exposed. The supraspinous and interspinous ligaments and the deep muscle layer in the contralateral side are left intact. A 4-mm wide gutter near the intervertebral joint is made and contralateral hinges are made under the deep muscle layer with minimum detachment of these muscles. The opened lamina is fixed with the titanium miniplate passed through the hole in the newly designed HA spacer. Mean operative time for the 22 operations was 193.5 minutes. Mean enlargement of the minimum spinal canal diameter was 50.9%. Neurological outcomes after 3 months were evaluated according to the Japanese Orthopaedic Association (JOA) scoring. Mean recovery rate of JOA score by Hirabayashi's method was 49.5%. Only 2 patients reported postoperative axial neck pain. Our method for unilateral open-door laminoplasty provides adequate decompression and tight fixation of laminae, and is less invasive to the posterior supporting elements of cervical spine.