12. Various Kind of Tremor in Human Cases and the Experimentally Produced Tremor in Monkeys

H. NARABAYASHI M.D. and K. KOSAKA, M.D.
M. SHIMAMURA M.D.

*Department of Neurology, and Department of Neurosurgery
Juntendo University
Fuchu Ryoiku Center*

13. Posttraumatic Tremor and Stereoecephalotomy

Chihiro OHYE, Akira FUKAMACHI and Hirotaro NARABAYASHI

*Department of Neurosurgery, University of Gunma
Neurological Clinic*

14. Stereotaxic Operation for Treatment of Hypotonic Hyperkinetic Syndrome

Takuhei KIDO, Norihiko KOMAI, Hideo NISHINA, Toru HAMADA,
Ryohei KOBAYASHI, Ryuso YABUSHITA, Seiki TAKATSUNA
Yoshikatsu SHIMAMOTO and Shingo KOSHIMICHI

Neurosurgical Clinic, Wakayama Rosai Hospital

Five cases of hypotonic hyperkinetic syndrome treated by thalamotomy (VL-basis) were reported. Outline of each cases were as follows.

Case 1 K. I., 62-year-old female

Symptoms; choreic movement of the left hand and foot with dysesthesia of the left side of the face.

Supposed etiology; vascular.

Target; the right ventrolateral basis of the thalamus.

Result; Complete relief of choreic movement with transient facial weakness.

Case 2. O. U., 62-year-old female

Symptoms; choreic movements of the right extremities, particularly dominant in the hand.

Supposed etiology; vascular.

Target; the left ventrolateral basis of the thalamus.

Result; complete relief of choreic movements with transient hemiparesis.

Case 3. T. M., 70-year-old male