18. Electroencephalograph and Halogenated Inhalation Anesthetics

Akira INAMOTO, Kenjiro MORI and Kikuo NIO

Department of Anesthesiology, Kyoto University

Caudate-induced spindles and recruiting response were studied under halothane and methoxyflurane anesthesia for cats.

The results were as follows,

- 1) Duration and amplitude of caudate-induced spindles were both increased as anesthesia proceeded.
- 2) Under 0.5% methoxyflurane anesthesia, threshold voltage, latency and duration of each spindle showed following changes as anesthesia proceeded,
  1. Threshold voltage showed initial increase, and slight decrease thereafter.
  2. Latency was shortened markedly in 10 minutes, then prolonged gradually and made constant after another 10 minutes.
  3. Duration was decreased in the first several minutes, then increased gradually and made constant after 30 minutes.
- 3) Individual spindle at light and relatively deep stage of anesthesia consisted of 7-9 and 5-7 CPS. waves respectively in its beginning, the latter was more hypersynchronous.
- 4) The caudate-recruiting responses in the thalamus and cortices were affected by halothane and methoxyflurane as the caudate-induced spindles.

The above results are closely similar to that obtained by the other workers about barbiturate anesthesia. We guess that the mode of halothane and methoxyflurane anesthesia is similar to that of barbiturates; blocking the "tonic activity" of the reticular system.

---

19. EEG under Fluothane Anaesthesia in the Patients of Cerebral Lesions

Kenjiro MORI and Akira INAMOTO

Dept. of Anaesthesiology, Kyoto University

Yutaka NISHIMOTO

Dept. of Internal Medicine, Kyoto University