N-3. A Case of "Cushing Ulcer" followed by the Stereotaxic Hypothalamotomy

Dennosuke JINNAI, Heitaro MOGAMI, Kazuyoshi TOMITA,
Kiyoo KAMIKAWA, Jiro MUKAWA, Shintaro MIYAGAWA
and Mitsuhiro HOSOTANI

*Department of Surgery and Neurosurgery, Osaka University
Medical School*

It has been known that so-called "Cushing ulcer" is caused by hypothalamic injury. Recently we had an opportunity to hold an autopsy of a case of the "Cushing ulcer". The patient was a nine-year-old boy of ferocious idiot. For treatment of the ferocity a posterior hypothalamotomy was performed stereotaxically. Two days after the operation the patient had a bloody flux and hematoemesis which sustained for a week. Laparotomy revealed a duodenal perforation and the gastro-jejunostomy was carried out. However, the patient died 4 days later due to continued gastrointestinal bleeding. This duodenal ulcer was characterized by sharp delineated perforation with diameter of 7×8 mm. Lesion in the hypothalamus was demonstrated by both specimens of the brain and X-ray films taken during the stereotaxic operation.

N-4. PEG or PVG ?

Keizo MATSUMOTO, Takashi OOMOTO and Akira NISHIMOTO
Dept. of Neurosurgery, Okayama University Medical School

N-5. Film Presentation "Target Points of Globus Pallidus and Putamen"

Yutaka HORI, Shozaburo UTSUMI, Seiji MIYAMOTO,
Atsuo OKAZAKI and Hideaki ARACHI
2nd Dept. of Surgery, Nara Medical College