Bacterium xylinum の鹽化物に對する抵抗性

住 江 金 之, 津 田 國 雄 (東京農業大學)

昭和 25 年 9 月 5 日 受理

The Effect of Chlorides to restrain Multiplication of *Bact. xylinum*.

By Kinshi SUMINOE, and Kunio TSUDA (Tokyo Agric. Univ.)

緒 言 *Bact. xylinum* は食酢を濁らせる主要な菌であるが、從來之を防ぐには醋酸を濃厚にするか火入殺菌するか食鹽を加えるかである。醋酸は 5% 以上になれば發育は阻害されるが、我國普通の食酢は醋酸 4% 内外であるから屢々夏季に濁る。瓶詰の場合には火入して濁らないで永く貯えられるが、樽詰、かめ詰などの場合には開栓後店頭販賣中に濁る。又食鹽は之まで濁り防ぎに用いられ或る程度濁りの來るのをおそくするが十分に濁らないようにするには多量に使用せねばならぬので鹽からくなり食酢として使えなくなる。著者の一人(住江)⁽¹⁾は曾て食酢濁り止めに鹽化石灰と食鹽と併用すると味に悪影響がなくて目的を達する事を知り其の後廣く實用に供せられているが、當時學術的の調査を缺いていたので更めて之を明かにする事とした。

實 驗 之 部

菌種 當研究室で分離した *Bacterium xylinum* var. を用いた。

(1) 食鹽に對する抵抗性：麹汁 (Ball. 7°) に種々の濃度に食鹽(化學用)を加え菌を移植し 25°~30° の恒温器にいて 14 日間おき繁殖の模様を見た。

Table 1. The Relation of Multiplication of *Bact. xylinum* to Concentrations of Sodium Chloride in Koji Extract (Balling 7°).

days \ CaCl ₂ %	0.5	0.75	1.00	1.25	1.50	1.75	2.00	2.25
1	+	+	+	+	-	-	-	-
2	+	+	+	+	+	±	-	-
3	++	++	++	++	+	±	-	-
4	++	++	++	++	+	±	-	-
5	++	++	++	++	++	±	-	-
6	++	++	++	++	++	±	-	-
7	++	++	++	++	++	±	-	-
8	++	++	++	++	++	±	-	-
9	++	++	++	++	++	++	-	-
10	++	++	++	++	++	++	-	-
11	++	++	++	++	++	++	-	-
12	++	++	++	++	++	++	-	-
13	++	++	++	++	++	++	-	-
14	++	++	++	++	++	++	-	-

- No multiplication, ± a little multiplication, + moderate multiplication,
+ islands formed, ++ islands developed, ++ scums formed.

即ち麹汁では食鹽 2% になると發育しない。

(2) 醋酸の存在に於ける食鹽又は鹽化石灰の影響：普通我邦の食酢は醋酸 4% 内外を含むので麹汁(Ball. 3°)に醋酸を 4% 加えたものに就き食鹽又は鹽化石灰を種々の濃度に加えて發育を検した。

Table 2. Relation of Multiplication of *Bact. xylinum* to Concentrations of Sodium Chloride in Koji Extract (Ball 3°) containing 4% of Acetic Acid.

days \ CaCl ₂ %	0.2	0.4	0.6	0.8	1.0
1	+	-	-	-	-
2	+	+	+	+	-
3	++	++	+	+	-
4	++	++	++	+	-
5	++	++	++	+	-

6	+	+	+	+	-
7	+	+	+	+	-
8	+	+	+	+	-
9	+	+	+	+	-
10	+	+	+	+	-

即ち醋酸が 4% ある場合には食鹽 0.8% まではやや繁殖し, 1% となると繁殖が止る.

Table 3. Relation of Multiplication of *Bact. xylinum* to Concentrations of Calcium Chloride in Koji Extract (Ball. 3°) containing 4% Acetic Acid.

CaCl ₂ %	0.2	0.4	0.6	0.8
days				
1	+	-	-	-
2	+	+	-	-
3	+	+	-	-
4	+	+	-	-
5	+	+	-	-
6	+	+	-	-
7	+	+	±	-
8	+	+	±	-
9	+	+	±	-
10	+	+	±	-

醋酸 4% の場合, 鹽化石灰 0.6% を加えると極めて僅に發育するが 0.8% になれば止る.

(3) 食鹽と鹽化石灰を併用した場合: 第 2 表に依るとこの菌の發育を止めるには醋酸 4% を含んだ場合食鹽を 1% 加えねばならぬ. 之は鹽からくて實用にならぬので鹽化石灰と併用して見た. 又市販の食酢には醋酸 4% 内外並に 3% のもあるので麵汁(Ball. 3°)に醋酸を 3%, 4%, 4.5% 等加えたものにつき檢した.

a) 醋酸 3% とし食鹽並に鹽化石灰を種々に組合せた場合

Table 4. Relation of Multiplication of *Bact. xylinum* to Concentrations of Calcium Chloride in Koji Ext. (Ball. 3°) containing 3% Acetic Acid, and 0.2 to 0.4% of Sodium Chloride.

NaCl % CaCl ₂ %	0.2				0.3					0.4			
	0.2	0.3	0.4	0.5	0.1	0.2	0.3	0.4	0.5	0.1	0.2	0.3	0.4
days													
1	-	-	-	-	-	-	-	-	-	-	-	-	-
2	+	-	-	-	+	-	-	-	-	-	-	-	-
3	+	-	-	-	+	-	-	-	-	-	-	-	-
4	+	-	-	-	+	-	-	-	-	+	-	-	-
5	+	+	-	-	+	+	-	-	-	+	-	-	-
6	+	+	-	-	+	+	+	-	-	+	+	±	-
7	+	+	-	-	+	+	+	±	-	+	+	±	-
8	+	+	+	-	+	+	+	±	-	+	+	±	-
9	+	+	+	-	+	+	+	±	-	+	+	±	-
10	+	+	+	-	+	+	+	±	-	+	+	±	-
11	+	+	+	-	+	+	+	±	-	+	+	±	-
12	+	+	+	-	+	+	+	±	-	+	+	±	-
13	+	+	+	-	+	+	+	±	-	+	+	±	-
14	+	+	+	-	+	+	+	±	-	+	+	±	-

b) 醋酸 4% と食鹽並に鹽化石灰を種々に組合せた場合

Table 5. Relation of Multiplication of *Bact. xylinum* to Concentrations of Calcium Chloride in Koji Ext. (Ball. 3°) containing 4% of Acetic Acid and 0.1 to 0.4% of Sodium Chloride.

NaCl % CaCl ₂ %	0.1					0.2					0.3					0.4				
	0.1	0.2	0.3	0.4	0.5	0.1	0.2	0.3	0.4	0.5	0.1	0.2	0.3	0.4	0.5	0.1	0.2	0.3	0.4	0.5
days																				
1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3	+	+	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4	+	+	+	-	-	+	-	-	-	-	-	-	-	-	-	+	-	-	-	-
5	+	+	+	±	-	+	-	-	-	-	+	+	-	-	-	+	-	-	-	-
6	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-	-

7	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-
8	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-
9	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-
10	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-
14	+	+	+	±	-	+	+	+	-	-	+	+	±	-	-	+	±	-	-

c) 醋酸を 4.5% とし食鹽と鹽化石灰を種々に組合せた場合

Table 6. Relation of Multiplication of *Bact. xylinum* to Concentrations of Calcium Chloride in Koji Ext. containing 4.5% of Acetic Acid and 0.2% to 0.4% Sodium Chloride.

days	NaCl% CaCl ₂ %	0.2				0.3				0.4			
		0.1	0.2	0.3	0.4	0.1	0.2	0.3	0.4	0.1	0.2	0.3	0.4
1		-	-	-	-	-	-	-	-	-	-	-	-
2		+	-	-	-	+	+	-	-	+	-	-	-
3		+	+	-	-	+	+	-	-	+	-	-	-
4		+	+	-	-	+	+	-	-	+	-	-	-
5		+	+	-	-	+	-	-	-	+	-	-	-
6		+	+	-	-	+	+	-	-	+	-	-	-
7		+	+	-	-	+	+	-	-	+	-	-	-
8		+	+	-	-	+	+	-	-	+	-	-	-
9		+	+	-	-	+	+	-	-	+	-	-	-
10		+	+	-	-	+	+	-	-	-	-	-	-

即ち食鹽 0.2% 並に 0.3% の時は鹽化石灰 0.3% 以上は濁らない。食鹽 0.4% の場合は鹽化石灰 0.2% あれば濁らない。

(1) 住江金之：特許 74964 號

Summary

(1) The effect of chlorides (chiefly NaCl and CaCl₂) to restrain multiplication of *Bact. xylinum* (the chief bacteria which causes turbidity of vinegar) is studied.

(2) It multiplies when 1.75% of sodium chloride is added to Koji ext. (Ball.7°), but does not multiply when 2% of sodium chloride is added.

(3) It multiplies when 0.8% of sodium chloride is added to Koji ext. (containing 1% of acetic acid) but does not multiply when 1% of sodium chloride is added.

(4) It multiplies when 0.6% of calcium chloride is added to Koji ext. (containing 4% of acetic acid) but does not multiply when 0.8% of calcium chloride is added.

(5) It does not multiply in Koji ext. (Ball.3°, containing 3% of acetic acid) in the cases, when 0.2% of sodium chloride and 0.5% of calcium chloride, 0.3% of sodium chloride and 0.5% of calcium chloride, or 0.4% of sodium chloride and 4% of calcium chloride are added.

(6) It does not multiply in Koji ext. (Ball.3°, containing 4% of acetic acid) in the cases, when 0.1% of sodium chloride and 0.5% of calcium chloride, 0.2% of sodium chloride and 0.5% of calcium chloride, 0.2% of sodium chloride and 0.4% of calcium chloride, or 0.3% of sodium chloride and 0.3% of calcium chloride are added.

(7) It does not multiply in Koji ext. (Ball.3°, containing 4.5% of acetic acid) in the cases, when 0.2% of sodium chloride and 0.3% of calcium chloride, 0.3% of sodium chloride and 0.3% of calcium chloride, 0.4% of sodium chloride and 0.2% of calcium chloride are added.