Lactonase Process for the Optical Resolution of Racemic Pantolactone

The process involves stereospecific hydrolysis step with a novel enzyme "lactonohydrolase" as the catalyst for the resolution of DL-pantolactone (DL-PL). The enzyme obtained from *Fusarium oxysporum* specifically hydrolyzes D-PL to D-pantoic acid (D-PA); thus DL-PL can be separated into D-PA and L-PL. When 700 g/L aqueous solution of DL-PL was incubated with the fungal mycelia as the catalyst, D-PL in the racemic mixture was almost completely hydrolyzed to D-PA (98%*e.e.*). The mycelia were immobilized into calcium alginate gels for the practical purpose. When the gels were incubated in 350 g/L of DL-PL for 21 h at 30°C with automatic pH control (pH 7.0), 90–95% of the D-PL was hydrolyzed. The resultant D-PA in the reaction mixture had a high optical purity (91–95%*e.e.*) and the L-PL remained was unmodified. After 180 reaction cycles (*i.e.*, 180 days), the gels retained about 60% of their initial activity.

Based on these results, the commercial production of D-PL has started since 1999, through which it was been shown that the present process is highly satisfactory not only economically but also environmentally (water –49%, CO₂ –30%, and BOD –62%, comparing the former chemical resolution method).

Bioreduction System for Large-scale Production of Chiral Alcohols

We developed a novel bioreduction system, in which

E. coli transformant cells coexpressing an NAD(P)H-dependent carbonyl reductase gene and that of glucose dehydrogenase (or formate dehydrogenase) as a cofactor regenerator are used as a catalyst, for asymmetric reduction of prochiral carbonyl compounds to chiral alcohols. Production of chiral 4-chloro-3-hydroxybutanoate esters (CHBE) is a typical successful example of this bioreduction system. Aldehyde reductase of *Sporobolomyces salmonicolor* and carbonyl reductase (S1) of *Candida magnoliae* were found to catalyze NADPH-dependent stereospecific reduction of 4-chloro-3-oxobutanoate esters (COBE) to (*R*)- and (*S*)-CHBE, respectively, and they were successfully applied to the coexpression system for the practical production. 300–350 g/L of (*R*)- or (*S*)-CHBE of high optical purity (92–100%*e.e.*) was stoichiometrically obtained on incubation of the transformant cells in a reaction mixture containing COBE, glucose and a catalytic amount of NADP.

This bioreduction system is applicable to the production of many other useful chiral alcohols, by replacing the carbonyl reductase gene for other appropriate reductase genes. For this purpose, we have obtained various kinds of carbonyl reductases showing different substrate- and stereo-specificities through screening. Expression of the genes encoding these enzymes together with cofactor regenerator gene in *E. coli* cells (or other host cells) has been shown to be useful for large-scale production of various kinds of chiral alcohols.

Microbial Biotechnology and Future Applications

Roy H. Doi

(Section of Molecular & Cellular Biology, University of California, Davis)

Microbial biotechnology will play a significant role in solving many of the problems that are facing the world. Ten major problems that are challenging the world have been discussed recently at a meeting in Copenhagen, Denmark and among the Copenhagen Consensus are three problems that are relevant to microbiologists (Table 1). These challenges are (a) Communicable Diseases, (b) Malnutrition and Hunger, and (c) Sanitation and Water. In these three areas of major concern, microbiologists will be able to contribute significantly in the future and play a role by application of current and future microbiological technologies. What types of technologies will be applied in the future?

Table 1. The 10 Challenges of the Copenhagen Consensus

Armed Conflict
Climate Change
◆Communicable Diseases
Education
Financial Stability
Governance and Corruption
◆Malnutrition and Hunger
Population and Migration
◆Sanitation and Water
Subsidies and Trade Barriers

From: J. Bohannon: *Science*, **304**, 1429 (2004)¹⁾

Table 2. Examples of Future Microbiological Technologies and Areas of Study

Nanotechnology:
Food safety tests and various diagnostics
Detection of bioterrorism weapons
Virochips and Bacteriochips:
Diagnostics for rapid detection of disease bearing microorganisms
Detection of bioterrorism weapons in air and water
Marine Microbiology:
Detection of unknown microorganisms and useful products
Culturing of "unculturable microorganisms"
Use of renewable biomass for industrial purposes and for energy:
Conversion of industrial microorganisms to cellulose utilizers
Intercellular complementation—cellulosomes produced by mixed cultures

Among several possibilities, Table 2 lists a few approaches.

Nanotechnology applied to microbiological systems²⁾ will allow the rapid and sensitive identification of DNA sequences that can establish the types of microorganisms that are present in foods and water and that will lead to safety in our food and water supplies.³⁾ As part of the feature of nanotechnology, rapid identification of dangerous microorganisms in terrorism activities can also be achieved.

The use of virochips and bacteriochips^{4),5)} containing known sequences of dangerous and potentially dangerous disease causing organisms will allow rapid identification of causes of outbreaks of disease, e.g., SARS or avian flu. This technology has already paid dividends in the rapid detection of the causative agent of SARS and with further development and knowledge of nucleic acid sequences, the virochips should be even more useful for rapid detection of disease causing viruses.

Marine microbiology appears to be ripe for future studies, since the recent survey of nucleic acid sequences by Venter and his colleagues has revealed the remarkable diversity of microorganisms in the marine world.⁶⁾ In approximately 1500 liters of sea water, they found 1800 species, 148 new bacterial phylotypes, 1.2 million previously unknown genes, and 782 new rhodopsin-like photoreceptors. Thus research in marine microbiology, if properly approached, may reveal a wealth of information in the future.

Finally, in our research program, we are interested in converting non-cellulose users into cellulose utilizers. We are doing this by transferring *Clostridium cellulovo-*

rans cellulosome genes⁷⁾ into a non-cellulose degrader such as *Bacillus subtilis*. We have recently constructed a *B. subtilis* strain that is capable of making minicellulosomes.⁸⁾ By use of this strategy, it appears possible to convert many microorganisms into cellulose utilizers. This could reduce the cost of production of such items as ethanol, amino acids, and organic acids since relatively inexpensive cellulosic biomass can be used as the starting carbon substrate. Furthermore, we have developed an "Intercellular Complementation" system in which one *B. subtilis* strain expresses and secretes a mini-scaffoldin and another expresses and secretes an endoglucanase EngE. By growing a mixed culture of the two *B. subtilis* strain, we have found that the mini-scaffoldin has interacted with EngE in the culture medium to form minicellulosomes. This strategy can be used to produce designer cellulosomes.

Thus many novel approaches will be used in the future that take advantage of microbial biotechnology. This should lead to the development of many useful technologies for health care, for the food and pharmaceutical industries, and for the alleviation of many of the environmental problems facing the world. The 80th anniversary of JSBBA has seen much progress during the past 80 years and the 100th anniversary should see the development and fruition of many more successful novel technologies.

Acknowledgments I thank Dr. Ray Rodriguez for stimulating discussions and several references pertinent to this lecture. Our research was supported in part by a grant from the U.S. Department of Energy and from the RITE Institute.

References

- 1) J. Bohannon: *Science*, **304**, 1429 (2004).
- 2) V. Vogel and W. E. Thomas: *ASM News*, **70**, 113–120 (2004).
- 3) A. Stikeman: *Technology Review*, May issue, 61–66 (2002).
- 4) D. Wang, L. Coscoy, P. C. Zylberberg, Avila, H. A. Boushey, D. Ganem, and J. L. DeRisi: *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 15687–15692 (2002).
- 5) D. Wang, A. Urisman, Y.-T. Liu, M. Springer, T. G. Ksiazek, D. D. Erdman, E. R. Mardis, M. Hickenbotham, V. Magrini, J. Eldred, J. P. Latreille, R. K. Wilson, D. Ganem, and J. L. DeRisi: *pLoS Biol.*, **1**, 257–260 (2003).
- 6) J. C. Venter, K. Remington, J. F. Heidelberg, A. L. Halpern, D. Rusch, et al.: *Science*, **304**, 66–74 (2004).
- 7) R. H. Doi and A. Kosugi: *Nature Revs.-Microbiol.*, **2**, 541–551 (2004).
- 8) H.-Y. Cho, H. Yukawa, M. Inui, R. H. Doi, and S.-L. Wong: *Appl. Environ. Microbiol.*, **70**, in press (2004).