Quantitative Evaluation of Total Cellular Number and Cellular Density in the Thymolymphatic Organs of Young Adult Albino Rats by Means of DNA Determination*

By

Yoshitoki Monden

Department of Anatomy (Director: Prof. B. Osogoe) and Laboratory
of Cytochemistry (Director: Prof. A. Shibata),
Yamaguchi Medical School, Ube

Attempts to estimate the total cellular number and the rate of cell production in the hemopoietic organs have been made in a series of studies on the rat by Kindred (1938, 1940, 1942 and 1955). The methods employed in these studies are based on cell counts and mitotic counts of given areas in sections of the hemopoietic organs of the rat. Such procedures, however, meet with difficulties particularly in sections of lymph node, because variations in cellular density and irregular distribution of mitotic figures in different regions of lymphatic organs complicate the quantitative approach in sections and may, for this reason, cause uncertainty in the results.

In the earlier communications from this laboratory, a simple method for estimating total cellular number in lymphoid organs by means of DNA determination of whole tissue and of each nucleus has been reported, together with some preliminary results (Monden, 1955; Osogoe et al., 1957; Osogoe, 1958; and Osogoe and A way a, 1958). The present paper makes a more detailed description of the results of measurements on normal adult rats that have been carried out thus far by this method.

Material and Methods

Male albino rats of a subline of the Wistar strain, weighing 200 g ($\pm 2\%$), served as the material. They were maintained on a stand-

* Supported in part by a grant to Professor Bunsuke Osogoe from the Ministry of Education (Grant-in-Aid for Fundamental Scientific Research; Cooperative).

Fig. 1. Thymolympathatic organs of the rat.

1. Superficial cervical.	14. Mesocolic.
2. Deep cervical.	15. Mesenteric.
3. Superficial axillary.	16. Cecal.
4. Deep axillary.	17. Posterior abdominal wall.
5. Superficial mediastinal.	18. Inguinal.
6. Deep mediastinal.	19. Popliteal.
7. Portal.	20. Peyer's patch in the cecum.
8. Postgastric.	21. Ditto in large intestine.
9. Right renal.	22. Ditto in small intestine.
10. Left renal.	23. Ditto in rectum.
11. Para-aortic.	24. Spleen.
12. Sacral.	25. Thymus.
13. Iliac.	

The terminology is the same as used in the paper of Sanders and Florey (1940).

ard laboratory diet¹ and water *ad libitum* supplemented once a week with cabbage or other vegetables. The chief reason why the albino rat was chosen for this work is that its diet and environment can readily be standardized.

For the present type of measurement, it is essential to use animals of constant weight, for with advancing age the lymphoid organs, the thymus in particular, gradually undergo involution. In addition, sex differences in growth rate also must be considered. Taking these into account, young adult male rats weighing around 200 g were employed as the standard animals to be examined in the present study.

(1) *Determination of the total DNA content in lymphoid organs*

All lymph nodes, thymus, spleen and Peyer's patches (Fig. 1) were rapidly taken out from a sacrificed animal and chilled with ice. After weighing the tissues, they were finely minced with a pair of scissors, and homogenized with 9 volume of ice-cold 2% citric acid in a Potter Elvehjem-type glass homogenizer and made up to a volume. An appropriate volumes of this homogenate was dispensed for the Schneider method to determine, in duplicate, the total DNA content of the homogenate using diphenylamine reaction for the test of DNA. A sample of calf thymus DNA (sodium salt; N:P=1.6; P content:9.1%) served as the standard.

(2) *Determination of the average DNA content per nucleus in lymphoid organs*

This was made by a simplified procedure of the method employed by Fukuda and Sibatani (1953a) for determination of the average DNA content of rat liver nuclei. The bulk of the homogenate obtained in (1) was further diluted 5-10 fold and filtered through 8 layers of gauze to remove the fiber and clumps of unbroken cells. The dilution facilitated the sedimentation of nuclei on centrifugation. Then it was centrifuged at very low speed (500 rpm) for 3-4 minutes to sediment the last trace of unbroken cells along with a small amount of nuclei which formed a thin white layer on the top of the sediment. The supernatant (S_0) now contained only isolated nuclei and was quite homogeneous with respect to the distribution of nuclei.

¹ The diet consisted chiefly of unpolished rice, pressed barley and dried small sardines, with a small amount of cod liver oil and minerals ($\text{CaCO}_3 + \text{NaCl}$).

Counting of nuclei was made with S_0 . For this purpose duplicate 0.5–1.0 ml samples of S_0 were diluted 5–10 fold with ice-cold 2% citric acid (S_1). Nuclei were counted on S_1 in Bürker hemocytometer. Five 0.1 mm^3 sections were counted twice and averaged. The final value of the number of nuclei was the mean of the two dilutions S_1 . Two 2 ml aliquots were taken from the original S_0 in separate tubes and centrifuged at 3000 rpm for 10 minutes to sediment all the nuclei. The sediment thus obtained was then analyzed for DNA as in (1). The number of nuclei in 2 ml of S_0 was calculated from S_1 , and the average DNA-P content per nucleus was obtained by dividing the total DNA-P content of S_0 by the corresponding number of nuclei.

(3) *Determination of the percentages of lymphocytes in lymphoid organs*

For this purpose, imprint preparations were made from various lymphoid organs and stained with Giemsa. Because of rather irregular distribution of lymphocytes and other cellular elements in lymphoid organs, differential cell counts were made from different portions of the imprint preparations and averaged. In each slide at least 1000 nucleated cells were counted.

Results and Discussion

The results of measurements of the mesenteric and other lymph

Table 1. Total nucleated cell counts in the mesenteric lymph nodes.

Rat No.	Weight of Organ (mg)	DNA-P Content per Organ (mg)	DNA-P Content per Nucleus ($\times 10^{-9} \text{ mg}$)	Total Nucleated Cells per Organ ($\times 10^6$)	Total Nucleated Cells per mm^3 of Tissue ($\times 10^6$)
1	380	0.37	0.70	528	1.39
2	280	0.34	0.62	548	1.96
3	630	0.80	0.58	1,383	2.20
4	490	0.54	0.59	915	1.87
5	290	0.30	0.92	326	1.12
6	380	0.65	0.72	902	2.37
7	400	0.55	0.52	1,057	2.64
8	400	0.54	0.83	650	1.63
9	350	0.58	0.79	734	2.10
10	300	0.47	0.80	587	1.96
Mean $\pm$ S.E.*	390 $\pm$ 33	0.51 $\pm$ 0.05	0.71 $\pm$ 0.04	763 $\pm$ 121	1.92 $\pm$ 0.15
σ	103	0.16	0.13	382	0.47
v	26.4	31.4	18.3	50.1	24.2

*S.E. = Standard error.

nodes, thymus, spleen and Peyer's patches on 10 animals each are listed in Tables 1-5. From these data, the number of total nucleated cells in the individual and whole lymphoid organs can readily be

Table 2. Total nucleated cell counts in the lymph nodes other than mesenteric lymph nodes.

Rat No.	Weight of Organ (mg)	DNA-P Content per Organ (mg)	DNA-P Content per Nucleus ($\times 10^{-9}$ mg)	Total Nucleated Cells per Organ ($\times 10^6$)	Total Nucleated Cells per mm ³ of Tissue ($\times 10^5$)
1	870	0.75	0.70	1,071	1.23
2	1,210	0.94	0.62	1,516	1.25
3	1,140	0.64	0.58	1,103	0.97
4	1,030	0.83	0.59	1,406	1.36
5	1,040	0.94	0.92	1,021	0.98
6	860	0.80	0.72	1,111	1.29
7	1,180	0.65	0.52	1,250	1.05
8	1,010	0.67	0.83	807	0.80
9	870	0.87	0.79	1,101	1.27
10	1,010	1.01	0.80	1,262	1.25
Mean $\pm$ S.E.	1,020 $\pm$ 45	0.81 $\pm$ 0.04	0.71 $\pm$ 0.04	1,165 $\pm$ 79	1.15 $\pm$ 0.06
σ	142	0.13	0.13	250	0.48
v	13.9	16.0	13.8	12.4	41.6

Table 3. Total nucleated cell counts in the thymus.

Rat No.	Weight of Organ (mg)	DNA-P Content per Organ (mg)	DNA-P Content per Nucleus ($\times 10^{-9}$ mg)	Total Nucleated Cells per Organ ($\times 10^6$)	Total Nucleated Cells per mm ³ of Tissue ($\times 10^5$)
1	410	0.63	0.83	760	1.84
2	380	0.51	0.89	571	1.50
3	360	0.75	0.54	1,390	3.85
4	300	0.91	0.85	1,070	3.57
5	370	0.82	0.75	1,093	2.95
6	470	0.55	0.62	887	1.89
7	430	1.08	0.88	1,227	2.85
8	450	1.17	0.72	1,625	3.61
9	450	0.87	0.68	1,280	2.85
10	550	0.69	0.66	1,045	1.90
Mean $\pm$ S.E.	417 $\pm$ 18	0.80 $\pm$ 0.04	0.74 $\pm$ 0.04	1,094 $\pm$ 144	2.68 $\pm$ 0.27
σ	57	0.13	0.13	455	0.85
v	13.8	16.2	17.6	41.6	31.2

calculated. The figures obtained were as follows: $(763 \pm 121) \times 10^6$ in the mesenteric lymph nodes; $(1,165 \pm 79) \times 10^6$ in the other lymph nodes; $(1,094 \pm 144) \times 10^6$ in the thymus; $(1,375 \pm 189) \times 10^6$ in the spleen; $(420 \pm 33) \times 10^6$ in the Peyer's patches; totaling $(4,823 \pm 234) \times 10^6$ in

Table 4. Total nucleated cell counts in the spleen.

Rat No.	Weight of Organ (mg)	DNA-P Content per Organ (mg)	DNA-P Content per Nucleus ($\times 10^{-9}$ mg)	Total Nucleated Cells per Organ ($\times 10^6$)	Total Nucleated Cells per mm ³ of Tissue ($\times 10^5$)
1	1,050	0.93	0.83	1,120	1.07
2	1,580	1.58	0.53	2,981	1.89
3	750	0.84	0.80	1,050	1.40
4	850	0.90	0.84	1,072	1.26
5	600	0.92	0.82	1,121	1.86
6	890	1.41	0.85	1,658	1.86
7	1,200	0.94	0.80	1,175	0.98
8	600	0.92	0.76	1,210	2.02
9	1,030	0.80	0.79	1,012	0.97
10	850	0.91	0.67	1,350	1.60
Mean $\pm$ S. E.	940 $\pm$ 93	1.02 $\pm$ 0.08	0.77 $\pm$ 0.04	1,375 $\pm$ 189	1.49 $\pm$ 0.13
σ	294	0.25	0.13	597	0.41
v	31.2	24.5	16.8	43.4	27.5

Table 5. Total nucleated cell counts in the Peyer's patches.

Rat No.	Weight of Organ (mg)	DNA-P Content per Organ (mg)	DNA-P Content per Nucleus ($\times 10^{-9}$ mg)	Total Nucleated Cells per Organ ($\times 10^3$)	Total Nucleated Cells per mm ³ of Tissue ($\times 10^3$)
1	220	0.28	0.58	483	2.19
2	220	0.28	0.53	528	2.40
3	200	0.19	0.75	253	1.27
4	250	0.26	0.78	333	1.33
5	280	0.31	0.59	525	1.88
6	400	0.42	0.84	500	1.25
7	340	0.29	0.59	492	1.45
8	370	0.21	0.74	284	0.77
9	300	0.28	0.66	424	1.41
10	220	0.28	0.72	388	1.76
Mean $\pm$ S. E.	280 $\pm$ 25	0.28 $\pm$ 0.02	0.68 $\pm$ 0.03	420 $\pm$ 33	1.57 $\pm$ 0.16
σ	73	0.06	0.10	104	0.51
v	26.9	21.4	14.7	24.8	32.5

the whole lymphoid organs (Tables 1-6).

The proportions of lymphocytes to total nucleated cells in imprint preparations were found to be $89.7 \pm 2.0\%$ in the mesenteric lymph nodes, $88.5 \pm 2.3\%$ in the other lymph nodes, $97.9 \pm 0.4\%$ in the thymus, $80.7 \pm 1.5\%$ in the spleen, and $91.0 \pm 1.0\%$ in the Peyer's patches (Table 7).

Table 6. Total weight and total number of nucleated cells of the whole lymphoid organs of normal male rats weighing around 200 g.

Rat No.	Body Weight (g)	Total Weight of Lymphoid Organs (g)	Total Nucleated Cells ($\times 10^3$)
1	200	2.93	3,962
2	203	3.67	6,154
3	202	3.08	5,187
4	202	2.92	4,792
5	202	2.58	4,125
6	201	3.00	5,055
7	205	3.55	5,201
8	199	2.83	4,566
9	203	3.00	4,557
10	200	2.93	4,630
Mean $\pm$ S. E.	201 ± 0.6	3.049 ± 0.103	$4,823 \pm 234$
σ	1.9	0.326	739
ν	0.9	10.6	15.3

Table 7. Percentages of lymphocytes in smears from different lymphoid organs of normal male rats weighing around 200 g. Mean of 10 animals. Mean $\pm$ S. E.

Cell Type	Lymph Nodes		Thymus	Spleen	Peyer's patches
	Mesenteric Nodes	Other Nodes			
Lymphocytes	89.7 ± 2.0	88.5 ± 2.3	97.9 ± 0.4	80.7 ± 1.5	91.0 ± 1.0
Non-lymphocytic Cells					
Plasma Cells	9.5 ± 1.9	10.7 ± 2.0	—	0.7 ± 0.1	—
Reticuloendothels	0.6 ± 0.2	0.5 ± 0.1	1.8 ± 0.4	1.8 ± 0.3	8.0 ± 0.5
Neutrophils	0.2 ± 0.05	0.2 ± 0.02	0.2 ± 0.09	4.6 ± 0.8	0.9 ± 0.1
Eosinophils	0.1 ± 0.02	0.1 ± 0.04	0.1 ± 0.05	1.2 ± 0.1	0.1 ± 0.03
Monocytes	—	—	—	0.3 ± 0.1	—
Erythroblasts	—	—	—	10.6 ± 2.0	—
Megakaryocytes	—	—	—	0.1 ± 0.04	—

Using the data noted above, the number of total lymphocytes in the individual and whole lymphoid organs was then computed. The computations showed that total lymphocytes numbered $(684 \pm 123) \times 10^6$ in the mesenteric lymph nodes, $(1,031 \pm 106) \times 10^6$ in the other lymph nodes, $(1,071 \pm 145) \times 10^6$ in the thymus, $(1,100 \pm 169) \times 10^6$ in the spleen, $(382 \pm 34) \times 10^6$ in the Peyer's patches, amounting to $(4,268 \pm 577) \times 10^6$ in the whole lymphoid organs² (Table 8).

Table 8. Total number of lymphocytes present in the chief thymolymphatic organs of normal male rats weighing around 200 g. Mean $\pm$ S. E.

Organ	Total Number of Nucleated Cells $\times (10^3)$	Percentage of Lymphocytes	Total Number of Lymphocytes $(\times 10^3)$
Mesenteric Lymph Nodes	763 $\pm$ 121	89.7 $\pm$ 2.0	684 $\pm$ 123
Other Lymph Nodes	1,165 $\pm$ 79	88.5 $\pm$ 2.3	1,031 $\pm$ 106
Thymus	1,094 $\pm$ 114	97.9 $\pm$ 0.4	1,071 $\pm$ 145
Spleen	1,375 $\pm$ 189	80.7 $\pm$ 1.5	1,100 $\pm$ 169
Peyer's patches	420 $\pm$ 33	91.0 $\pm$ 1.0	382 $\pm$ 34
Total Thymolymphatic Organs	4,823 $\pm$ 234*		4,268 $\pm$ 577

* Calculated from the figures for total nucleated cells present in the chief thymolymphatic organs of individual animals and not from the data given above in the same column (cf. Table 6).

Since the average number of total lymphocytes in the circulating blood has been found to be $(132 \pm 12.0) \times 10^6$ from a study of 50 normal male rats weighing around 200 g (Monden, 1958), the ratio of this value to the number of total lymphocytes in the thymolymphatic system is calculated to be about one to thirty-two³. This indicates that the lymphocyte reserve in the thymolymphatic system is 32 times as great as the number of lymphocytes present in the blood.

The cellular density in lymphoid organs, that is, the number of nucleated cells per unit volume of the tissue, was then computed from the weight of the organ and the number of total nucleated cells per organ, assuming the specific gravity of the lymphoid tissue to be

² In the earlier communications (Monden, 1955; Osogoe et al., 1955; and Osogoe, 1958), the corresponding figure was computed to be $(4,120 \pm 651) \times 10^3$. A slight increase in the figure is due to increase in the number of observations and to more accurate calculations.

³ In the earlier communications (Monden, 1955; Osogoe et al., 1957; and Osogoe, 1958), the corresponding ratio was estimated to be 1:36.

approximately 1.00. The values obtained for total nucleated cells per 1 mm³ of tissue were as follows: $(1.92 \pm 0.15) \times 10^6$ in the mesenteric lymph nodes; $(1.15 \pm 0.06) \times 10^6$ in the other lymph nodes; $(2.68 \pm 0.27) \times 10^6$ in the thymus; $(1.49 \pm 0.13) \times 10^6$ in the spleen; and $(1.57 \pm 0.16) \times 10^6$ in the Peyer's patches (Table 1-5). As previously expected, the cellular density is highest in the thymus and it decreases in the following order: mesenteric nodes, Peyer's patches, spleen, lymph nodes other than mesenteric nodes.

It should be emphasized here that all of the numerical analyses made in the present study showed relatively small variation in the values obtained. As seen in Tables 1-6, the coefficient of variation in each measurement did not exceed 50% and lies in most instances between 16 and 43%. It is of interest to note that the values found for total lot of lymphoid organs showed less variations than those obtained for the individual lymphoid organs (cf. Tables 1-5 and Table 6).

It is generally believed that the amount of lymphoid tissue is subject to considerable individual variation even in the animals which are of the same weight or age class and are normal according to ordinary laboratory standards. In his extensive quantitative studies of the lymphoid organ by chemical determination of the nuclear substance (nucleic acid phosphorus), Andreassen (1943) demonstrated that, in adult rats, the coefficient of variation in the amount of lymphoid tissue was 44% for the total lymph nodes, whereas the corresponding values for the individual lymphoid organs greatly varied between 43 and 341%. The value for the thymus was 66%. (The corresponding figure is 31.2% in our material.)

That the variability of our material with regard to the amount of lymphoid tissue is generally much less than that of Andreassen is of particular interest. It must be borne in mind, however, that our material is too small in number to draw definite conclusions as regards the variability of the lymphoid organs. In order to gain further information on this point, an extensive study is now being made by Sasaki in our laboratory. The results obtained so far seem to confirm the present observations (Sasaki, unpublished observations).

The present method of evaluating total cellular number in the tissue is based principally on the constancy of the DNA content per nucleus in the diploid nuclei. The degree of accuracy or reliability of this method, therefore, depends upon whether the estimated values for the DNA

content per nucleus are reasonable or not. Thomson et al. (1953), who made a number of accurate measurements on various tissues of rats weighing from 195 to 250 g, claim that the average DNA-P content per nucleus from the tissues, such as spleen, kidney, pancreas, small intestine, leukocytes, bone marrow, thymus, heart, lung and salivary glands, lies between 0.65×10^{-9} mg and 0.70×10^{-9} mg. As regards the DNA-P content per nucleus in the rat liver, however, divergent values are reported in the literature as follows: 0.60×10^{-9} mg by Cunningham et al. (1951); 0.80×10^{-9} mg by Mirsky and Ris (1952); 0.86×10^{-9} mg by Vendrely, R. and C. Vendrely (1949, 1952); 0.90×10^{-9} mg by Thomson et al. (1953); and 1.01×10^{-9} mg by Harrison (1951). Of particular interest in this connection is the work of Fukuda and Sibatani (1953b), who have demonstrated that during the earlier course of the postnatal growth of the rat, the average DNA content of liver nuclei shows a certain age variation, and that this value is determined by the maximum body weight which the animals have attained irrespective of growth rate or nutritional conditions. This is one of the main reasons why the animals of constant weight must be used in the present type of measurements.

From the above considerations, it is expected that the values determined for average DNA-P content per nucleus in the different lymphoid tissue of rats of approximately constant weight would agree fairly well with each other. As a matter of fact, no significant differences are to be seen among the following figures obtained: $(0.71 \pm 0.04) \times 10^{-9}$ mg in the whole lymph nodes; $(0.74 \pm 0.04) \times 10^{-9}$ mg in the thymus; $(0.77 \pm 0.04) \times 10^{-9}$ in the spleen; and $(0.68 \pm 0.03) \times 10^{-9}$ mg in the Peyer's patches (Tables 1-5). Furthermore, these values are in good agreement with the figures of Thomson et al. (1953), who claim that the DNA-P content per nucleus lies between 0.65×10^{-9} mg and 0.70×10^{-9} mg in the rat tissues as already cited. Therefore, the values obtained in the present study for the DNA-P content per nucleus in the lymphoid organs may all be considered to be quite reasonable.

In the present study, the calculated figures for total lymphocytes in the individual as well as in the whole lymphoid organs are slightly greater than the corresponding figures reported in the preliminary communications (Monden, 1955; Osogoe et al., 1957; Osogoe, 1958), owing to increase in the number of observations and to more accurate calculations. Hence, it seems desirable to make renewed calculations on the daily production of lymphocytes by mitosis in the

thymolymphatic organs based on the present data. For the daily mitotic rates in the individual lymphoid organs the figures of Ito reported in the preliminary papers (Osogoe et al., 1957; Osogoe, 1958; and Osogoe and A way a, 1958) were used, and the computations were made on a similar assumption as in the earlier studies that the mitotic rate is the same for both the lymphocytic and the non-lymphocytic elements (Table 9). The calculations showed that

Table 9. Average number of lymphocyte mitoses per day in lymphoid organs of normal male rats weighing around 200 g. Mean $\pm$ S. E.

Organ	Total Number of Lymphocytes ($\times 10^3$)	Total Percentage of Mitoses per Day*	Number of Mitoses per Day ($\times 10^3$)
Thymus	1,071 $\pm$ 145	22.23 $\pm$ 1.47	238 $\pm$ 47.6
Mesenteric Lymph Nodes	684 $\pm$ 123	7.85 $\pm$ 0.70	54 $\pm$ 14.5
Other Lymph Nodes	1,031 $\pm$ 106	6.65 $\pm$ 0.51	69 $\pm$ 12.4
Spleen	1,100 $\pm$ 169	6.65 $\pm$ 0.51**	73 $\pm$ 11.8
Peyer's patches	382 $\pm$ 34	6.65 $\pm$ 0.51**	26 $\pm$ 2.5
Total Lymphoid Organs	4,268 $\pm$ 577		460 $\pm$ 88.8 (361 $\pm$ 74.5***)

* The data from the observations of Ito reported in the preliminary communications (Osogoe et al., 1957; Osogoe, 1958; and Osogoe and A way a, 1958).

** The daily mitotic rate in the spleen as well as in the Peyer's patches is assumed to be equal to that observed in the lymph nodes other than the mesenteric nodes.

*** The figures when the spleen and Peyer's patches are excluded.

the number of mitoses per day amounted to $(238 \pm 47.6) \times 10^6$ in the thymus, $(54 \pm 14.5) \times 10^6$ in the mesenteric lymph nodes, $(69 \pm 12.4) \times 10^6$ in the other lymph nodes, $(73 \pm 11.8) \times 10^6$ in the spleen and $(26 \pm 2.5) \times 10^6$ in the Peyer's patches, totaling $(460 \pm 88.8) \times 10^6$ in the whole lymphoid organs. Here, the daily mitotic rate of lymphocytes in the spleen and Peyer's patches, in which no mitotic counts had been made, was assumed to be equal to that observed in the lymph nodes other than the mesenteric nodes.

The number of lymphocytes needed for daily growth of the lymphoid tissue was roughly estimated to total 26.6×10^6 in the whole lymphoid organs, assuming the rate of growth of the tissue to be 0.624% according to Kindred (1942). Subtracting this figure from the number of total lymphocyte mitoses per day in the whole lymphoid organs, the average number of excess mitoses per day in the

whole thymolymphatic organs was found to be 433×10^6 . This is 3.3 times as great as the number of lymphocytes present in the circulating blood⁴.

If we assume that all excess lymphocytes newly produced by mitosis in the lymphoid organs enter into the general circulation, the turnover or renewal time of the lymphocytes in the circulating blood may be estimated to be 0.30 day or 7.2 hours and the "Daily Replacement Factor" (D.R.F.) of Yoffey (1954) for the blood lymphocytes to be 3.3. In the earlier studies (Osogoe et al., 1957 and Osogoe, 1958), in which the spleen and Peyer's patches were not included, the D.R.F. was estimated to be 2.87 or 2.6.

Summary

1. In a series of 10 male albino rats of the Wistar strain weighing approximately 200 g each, total cellular number and cellular density in the thymolymphatic organs were estimated chemically by determining DNA-P content of the whole tissue and of each nucleus, using the Schneider method and diphenylamine test.

2. The values obtained for total nucleated cells in the lymphoid organs were as follows: $(763 \pm 121) \times 10^6$ in the mesenteric lymph nodes; $(1,165 \pm 79) \times 10^6$ in the other lymph nodes; $(1,094 \pm 144) \times 10^6$ in the thymus; $(1,375 \pm 189) \times 10^6$ in the spleen; $(420 \pm 33) \times 10^6$ in the Peyer's patches; and $(4,823 \pm 234) \times 10^6$ in the whole thymolymphatic organs.

3. The proportion of lymphocytes to total nucleated cells in imprint preparations was found to be: $89.7 \pm 2.0\%$ in the mesenteric lymph nodes; $88.5 \pm 2.3\%$ in the other lymph nodes; $97.9 \pm 0.4\%$ in the thymus; $80.7 \pm 1.5\%$ in the spleen; and $91.0 \pm 1.0\%$ in the Peyer's patches.

4. The number of total lymphocytes was calculated to be: $(684 \pm 123) \times 10^6$ in the mesenteric lymph nodes; $(1,031 \pm 106) \times 10^6$ in the other lymph nodes; $(1,071 \pm 145) \times 10^6$ in the thymus; $(1,100 \pm 169) \times 10^6$ in the spleen; $(382 \pm 34) \times 10^6$ in the Peyer's patches; and $(4,268 \pm 577) \times 10^6$ in the whole thymolymphatic organs. It was estimated that the lymphocyte reserve in the whole thymolymphatic organs is 32 times as great as the number of lymphocytes present in the circulating blood.

5. The estimated figures for total nucleated cells per 1mm^3 of

⁴ For this calculation an estimate of $(132 \pm 12.0) \times 10^6$ was used for the total number of lymphocytes present in the circulating blood (cf. Monden, 1958).

tissue (cellular density) were as follows: $(1.92 \pm 0.15) \times 10^6$ in the mesenteric lymph nodes; $(1.15 \pm 0.06) \times 10^6$ in the other lymph nodes; $(2.68 \pm 0.27) \times 10^6$ in the thymus; $(1.49 \pm 0.13) \times 10^6$ in the spleen; and (1.57 ± 0.15) in the Peyer's patches. Thus, the cellular density is highest in the thymus and it decreases in the following order: mesenteric nodes, Peyer's patches, spleen, lymph nodes other than the mesenteric nodes.

6. On the basis of the figures obtained, a renewed calculation was made of the daily rate of production of lymphocytes by mitosis in the whole thymolymphatic organs.

References

- Andreasen, E. 1943 Studies on the Thymolymphatic System. Acta path. et microbiol. Scandinav., Suppl. XLIX. Ejnar Munksgaard, Copenhagen.
- Cunningham, L., A. C. Griffin and J. M. Luck 1950 The desoxypentose nucleic acid content of nuclei of normal, precancerous and neoplastic rat tissues. J. Gen. Physiol., 34: 59-63.
- Fukuda, M. and A. Sibatani 1953a Biochemical studies on the number and composition of liver cells in postnatal growth of the rat. J. Biochem., 40: 95-110.
- Fukuda, M. and A. Sibatani 1953b Relation between the body weight and the average DNA content of liver nuclei in postnatal growth of the rat. Exp. Cell Res., 4: 236-238.
- Harrison, M. F. 1951 Relation between polyploidy and amounts of desoxyribonucleic acid per nucleus in the liver and kidney of adult rat. Nature, 168: 248-250.
- Kindred, J. E. 1938 A quantitative study of the lymphoid organs of the albino rat. Am. J. Anat., 62: 453-473.
- Kindred, J. E. 1940 A quantitative study of the hematopoietic organs of young albino rats. Am. J. Anat., 67: 99-149.
- Kindred, J. E. 1942 A quantitative study of the hematopoietic organs of young adult albino rats. Am. J. Anat., 71: 207-243.
- Kindred, J. E. 1955 Quantitative studies on lymphoid tissues. Ann. New York Acad. Sci., 59 (Art 5): 746-756.
- Mirsky, A. E. and H. Ris 1952 The desoxyribonucleic acid content of animal cells and its evolutionary significance. J. Gen. Physiol., 34: 451-462.
- Monden, Y. 1955 Total number of lymphocytes contained in the thymo-lymphatic system of rats as estimated by means DNA determination. Acta haemat. Jap., 18: 617-624. (In Japanese with English summary.)
- Monden, Y. 1958 A quantitative study of cell populations in the circulating blood of young adult albino rats. (To be published.)
- Osogoe, B. 1958 A quantitative study of lymphocyte production, with special reference to its relation to myelopoiesis. Saishin-Igaku, 13: 1028-1041. (In Japanese.)
- Osogoe, B. and K. Awaya 1958 Turnover of cell populations in blood and blood-forming tissue. Acta haemat. Jap., 21 (Suppl.): 408-418.

- Osogoe, B., Y. Monden et H. Ito 1957 Étude quantitative de la production cellulaire par le système thymo lymphatique du rat. *Sang*, 28:792-737.
- Sanders, A. G. and H. W. Florey 1940 The effects of the removal of lymphoid tissue. *Brit. J. Exp. Path.*, 21:275-287.
- Sasaki, M. 1958 Unpublished observations.
- Thomson, P. Y., F. C. Heagy, W. C. Hutchison and J. N. Davidson 1953 The desoxyribonucleic acid content of the rat cell nucleus and its expressing the results of tissue analysis, with particulare reference to the composition of liver tissue. *Biochem. J.*, 53:460-474.
- Vendrelly, R. et C. Vendrelly 1949 La teneur dun oyau cellulaire en acide désoxyribonucléique à travers les organs, les individus et les espèces animals. *Experientia*, 5: 327-333.
- Vendrelly, R. 1952 L'acide désoxyribouncléious du noyau des cellules animals. *Bull. biol. Franc. Belg.*, 86:1-87.
- Yoffey, J. M. 1954 Bone marrow. *Brit. Med. J.*, 1954, Vol. 2:193-204.