**Stereological Studies on Several Ducts and Vessels
by Injection Method of Acrylic Resin**

**XV. Distributing arterial branches of the submandibular
lymph nodes in guinea pig**

by

T. Miyata, K. Kamakura and S. Kaneko

Department of Anatomy, Osaka Dental College, Osaka

(Director: Prof. Y. Taniguchi)

(With seven figures in one plate)

This work was announced at the 37th Kinki-district meeting of
the Japanese Association of Anatomists, 1965, Osaka

Miyata gave a lecture about the comparative study of arterial supply to the submandibular lymph nodes in some mammals in 1962. The present authors made observations about distributing arterial resources to the submandibular nodes of guinea pig.

Materials and Method

Fifteen adult guinea pigs have been used for observations. Plastic injection was applied by method of Taniguchi, Ohta and Tajiri (1952 and 1955). The material was injected through a cannula which was inserted into the left ventricle.

Observations

Submandibular lymph nodes are one or two, being located mediolaterally.

Distributing arterial branches are (1) a ramus of the submental artery, (2) a ramus of the facial artery and (3) a ramus of R. massetericus of the facial artery. The first one is observed in 10 cases of 19 examples, the middle in 7, and the last (Figs. 3 & 7) in 2.

(1) R. ln. submandibularis of the submental artery (Figs. 1, 2 & 3)

The ramus (0.24-0.36, M. 0.30 mm) leaves the artery about 1.8-8.2 mm distal to its origin, between the M. sternohyoideus and the

anterior venter of the M. digastricus, and passes downward along medial to the venter, and supplies the nodes which are located at ventral to the submandibular salivary gland.

(2) R. ln. submandibularis of the facial artery (Figs. 4, 5, 6 & 7)

The ramus (0.24-0.31, M. 0.26 mm) leaves the artery right before it passes laterally from the mandibular base, and continues backward to the nodes giving twigs to neighbouring muscles, along inferior to the anterior venter of the M. digastricus. Nodes of both sides in one case are supplied with the branch of only right side.

(3) R. ln. submandibularis of R. massetericus of the facial artery (Figs. 3 & 7)

The ramus (0.12-0.14, M. 0.13 mm) leaves the massetericus which passes backward along the inferior margin of the M. masseter, and enters the nodes.

Nodes in two cases, in which the branch is observable, are two, located mediolaterally.

Discussion

The observations are discussed with those of rabbit, which were reported by Miyata (1962).

The node branches of the submental artery are longer and in some cases terminal ones, passing backward in rabbit, but are somehow shorter in guinea pig.

General course of the node branch of the facial artery is tortuous in rabbit, but straight in guinea pig.

The node branches of the R. massetericus of the facial artery are observed only in a few case of both rabbit and guinea pig.

Summary

In guinea pig, Rr. lnn. submandibulares have their origins in the submental artery, the facial artery, and in some cases the R. massetericus of the facial.

The authors should like to express their thanks to Professor Y. Taniguchi for guidance and encouragement shown to them, and also to Assistant Professor Y. Ohta for his help in surveying specimens and preparing the manuscript.

References

- 1) Miyata, T. 1962: Arterial branches supplying Lymphodi submandibulares in some mammals. (Abstract of the 67th Annual Session of the Japanese Association of Anatomists, 1962). Acta Anat. Nippon., 37, 17.
- 2) Taniguchi, Y., Y. Ohta and S. Tajiri, 1952: New improved method for injection of acrylic resin. Okajimas Fol. anat. jap., Bd. 24, 259-267.
- 3) Taniguchi, Y., Y. Ohta, S. Tajiri, H. Okano and H. Hanai, 1955: Supplement to new improved method for injection of acrylic resin. Okajimas Fol. anat. jap., Bd. 27, 401-406.

Abbreviations

L	Submandibular lymph node
M	Mandibula
f	facial artery
fm	R. massetericus of facial artery
s	submental artery
×	R. ln. submandibularis

Explanation of figures

- Fig. 1. R. ln. submandibularis of the submental artery.
- Fig. 2. R. ln. submandibularis of the submental artery supplies one lymph node of the lateral side, prior to another medial node.
- Fig. 3. R. ln. submandibularis of the submental artery and the same of R. massetericus of the facial artery.
- Fig. 4. R. ln. submandibularis of the facial artery supplies one lymph node of the lateral side, prior to another medial node.
- Fig. 5. R. ln. submandibularis of the facial artery supplies two lymph nodes.
- Fig. 6. R. ln. submandibularis of the facial artery.
- Fig. 7. R. ln. submandibularis of the facial artery and the same of R. massetericus of the facial artery.