IRT における能力の最尤およびベイズ推定量の漸近的性質

小笠原 春彦

(小樽商科大学商学部)

キーワード: キュムulant、ステューデント化、加重スコア法

Asymptotic properties of the maximum likelihood and Bayes modal estimators of ability in IRT

Haruhiko Ogasawara

(Department of Information and Management Science, Otaru University of Commerce)

Keywords: cumulants, studentization, weighted score.

For an ability test developed using item response theory (IRT), estimation of the proficiencies of examinees separately from the calibration for items is a main purpose of the test. When the parameters of items have been well estimated in an item bank using a large sample size, the item parameters may be seen as population ones though they are actually estimates under possible model misspecification (p.m.m.). With the assumption of known item parameters, the ability of an examinee, seen as an unknown fixed parameter, is estimated by various ways. Maximum likelihood (ML) estimation is a standard method (Lord, 1953; Birnbaum, 1968, Section 20.3). Lord (1983) gave the asymptotic biases of the MLE of an ability and its monotone transformations under correct model specification (c.m.s.). Under p.m.m., Ogasawara (2012) derived the asymptotic cumulants up to the fourth order and the higher-order asymptotic variances of the MLEs of an ability and its transformations with and without studentization.

In addition to ML estimation, Bayesian estimation has been used, where the maximum a posteriori (MAP) estimator (Samejima, 1969, Chapter 2; Bock & Aitkin, 1981) and the expected a posteriori (EAP) estimator (Birnbaum, 1958, 1967; Bock & Aitkin, 1981) are typical ones. The MAP estimator is also called the Bayes modal estimator (BME). Lord (1984) gave the asymptotic bias of the BME with the standard normal prior under c.m.s. One of the advantages of the BME is that finite values of the estimators are available in the cases of no correct/incorrect responses while the corresponding MLEs are infinite.

The weighted likelihood estimator (WLE) given by Warm (1989) looks like a BME, where the weight corresponding to the first derivative of the logarithm of a prior with respect to the ability parameter is introduced such that the asymptotic bias disappears. Since it is difficult to derive the prior explicitly, the WLE may be seen as a pseudo BME. Another BME is given by using the Jeffreys (1946; 1961, Section 3.10) uninformative prior written as JME in this paper. It is known that the WLE becomes equal to the JME in the case of the 2-parameter logistic model (2PLM).

The MLE, BME, WLE and JME can be seen as special cases of the estimator using the general weight,

given as a function of an ability, where the weight corresponds, in form, to the log-prior derivative with respect to the ability parameter. In the case of the MLE, the weight is defined as null. The main purpose of this paper is to derive the asymptotic properties such as the asymptotic cumulants up to the fourth order with the higher-order asymptotic variance and the mean square error for the pseudo Bayes estimator using the general weight. We also show their applications to the four typical estimators mentioned above. The asymptotic properties will be given for the estimators with and without studentization. It is known that the so-called sandwich estimator of the asymptotic variance is robust against p.m.m. in the case of independent and identically distributed (i.i.d.) cases (e.g., Ogasawara, 2010). Unfortunately, for estimation of ability, the items in an ability test are generally unidentical and do not give the advantage of the sandwich estimator (note that generally only one observation per examinee is available for an item). So, studentization is performed using the estimated information with the assumption of c.m.s. However, the asymptotic properties of the estimators will be given under p.m.m. as well as under c.m.s.

The results given in this paper will be useful when we choose good point estimators among the typical ones used in practice by e.g., seeing the amounts of their mean square errors, which will be algebraically given and numerically illustrated. Estimators are also used for interval estimation of the population value of an ability, where the asymptotic distributions of the estimators after studentization are required. For interval estimation more accurate than the usual Wald confidence interval (CI), the asymptotic cumulants in addition to the usual asymptotic variance are needed and will be given with numerical illustration.

References: Ogasawara, H. (2012). Asymptotic expansions for the ability estimator in item response theory. *Computational Statistics*, 27, 661-683.

Ogasawara, H. (2013). Asymptotic properties of the Bayes and pseudo Bayes estimators of ability in item response theory. *Journal of Multivariate Analysis*, 114, 359-377.