16. *Fragile Sites in Newborns*

By Tetsuji KADOTANI, Yoko WATANABE, and Noriko KUROSAKI

The Kadotani Medical Research Foundation,
Saijohigashi, Saijo, Higashihiroshima 724

(Communicated by Sajiro MAKINO, M. J. A., March 14, 1988)

Since Dekaban (1965) reported on heritable fragile site on the human chromosomes and Lubs (1969) reported on the fragile X chromosome, numerous reports have been represented and revealed that the fragile X chromosome was associated with the commonest inherited form of the mental retardation in man. Recently Sutherland (1979) studied the fragile site in the minimal essential medium without folic acid (MEM-FA) and revealed that the frequency of fragile sites in human chromosomes was increased. However the study of the fragile site in the normal newborn subject has been still meager. This paper is represented on 20 cases of the normal newborns, in order to reveal the position and frequency of fragile sites on normal newborns.

Cytological findings. On the 20 cases of the newborns, the lymphocytes taking from the vein of umbilical cord were cultured for 72 hrs in the medium MEM-FA supplemented with 5% fetal serum (Sutherland, 1979). The slides were made by means of the air-drying technique. The chromosome observation was made with the frequency of the break and/or gap with 50 well-delineated metaphases for each newborn. The locations of the break points were identified by conventional Giemsa and G-banding staining after the destain, and the break points observed more than two times each in 50 cells were picked up as the fragile site. And those fragile sites were divided into two groups; the common fragile site-group and the folic acid sensitive fragile site-group, according to the list of the fragile site in 8th International Workshop on Human Gene Mapping in the report of Hori (1986).

Out of 1000 cells cultured in the medium MEM-FA, four fragile sites were shown; one site at 1p31 and three sites at 3p14. These break points were the common fragile sites and the sites at 3p14 in three newborns were hot points. The folic acid fragile site was not shown in this study.

Remarks and conclusion. The frequency of the chromosome break points was studied in the medium MEM-FA in lymphocytes of 20 normal newborns. For the fragile sites of neonates, Sutherland (1982) screened 1019 unselected newborns without finding fragile X in any child and with finding one autosomal folate sensitive fragile site. And furthermore, in 1985 it was reported by Sutherland that four folic acid sensitive fragile sites were detected in 2439 neonates; 12q13, 8q22, 12q13, and 11q13. In our study, no folate sensitive fragile site was detected in 20 newborns.

Chromosomal fragility occurs nonspecifically and sporadically at a very low level, and the common fragile sites are present in all individuals in population (1985, Sutherland *et al.*). Zhou *et al.* (1984) called the cluster of gaps or breaks in certain regions of chromosomes a "hot point". They reported that thirteen of the 22 peasants had a break-point at 3p14. In our study, eight cases of 20 newborns had one or two break points at 3p14 and three cases had one break

Table I. Frequencies of hot points in newborns

Case	Sex	No. of cells	No. of 3p14	No. of 6q26	No. of 16q23
1	F	50	1	0	0
2	M	50	0	0	0
3	F	50	0	0	0
4	M	50	0	0	0
5	M	50	0	1	0
6	F	50	0	0	0
7	F	50	2	0	0
8	M	50	0	0	0
9	F	50	0	0	0
10	F	50	0	0	0
11	F	50	0	0	0
12	F	50	0	0	0
13	M	50	2	0	0
14	M	50	2	1	0
15	M	50	1	0	0
16	F	50	1	0	0
17	F	50	1	1	0
18	M	50	0	0	0
19	M	50	1	0	0
20	M	50	0	0	0
Total		1000	11	3	0

Fig. 1. The same partial metaphases of newborn. A: Giemsa staining.
B: G-banding staining after destain.

point at 6q26 per 50 cells (Table I). The mean frequency of hot points at 3p14 in newborns was 1.1 breaks/100 cells. This frequency was less than 2.2 breaks/100 cells of normal peasants aged 12 to 62 years reported by Zhou *et al.* (1984). On the results of this study, it was shown that the fragile site was scarcely detected in 20 newborns.

Acknowledgements. We are cordially obliged to Emeritus Professor Sajiyo Makino, M. J. A., the senior director of the Kadotani Medical Research Foundation, for improvement of this manuscript. Financial aid from the Japan Academy is gratefully acknowledged here.

References

- Dekaban, A. (1965): J. Nucl. Med., **6**, 740-746.
Hori, T. (1986): Jpn. J. Cancer Chemother, **13**(3), part II, 667-676.
Lubs, H. A. (1969): Am. J. Hum. Genet., **21**, 231-244.
Sutherland, G. R. (1979): *ibid.*, **31**, 125-135.
—— (1982): *ibid.*, **34**, 452-458.
—— (1985): Ann. Hum. Genet., **49**, 153-161.
Sutherland, G. R. *et al.* (1985): Hum. Genet., **69**, 233-237.
Zhou, X. *et al.* (1984): *ibid.*, **67**, 249-251.