Flavonoids in plant rhizospheres: secretion, fate and their effects on biological communication

Akifumi Sugiyama*, Kazufumi Yazaki

Research Institute for Sustainable Humanosphere, Kyoto University, Uji, Kyoto 611-0001, Japan

*E-mail: akifumi_sugiyama@rish.kyoto-u.ac.jp Tel: +81-774-38-3617 Fax: +81-774-38-3623

Received June 30, 2014; accepted September 17, 2014 (Edited by T. Koeduka)

Abstract Flavonoids, one of the most-described group of plant “specialized metabolites”, consist of more than 10,000 structurally diverse compounds. Most flavonoids accumulate in plant vacuoles as glycosides, with some released by the roots into rhizospheres. These flavonoids are involved in biological communications with rhizobia, arbuscular mycorrhizal fungi, plant growth promoting rhizobacteria, pathogens, nematodes, and other plant species. Both aglycones and glycosides of flavonoids are found in root exudates and in soils. This review describes researches on the mechanisms of flavonoid secretion and the fate of flavonoids released into rhizospheres. This review also discusses the direction of future research that may elucidate the specific roles of flavonoids in biological communications in rhizospheres, enabling the utilization of flavonoid activities and functions in agricultural practice.

Key words: Biological communication, flavonoid, rhizosphere, root exudates.

Introduction

Plants synthesize a wide range of low molecular weight compounds, which are active in plant defenses against biotic and abiotic stresses and also act as attractants or repellents of other organisms. These compounds have recently been described as “specialized” rather than “secondary” metabolites, and their number has been estimated at more than 200,000 (Dixon and Strack 2003). In contrast to the primary metabolites, which are synthesized by almost all organisms, each specialized metabolite is usually synthesized by a few plant families, or even by a few species, suggesting that, during evolution, these plants acquired the ability to synthesize new specialized metabolites, which conferred adaptive advantages to these plants (Pichersky and Lewinsohn 2011).

Flavonoids are a well-described group of plant specialized metabolites, consisting of more than 10,000 different compounds (Ferrer et al. 2008). Flavonoids are a group of phenylpropanoid compounds synthesized from 4-coumaroyl-CoA and malonyl-CoA, with the flavonoid skeleton often undergoing various types of chemical modification, such as glycosylation, hydroxylation, acylation, methylation, malonylation, and prenylation, which result in diverse structures and functions. Flavonoids are classified into several groups

based on their structures and the functional groups on the C-ring, including the chalcones, flavanones, flavones, flavonols, anthocyanidins and catechins (Aoki et al. 2000) (Figure 1). Among them are the isoflavonoids, which occur mostly in legume plants and are biosynthesized from flavanones such as liquiritigenin and naringenin via 2-hydroxylation catalyzed by the cytochrome P450 enzyme isoflavone synthase (IFS).

Flavonoid functions in plants include the regulation of auxin transport, the modulation of reactive oxygen species (ROS), the coloring of flowers, and in protection against UV light (Falcone Ferreyra et al. 2012). Flavonoids are also important compounds for human life, functioning as phytoestrogens, anti-oxidants, anti-inflammatory and anti-carcinogenic compounds and in cancer prevention (Garcia-Lafuente et al. 2009; Miadokova 2009).

Due to their importance in biological processes in planta as well as in enhancing human quality of life, flavonoids have been intensively studied in various fields. This review focuses on flavonoids in rhizospheres, which are small regions around the roots of plants (Hartmann et al. 2008), where flavonoids play important roles in biological communications. Because there have been several recent reviews on flavonoids in rhizospheres (Cesco et al. 2012; Cheynier et al. 2013;

Abbreviations: ABC, ATP-binding cassette; AMF, arbuscular mycorrhiza fungi; CHS, chalcone synthase; IFR, isoflavone reductase; IFS, isoflavone synthase; MATE, multidrug and toxic compound extrusion; PAL, phenylalanine ammonia-lyase; PGPR, plant growth promoting rhizobacteria.

This article can be found at <http://www.jspcmb.jp/>

Published online December 4, 2014

Figure 1. Biosynthetic pathway for flavonoids. PAL, phenylalanine ammonia-lyase; C4H, cinnamate 4-hydroxylase; 4CL, 4-coumarate:CoA ligase; CHS, chalcone synthase; CHI, chalcone isomerase; IFS, isoflavone synthase; FS, flavone synthase; F3'H, flavonoid 3'-hydroxylase; F3H, flavanone 3-hydroxylase; FLS, flavonol synthase.

Hassan and Mathesius 2012; Weston and Mathesius 2013), this review does not include all relevant studies, but concentrates on the potential of flavonoids to act as plant-mediated signaling compounds for sustainable agriculture.

Secretion and fate of flavonoids in rhizospheres

Secretion of flavonoids from plant roots

Flavonoids are synthesized by coupling of the phenylpropanoid and polyketide pathways, with 4-coumaroyl-CoA and malonyl-CoA, respectively, being the direct precursors (Figure 1). In the phenylpropanoid pathway, phenylalanine and tyrosine are metabolized to form flavonoid skeletons as well as lignins and lignans. Genes involved in this pathway leading to the core structures of flavonoids have been characterized in detail in various plants (Saito et al. 2013). These genes have been used for metabolic engineering in various plant species as well as in microorganisms (for review see Suzuki et al. in press). Flavonoid diversity is largely due to the modification of the core structure by enzymes such as glycosyltransferases, prenyltransferases, methyltransferases and acyltransferases. These modifications confer biological activities on the resultant compounds, as well as being important for their transport and accumulation in plant cells. In general, flavonoids exist as glycosides in vacuoles, with various transporters responsible for the vacuolar accumulation of flavonoid glycosides; these include, for example, members of the ATP-binding cassette (ABC) and multidrug and toxic compound extrusion (MATE) transporter families (Shitan and Yazaki 2013; Zhao and Dixon 2010), while such transporters have not been identified from soybean so far. The most common isoflavone in soybeans (*Glycine max*) is daidzein, an aglycone commonly glucosylated by GmIF7GT, UDP-glucose:isoflavone 7-O-glucosyltransferase (Noguchi et al. 2007), followed by malonylation by GmIF7MaT, malonyl-CoA:isoflavone 7-O-glucoside-6''-O-malonyltransferase (Suzuki et al. 2007). These glucosides and malonylglucosides show high accumulation in vacuoles (Barz and Welle 1992).

Flavonoid aglycones and glycosides can be found in root exudates. The concentrations of flavonoids vary widely in plants grown axenically in mineral nutrient solution, depending on plant species, plant growth conditions and sampling techniques ((Cesco et al. 2010), our unpublished results). In *Arabidopsis*, root exudate contents and concentrations were shown to correlate with the metabolic activities of bacterial communities in rhizospheres (Chaparro et al. 2013, 2014).

There are two major mechanisms for flavonoid secretion from roots: 1) passive processes, such as root turnover, root injury and root decomposition from root cap and border cells, and 2) active processes; i.e., root exudation (Shaw et al. 2006)(Figure 2). Using plasma membrane vesicles prepared from soybean roots, these roots were shown to possess an ATP-dependent transporter that functions in the secretion

Figure 2. Biosynthesis, accumulation and secretion of isoflavones in soybean roots. Soybeans biosynthesize isoflavones, primarily daidzein and genistein, both of which are glucosylated and subsequently malonylated in the cytosol. The resulting glucosides and malonylglucosides accumulate in the vacuoles. The possible degradation products in the rhizosphere are also shown.

of genistein into the rhizosphere (Sugiyama et al. 2007). Biochemical transport assays using various inhibitors have suggested that an ABC transporter is the primary candidate for this process. This ABC-type transporter was shown to be specific for isoflavonoid aglycones, whereas similar biochemical transport assays using isoflavonoid glucosides such as daidzin and genistin as substrates did not show ATP-dependent transport into plasma membrane vesicles (our unpublished results). These results suggest that soybean roots have different mechanisms for the secretion of aglycones and glucosides (Figure 2). For example, although daidzein and genistein are synthesized in the cytosol, their glucosides, such as daidzin, genistin, malonyldaidzin and malonylgenistin, accumulate in vacuoles. The mechanisms by which metabolites that accumulate in the vacuoles are secreted into apoplasts remain to be determined; one possible mechanism could be vesicle-mediated secretion as illustrated in Figure 2. Several specialized metabolites of plants, such as nicotine, accumulate in vacuoles and are secreted into apoplasts. Characterization of these secretion mechanisms is of particular importance in future studies.

Apoplastic β -glucosidase has been identified in soybean roots, and an isoflavone conjugate-hydrolyzing β -glucosidase, GmICHG, indicated in Figure 2, has been purified from roots of soybean seedlings (Suzuki et al. 2006). Immunocytochemical experiments using antibodies against this protein showed that GmICHG

localized in the apoplasts is present in the epidermis, endodermis, and stele tissues of the root, as well as in root hairs. Isoflavonoid aglycones are a more active form, mediating legume-rhizobial interactions as well as in defenses against pathogens in rhizospheres. Apoplastic β -glucosidase may help plants rapidly secrete large quantities of the active forms of flavonoids, stored in vacuoles, into rhizospheres.

Flavonoids are also secreted by non-leguminous plants into rhizospheres. For example, at least 54 flavonoid compounds have been identified in *Arabidopsis* (Saito et al. 2013), with flavanone detected among the secreted phenolic compounds, such as coumarin, syringic acid and vanillic acid, in root exudates (Toussaint et al. 2012). An analysis of the effects of various mutations in ABC transporter genes of *Arabidopsis* on the composition of root exudates and soil microbial communities showed that root exudates from the *abcg30/pdr2* mutant contained more phenolic compounds and fewer sugars than the exudates of wild type plants (Badri et al. 2009). Because this mutation altered the expression of many genes, the differences in root exudate composition may not have been due to the direct effects of mutation in this ABC transporter gene but to the pleiotropic effects of the gene mutation. Although candidate genes involved in the secretion of flavonoids and phenolics have been identified, transporters responsible for flavonoid exudation into the rhizosphere have not yet been determined.

Fate of flavonoids in the rhizosphere

In natural soils, flavonoids are released from plant roots, with manure application potentially increasing flavonoid contents in agricultural soils. The flavonoid composition of soil was quantitatively investigated in grassland fields in the presence and absence of applied manure (Hoerger et al. 2011b). The amounts of formononetin, daidzein, biochanin A and genistein were higher in topsoil (0–10 cm) than in subsoil (10–20 cm), but flavonoid contents were not significantly increased by the application of manure. Moreover, manure application under conditions of good agricultural practice did not result in additional excretion of flavonoids into drainage water.

Legume plants are widely used as green manure. Formononetin, medicarpin and kaempferol have been detected in soils during the growth of white clover (*Trifolium repens*), presumably due to root exudation (Carlsen et al. 2012). High concentrations of the glycosides of kaempferol and quercetin were observed within one day of application of white clover to soil as green manure. However, after 16 days, most of the flavonoids were aglycones, with substantial amounts of kaempferol persisting in the soil while other flavonoids were degraded.

Because flavonoids can be aerobically metabolized by bacteria such as rhizobia and *Pseudomonas*, the measurement of the actual concentration of flavonoids in soil is technically challenging. Chemical properties of soil also affect the fate of flavonoids in soil, e.g., the concentration of catechin was about 4-fold higher in alluvial soil at pH 7 than in volcanic ash at pH 4 (Furubayashi et al. 2007). Minerals in soil such as Fe and Cu were shown to promote the degradation of catechin, while Ca was shown to protect flavonoids from degradation (Pollock et al. 2009), suggesting that mineral composition and contents in soil are important in determining the fate of flavonoids. Soil microbes, as a predominant factor, strongly affect the fate of flavonoids in soil. A study of the half-lives of various flavonoids in sterile and non-sterile soils showed that, in the latter, naringenin and formononetin were degraded within 4 and 24 h, respectively, with formononetin having a longer lag time in non-sterile soils (Shaw and Hooker 2008). Two major isoflavonoids from soybean, daidzein and genistein, had short half-lives in soil, of 37.2 and 6.0 min, respectively (Guo et al. 2011). These extremely short half-lives may have been due to the experimental design, in which isoflavonoids were added to soils collected from fields in which soybeans were continuously grown, with these soils likely containing active isoflavone-catabolizing bacteria.

In general, flavonoid glucosides are degraded more rapidly in non-sterile than in sterile soils. For example, the half-lives of flavonoid glycosides produced by *Alliaria petiolata* were shorter under non-sterile (3 to 12 h) than

sterile (12 to 46 h) conditions. Moreover, these flavonoid glycosides were not detected in bulk soils, confirming their instability in soil (Barto and Cipollini 2009). The half-lives of flavonoid glucosides may be short due to their rapid hydrolysis by glycosidase enzymes from both plants and microbes (Hartwig and Phillips 1991). One exception was formononetin-7-O-glycoside, which was detected in the rhizosphere soil of alfalfa (*Medicago sativa*), suggesting that this flavonoid glycoside was secreted from roots in large quantities or was unexpectedly stable in rhizosphere soil (Leon-Barrios et al. 1993).

Adding to the difference between sterile and non-sterile conditions, rhizosphere microbial communities whose composition depends on the growth condition of plants (Chaparro et al. 2014; Mougél et al. 2006; Sugiyama et al. 2014) are also important in affecting the fate of flavonoids in the rhizosphere. The rate of flavonoid degradation by rhizosphere microbial communities may increase during plant growth because these rhizosphere soils are exposed to the continuous release of flavonoids from roots (Shaw et al. 2006). When microbial communities are exposed to high concentrations of flavonoids, flavonoid-degrading microbes will likely prevail around the roots. The addition to soils of *Arabidopsis* root exudates collected from liquid medium was found to modify the microbial composition of these soils, depending on the fractions of root exudates; i.e. phenolic compounds exhibited positive correlation with more operational taxonomic units than other fractions (Badri et al. 2013). Because the microbial community is one of the most important determinants of the fate of flavonoids in soils, the half-life of a particular flavonoid in soil may largely differ depending on the soil microbes.

Extraction methods are another key factor for the measurement of flavonoid concentrations in soils. Organic solvents such as methanol, ethanol, acetone and ethyl acetate have been widely used to extract flavonoids from soil. A systematic comparison of various solvents for the extraction of formononetin from soil, with quantification relative to ¹³C-labeled internal standards, found that a mixture of methanol and acetonitrile had the highest extraction efficiency, followed by methanol (Hoerger et al. 2011a). Moreover, extraction was increased at higher temperature but not at higher pressure. Due to differences in soils and extraction methods it is not feasible to compare reported flavonoid concentrations and fate; as summarized in a review, contradictory results have been obtained (Cesco et al. 2012).

Effect on biological communication

Interaction with rhizobia

Interactions between legume plants and rhizobia are of particular importance in agriculture and ecology. The molecular mechanisms of bacterial and plant

perception, signaling, organogenesis and nitrogen fixation have been intensively studied, with many components essential for these symbiotic interactions identified (Popp and Ott 2011; Wang et al. 2012; Werner 2007; Yokota and Hayashi 2011). Due to the chemotaxis of rhizobia to root exudates, these rhizobia can find their host legume plants in soils. The bacterial proteins responsible for the recognition of plant signal molecules were identified as being NodD, a LysR-type regulator acting as a transcriptional activator for other nod genes, and NodA, NodB, and NodC, which together synthesize the backbone structure of a Nod factor, lipochitooligosaccharide, responsible for inducing symbiotic signals in the roots of host plants (Sugiyama and Yazaki 2012). Using a reporter system consisting of a nod promoter fused to LacZ, signaling molecules secreted by legume plants were identified as flavonoids, such as luteolin in alfalfa, 7,4'-dihydroxyflavone and geraldone in white clover, and daidzein and genistein in soybean, (Djordjevic et al. 1987; Kossak et al. 1987; Peters et al. 1986; Redmond et al. 1986). The induction of *nod* gene expression by flavonoids was studied using alfalfa-*S. meliloti* interaction as a model (Peck et al. 2006), showing that only host specific flavonoid, luteolin, up-regulates the expression of *nod* genes, while noninducing flavonoids such as naringenin compete with luteolin for binding to NodD1.

Upon the perception of Nod factors, legume plants rapidly respond to signals, leading to nodule formation on roots. Genes induced by these bacterial factors have been identified, but less is known about which metabolites, especially specialized metabolites, are increased, decreased, or accumulate during the nodulation process. Flavonoid biosynthesis genes have been shown to be induced during the early stage of nodulation; for example, *B. japonicum* induces the expression of phenylalanine ammonia-lyase (PAL) and chalcone synthase (CHS) in soybeans (Estabrook and Sengupta-Gopalan 1991). Nod factor also up-regulates the expression of isoflavone reductase (IFR) on suspension cells of alfalfa (Savoure et al. 1994), as well as inducing the expression of peroxidases, chitinases, WRKY transcription factors and IFS in the roots of *L. japonicus* (Nakagawa et al. 2011). The presence of flavonoids in the roots was shown to be essential in the stimulation of Nod factor synthesis in infection threads (Zhang et al. 2009). During infection, the contents of kaempferol are increased, inhibiting auxin transport and causing auxin accumulation at nodule initiation sites, thus stimulating cell division and nodule organogenesis (Mathesius et al. 1998; Wasson et al. 2006). In *Medicago truncatula*, the silencing of genes in different branches of the flavonoid pathway, such as chalcone synthase, the key entry-point enzyme for flavonoid biosynthesis, IFS and flavone synthase, showed that flavonols like kaempferol

act as auxin transport inhibitors during nodulation, while flavones such as 7,4'-dihydroxyflavone act as nod-gene inducers (Zhang et al. 2009).

Differences in flavonoid composition have been observed in legume roots infected with symbiotic rhizobia. In alfalfa, for example, the inoculation of rhizobia was shown to strongly alter the flavonoid composition in roots, with medicarpin and its glucoside as well as formononetin-7-O-(6''-O-malonylglycoside) found only in infected roots (Dakora et al. 1993). Co-inoculation of the common bean (*Phaseolus vulgaris*) with *Azospirillum* and *Rhizobium* was found to increase the exudation of nod-gene-inducing flavonoids from roots, as well as relieving symbiotic interactions under conditions of salt stress (Dardanelli et al. 2008). These results suggest that plants respond to bacteria in rhizospheres by altering the flavonoid composition of both roots and root exudates, possibly to facilitate and fine-tune symbiotic interactions with bacteria.

In addition to legume plants, more than 200 species from eight families of non-legume plants, collectively called actinorhizal plants, can interact symbiotically with nitrogen-fixing actinomycetes, particularly Frankia species. Flavonoids also play important roles in these interactions. Flavonoids were shown to accumulate in actinorhizal nodules (Laplaze et al. 1999). Among the flavonoid-like compounds extracted from red alder (*Alnus rubra*), one possible flavanone enhanced nodulation, whereas two other compounds were shown to act as inhibitors (Benoit and Berry 1997). To assess the roles of flavonoids in Frankia-actinorhizal symbiosis, eight flavonoid compounds were extracted from the fruits of *Myrica gale*, with two dihydrochalcone compounds found to enhance the growth and nitrogen fixation of compatible, but not incompatible, Frankia strains (Popovici et al. 2010). Transcriptome analysis revealed that genes involved in flavonoid synthesis and auxin transport were activated in the roots of *C. glauca*, suggesting the important roles of flavonoids and auxin during the Frankia infection process and nodule organogenesis (Hoche et al. 2011). The signaling pathway involved in Frankia-actinorhizal symbiosis has not been fully characterized, but transcriptome analysis showed that similar expression patterns of symbiotic genes in legume and actinorhizal plants, suggesting the involvement of a common or similar pathway for actinorhizal symbiosis.

Interaction with arbuscular mycorrhizal fungi

Mycorrhizal fungi are major components of the soil microbial community, contributing to the transfer of nutrients, especially phosphorus, from soils to plants. Among the mycorrhizal fungi, arbuscular mycorrhizal fungi (AMF) symbiotically interact with around 80% of plant species widely distributed in the terrestrial

ecosystem (Parniske 2008), whereas families such as the Brassicaceae and Chenopodiaceae are not hosts of these fungi (Smith and Read 1997). Fossil records have suggested that arbuscular mycorrhizal symbiosis originated at least 420 to 460 million years ago, coinciding with the origin of the first terrestrial plants, suggesting that ancestral AMF was involved in land plant survival (Redecker et al. 2000; Remy et al. 1994; Simon et al. 1993). Compounds in root exudates have been shown to facilitate interactions with AMF. Using the root exudates of *L. japonicus* grown in nutrient solution, the chemical structure of the signaling compound was identified as a strigolactone (Akiyama et al. 2005), previously found to be a seed germination factor for parasitic weeds such as *Striga* and *Orobanche*.

Various studies have analyzed flavonoid functions in interactions with AMF. Flavonoids were shown to stimulate spore germination, hyphal branching and root colonization (Kikuchi et al. 2007; Scervino et al. 2005, 2007; Siqueira et al. 1991; Steinkellner et al. 2007). Among flavonoids in tomato root exudates, chrysin, luteolin and morin increased the number of entry points and the root colonization of *Gigaspora rosea*, *Gigaspora margarita*, *Glomus mosseae*, and *Glomus intraradices*, whereas kaempferol, isorhamnetin, and rutin showed no effect (Scervino et al. 2007). In addition, a *M. truncatula* mutant with coumestrol accumulation was found to be hyperinfected by AMF (Morandi et al. 2009). These results suggest that flavonoids have positive effects on mycorrhizal symbiosis, but are not considered essential in mediating symbiosis with AMF, inasmuch as root exudates of maize mutants deficient in chalcone synthase exhibited branching activity on hyphae of germinating spores similar to wild type, with both equally colonized with AMF (Becard et al. 1995; Buee et al. 2000). Various phytochemicals, not only flavonoids, may have critical functions in AMF, depending on the plant species. During infection, flavonoids are thought to regulate plant defense reactions, but flavonoid pathway induction during AMF infection was lower than that observed during pathogen infection (Morandi 1996). Flavonoids in the white lupin (*Lupinus albus*), which does not symbiotically interact with AMF, were shown to inhibit interactions with AMF. Moreover, pyranaisoflavones such as licoisoflavone B, sophoraisoflavone A, alpinumisoflavone were found to be strong inhibitors of hyphal branching and germ tube growth (Akiyama et al. 2010).

Interactions with plant growth promoting rhizobacteria

Rhizospheres contain several microorganisms that promote plant growth and health through direct or indirect mechanisms. Beneficial plant-microbe interactions include symbiosis with rhizobia and AMF, as shown above, forming a special structure in roots.

In some cases, these interactions are associated with a broad range of bacteria colonizing the surface of roots or root tissues, mainly in apoplasts. The latter group of bacteria, which do not form special structures, are called plant growth promoting rhizobacteria (PGPR) and have been found to stimulate the growth and health of host plants via various mechanisms, such as phosphate solubilization, siderophore production, phytohormone production, inhibition of pathogen infection, and induction of plant defenses (Vacheron et al. 2013). Many plants, both monocots and dicots, act as hosts for these bacteria, and many bacterial genera, such as *Pseudomonas*, *Azospirillum* and *Bacillus*, and even those containing pathogenic bacteria, function as PGPR. PGPR flourish around the roots due to the exudation of plant metabolites and mucilages. Because plants secrete large amounts of sugars, organic acids and amino acids into the rhizosphere, PGPR utilize these metabolites as nutrients. In addition, PGPR such as *Pseudomonas* and *Bacillus* utilize flavonoids as carbon sources. In *Pseudomonas putida*, quercetin is metabolized to form phloroglucinol and 3,4-dihydroxy cinnamic acid, with the latter further metabolized to protocatechuic acid (Pillai and Swarup 2002). These catabolites and intermediates are secreted by these bacteria, making them secondary metabolites supplied to the rhizosphere.

Pseudomonas fluorescens CHA0 produces the antifungal compounds 2,4-diacetylphloroglucinol and pyoluteorin (Keel et al. 1992; Maurhofer et al. 1994), which contribute to the biocontrol activity of this bacterium. Flavonoids in root exudates have been reported to modulate the expression of *phlA* and *pltA*, genes involved in the synthesis of 2,4-diacetylphloroglucinol and pyoluteorin, respectively. Of 63 plant compounds, including flavonoids and other phenolic compounds, 25 were found to alter the expression of *phlA*, the 2,4-diacetylphloroglucinol biosynthesis gene, and 27 compounds modulated the expression of *pltA*, the pyoluteorin biosynthesis gene (de Werra et al. 2011). These results suggest that phenolic compounds in root exudates are important in regulating the production of antifungal compounds and biocontrol activity.

Relatively little is known about the impact of PGPR on the metabolism of host plants; however, several reports showed that the presence of PGPR altered flavonoid profiles in root exudates (for review see Shaw et al. 2006). For example, *Chryseobacterium balustinum* Aur9 is a PGPR that promotes nitrogen fixation by rhizobia and increases total nitrogen concentration in *Lupinus albus* and soybeans (Garcia et al. 2004; Manero et al. 2003). Under sterile conditions, various flavonoids, such as apigenin, quercetin, naringenin, daidzein, genistein and isoliquiritigenin, were detected in the root exudates of soybeans; however, quercetin and naringenin were not detected in root exudates of soybeans inoculated with

C. balustinum (Dardanelli et al. 2010). Inoculation of *C. balustinum* may have altered the expression of genes involved in the synthesis, accumulation and secretion of flavonoids into the rhizosphere, whereas the disappearance of these flavonoids may indicate their catabolism by *C. balustinum*. However, the ability of *C. balustinum* to utilize these flavonoids has not yet been determined. *C. balustinum* inoculation of the common bean resulted in the disappearance from these exudates of quercetin and isoliquiritigenin, but not of naringenin (Dardanelli et al. 2012). In contrast, apigenin, which was absent from the root exudates of control plants, was present after *C. balustinum* inoculation.

Interactions with pathogens

Plants produce antimicrobial compounds in response to attacks by pathogenic microorganisms. These compounds are called phytoalexins, with many of these compounds being flavonoids. In legume plants, most phytoalexins are isoflavonoids, such as pterocarpan and isoflavanes (Aoki et al. 2000). For example, pea (*Pisum sativum*) produces maackiain and pisatin, which have antifungal activity against *Nectria haematococca*. Maackiain produced by

alfalfa and pea also inhibited the oomycete pathogen *Pythium graminicola* (Jimenez-Gonzalez et al. 2008) and the fungal pathogen *Rhizoctonia solani* (Guenoune et al. 2001; Pueppke and Vanetten 1974). *L. japonicus* contains the phytoalexin vestitol, with most of the genes involved in its synthesis identified (Shimada et al. 2007). In addition to its activity against microbial pathogens, vestitol was secreted into the rhizosphere upon the intrusion of parasitic weed (*Striga hermonthica*), acting as a chemical barrier (Ueda and Sugimoto 2010).

Glyceollins, a family of prenylated pterocarpan derived from daidzein, are specifically induced in soybeans response to pathogens such as *Phytophthora sojae* and *Macrophomina phaseolina* (Ng et al. 2011). The crucial step in glyceollin biosynthesis is the prenylation of glycinol, a reaction catalyzed by the prenyltransferases G2DT and G4DT (Akashi et al. 2009). Glyceollin I is biosynthesized through prenylation at the 4 position, and glyceollins II and III are biosynthesized through prenylation at the 2 position. Yeast extract treatment of cultured soybean cells induced the accumulation of glyceollins I and III, as well as the precursor glycinol, and induced the expression of the genes involved in glyceollin

Figure 3. Biosynthesis of glyceollin I in soybean cells. Putative transport processes are also shown. CYP represents cytochrome P450. Additional CYP is involved in the 2'-hydroxylation of daidzein to form 2'-hydroxydaidzein, but not included in this figure due to space limitation.

biosynthesis (Akashi et al. 2009). G4DT was shown to localize at the plastids (Akashi et al. 2009), similar to other flavonoid prenyltransferases identified to date (Yazaki et al. 2009). However, glycinol is biosynthesized by 3,9-dihydroxypterocarpan 6a-hydroxylase, a cytochrome P450 localized at the ER (Schopfer et al. 1998), suggesting an as yet uncharacterized mechanism of glycinol transport into the plastids. Moreover, 4-dimethylallylglycinol, the product of G4DT, is cyclized by another cytochrome P450 localized at the ER (Welle and Grisebach 1988). These different localizations of successive biosynthetic enzymes suggest possible intracellular movement of intermediates between the ER and plastids (Figure 3), a movement that may also occur during the synthesis of sophoraflavanone G in *S. flavescent* (Sasaki et al. 2008; Yamamoto et al. 2000, 2001). Recently, using mutants in tocopherol biosynthesis and the transorganellar complementation approach, in which plastid-localized enzymes were retargeted to the ER, non-polar substrates in plastids were found to be accessed by ER-localized enzymes (Mehrshahi et al. 2013). These observations suggest interorganellar metabolic pathways, with a variety of compounds including pterocarpan and prenylated flavonoids present in one organelle modifiable by enzymes localized to another organelle.

Phytoalexins are synthesized in response to pathogen infection and exhibit anti-pathogen effects, such as inhibiting the elongation of fungal germ tubes and mycelial growth and inducing hypersensitive response-mediated cell death (Ahuja et al. 2012; Hassan and Mathesius 2012). In addition, several phytoalexins are secreted into the rhizosphere, or at least into the medium. For example, infection of soybean hairy roots with *Fusarium solani*, a soil-born fungal pathogen of soybeans, was found to induce glyceollin secretion (Lozovaya et al. 2004), and fungal infection of the hairy roots of peanuts (*Arachis hypogaea*) was found to induce trans-resveratrol secretion (Medina-Bolivar et al. 2007). However, the induction of phytoalexin secretion may not be a general phenomenon, because the phytoalexin medicarpin was induced in *M. truncatula* but the secretion into the medium was not induced (Farag et al. 2008). Thus, the mechanisms underlying phytoalexin secretion in these plants are tightly regulated, with transporters or other transport mechanisms, such as vesicle transport, being among the most prominent targets in future research.

Interactions with nematodes

Plant-parasitic nematodes like root knot and cyst nematodes cause tremendous damage to agricultural crops. Upon infection by these nematodes, galls or cysts are formed in root tissues, which show characteristic multiple cell divisions and endoreduplication. The involvement of flavonoids in these interactions with

nematodes are not clearly understood, but various phenolic compounds, especially flavonols such as kaempferol and quercetin and their glycosides, were shown to inhibit the chemotaxis and motility of nematodes such as *Meloidogyne incognita* (Wuyts et al. 2006). Invasion with root knot and cyst nematodes was also shown to induce genes involved in flavonoid biosynthesis, with the accumulation of these flavonoids regulating auxin transport (Grunewald et al. 2009; Jones et al. 2007), although conflicting results have been observed using mutants deficient in flavonoid biosynthesis. For instance, none of the flavonoid-deficient mutants of *Arabidopsis* tested, such as *tt4*, *tt5*, *tt6* and their double mutants, showed reductions in the number of infecting nematodes or adult female nematodes in infected roots. Furthermore, several mutant lines produced significantly higher numbers of adult female nematodes, suggesting that these lines permitted the development of more syncytia that supported female development (Jones et al. 2007). In contrast, flavonoid-deficient mutants of *M. truncatula* showed smaller gall formation and fewer cell divisions, although the number of galls was not reduced (Wasson et al. 2009). Although the impact of flavonoids secreted into the rhizosphere on plant-nematode interactions is not completely understood, other plant metabolites secreted into the rhizosphere have significant effects on plant-nematode interactions. Interestingly, for example, β -caryophyllene released from maize roots in response to feeding by a maize pest, the larvae of the beetle *Diabrotica virgifera virgifera*, resulted in the attraction of an entomopathogenic nematode to these maize roots (Rasmann et al. 2005). In contrast to volatile terpenoids, which can move away from plant roots due to their volatility, flavonoids remain in the vicinity of plant roots, suggesting their involvement in attachment and/or infection, rather than in attraction.

Interactions with other plants

Allelopathy is defined as the suppression or inhibition of plant growth and/or development by other plants through the release of organic compounds. In addition to well-characterized allelochemicals, including momilactones A and B from rice and juglone from black walnut (*Juglans nigra*) (Kato-Noguchi and Peters 2013; Soderquist 1973), flavonoids have been implicated as allelochemicals in the rhizosphere. *Dittrichia viscosa* is a ruderal plant, aggressively occupying disturbed areas with allelopathic effects. The effect of flavonoids secreted from the roots of *D. viscosa* on the growth of lettuce seedlings was investigated in detail, with apigenin, 6-methoxy-kaempferol, rhamnetin and isorhamnetin found to reduce the root length of lettuce seedlings, whereas kaempferol-7-methyl ester, 6-methoxy-kaempferol and taxifolin reduced root biomass (Levizou

Figure 4. Schematic overview of the functions of flavonoids in the rhizosphere. Representatives of biologically active flavonoids mediating various biological communications are shown. Pink circles represent nodules.

et al. 2004). In contrast, several flavonoids found in root exudates of *D. viscosa* stimulated lettuce root growth (Levizou et al. 2004). Many types of compounds, including alkaloids, phenolics, cyanoglucosides, polyamines and hydroxyamic acids, have been reported to function as allelochemicals in barley (*Hordeum vulgare*) (Kremer and Ben-Hammouda 2009). The precise functions of barley flavonoids in allelopathy have not yet been determined, although flavonoids such as cyanidin and catechin may inhibit the germination and cell growth of other plants, as well as interfering with auxin function (Kremer and Ben-Hammouda 2009).

As an example of the agricultural application of allelopathic flavonoids, a legume plant, *Desmodium uncinatum*, was used to protect cereal crops from infection by *Striga*, a parasitic weed that causes tremendous crop damage in Africa (Khan et al. 2010). The roots of *D. uncinatum* secrete a C-glycosylflavonoid, isoschaftoside, into the rhizosphere (Hopper et al. 2010). C-glycosylflavonoids are rather unusual metabolites compared with the more common O-glycosylflavonoids. In contrast to O-glycosylflavonoids, which are rapidly hydrolyzed in soil, C-glycosylflavonoids are more stable in soil, better inhibiting *Striga* infection (Hooper et al. 2010).

Conclusion and future perspective

Large gaps exist in our knowledge of flavonoid biosynthesis and their function in the rhizosphere, including the mechanisms of secretion, the rate

of secretion, and the fate and degradation of these compounds. In particular, few genes responsible for the secretion of flavonoids into the rhizosphere have been identified, although various flavonoids are actively secreted by roots. Relatively little is currently known about the kinetics of flavonoid degradation and the fate and further catabolism of these degradation products. Although the mechanism by which flavonoids induce *nod* genes in rhizobia have been well characterized, the mechanisms involved in flavonoid interactions with other organisms are not clearly understood. It is highly probable that the degradation compounds of flavonoids, not the flavonoids themselves, are the key regulators in these interactions. The half-lives of flavonoids in soils are very short and, to our knowledge, the functions of their catabolites in the rhizosphere have not been determined. Meta-metabolome analysis in rhizosphere soil is of particular importance in future research, in addition to metagenome and metatranscriptome analyses. Chemical analyses of the types and amounts of flavonoids and their degradation products are technically difficult, as judged by previous conflicting results, in which different extraction methods, soil types, and microbial communities were used. Therefore, future research should include systematic approaches to soil analysis, especially of microbial communities, in addition to measurements of flavonoids and their degradation products. Determination of the mobility, space distribution and structural changes of flavonoids in correlation with

soil types and microbes in the rhizosphere is crucial in understanding the functions of these plant-derived compounds in biological communications.

Harnessing the beneficial interactions between plants and other organisms to manage nutrition and disease, thus improving the crop yields, is of particular importance in sustainable agriculture, reducing the use of chemical fertilizers and pesticides (Figure 4). Transgenic plants and natural variations with altered flavonoid accumulation and exudation can be used to study the effects of flavonoid exudation in the rhizosphere. Flavonoid accumulation was recently shown to enhance plant tolerance to drought stress (Nakabayashi et al. 2014). Thus, molecular breeding methods that increase flavonoid accumulation and/or alter flavonoid composition are promising for crop improvements. The rhizosphere is recognized as being a frontier in plant breeding (Wissuwa et al. 2009), as shown by the soybean breeding program in Brazil, in which the disuse of nitrogen fertilizers led to a *Bradyrhizobium* strain more efficient in nitrogen fixation (Drinkwater and Snapp 2007). Manipulation of plant metabolites, especially those secreted into the rhizosphere, could be the basis for the next-generation rhizosphere breeding program for better utilization of rhizosphere biological communication.

Plants can be grown under various conditions, including axenic culture in the laboratory; in vermiculites; in commercial or field soils in pots, either in growth chambers or greenhouses; or in fields. The biological interactions observed in tightly controlled, or even highly artificial, conditions cannot be readily reproduced under field conditions, with this being one of the biggest obstacles in applying scientific insights to agriculture. To pave the way towards applying molecular components to mediate the beneficial interactions, basic mechanisms discovered in studies performed in well-controlled environments should further be tested under greenhouse conditions, followed by tests in the field. The *raison d'être* of plant specialized metabolites was first proposed as being aerial biological communication (Fraenkel 1959). Half a half century later, substantial evidence has accumulated to show that their *raison d'être* is biological communications in rhizospheres, suggesting their potential application in agricultural practice.

Acknowledgements

We gratefully acknowledge a grant from the Ministry of Agriculture, Forestry, and Fisheries of Japan (Genomics-based Technology for Agricultural Improvement, SFC2001) (AS), and JSPS KAKENHI Grant Number 26712013 (AS).

References

Ahuja I, Kissen R, Bones AM (2012) Phytoalexins in defense against pathogens. *Trends Plant Sci* 17: 73–90

- Akashi T, Sasaki K, Aoki T, Ayabe S, Yazaki K (2009) Molecular cloning and characterization of a cDNA for pterocarpan 4-dimethylallyltransferase catalyzing the key prenylation step in the biosynthesis of glyceollin, a soybean phytoalexin. *Plant Physiol* 149: 683–693
- Akiyama K, Matsuzaki K, Hayashi H (2005) Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. *Nature* 435: 824–827
- Akiyama K, Tanigawa F, Kashiwara T, Hayashi H (2010) Lupin pyranosylflavones inhibiting hyphal development in arbuscular mycorrhizal fungi. *Phytochemistry* 71: 1865–1871
- Aoki T, Akashi T, Ayabe S (2000) Flavonoids of leguminous plants: Structure, biological activity, and biosynthesis. *J Plant Res* 113: 475–488
- Badri DV, Chaparro JM, Zhang R, Shen Q, Vivanco JM (2013) Application of natural blends of phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *J Biol Chem* 288: 4502–4512
- Badri DV, Quintana N, El Kassis EG, Kim HK, Choi YH, Sugiyama A, Verpoorte R, Martinoia E, Manter DK, Vivanco JM (2009) An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota. *Plant Physiol* 151: 2006–2017
- Barto EK, Cipollini D (2009) Half-lives and field soil concentrations of *Alliaria petiolata* secondary metabolites. *Chemosphere* 76: 71–75
- Barz W, Welle R (1992) Biosynthesis and metabolism of isoflavones and pterocarpan phytoalexins in chickpea, soybean and phytopathogenic fungi. Plenum Press Div Plenum Publishing Corp, New York
- Becard G, Taylor LP, Douds DD Jr, Pfeffer PE, Doner LW (1995) Flavonoids are not necessary plant signal compounds in arbuscular mycorrhizal symbioses. *Mol Plant Microbe Interact* 8: 252–258
- Benoit LE, Berry AM (1997) Flavonoid-like compounds from seeds of red alder (*Alnus rubra*) influence host nodulation by *Frankia* (*Actinomycetales*). *Physiol Plant* 99: 588–593
- Buee M, Rossignol M, Jauneau A, Ranjeva R, Becard G (2000) The pre-symbiotic growth of arbuscular mycorrhizal fungi is induced by a branching factor partially purified from plant root exudates. *Mol Plant Microbe Interact* 13: 693–698
- Carlsen SCK, Pedersen HA, Spliid NH, Fomsgaard IS (2012) Fate in soil of flavonoids released from white clover (*Trifolium repens* L.). *Appl Environ Soil Sci* 2012: 743413
- Cesco S, Mimmo T, Tonon G, Tomasi N, Pinton R, Terzano R, Neumann G, Weisskopf L, Renella G, Landi L, et al. (2012) Plant-borne flavonoids released into the rhizosphere: impact on soil bio-activities related to plant nutrition. A review. *Biol Fertil Soils* 48: 123–149
- Cesco S, Neumann G, Tomasi N, Pinton R, Weisskopf L (2010) Release of plant-borne flavonoids into the rhizosphere and their role in plant nutrition. *Plant Soil* 329: 1–25
- Chaparro JM, Badri DV, Bakker MG, Sugiyama A, Manter DK, Vivanco JM (2013) Root exudation of phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and correlate with soil microbial functions. *PLoS ONE* 8: e55731
- Chaparro JM, Badri DV, Vivanco JM (2014) Rhizosphere microbiome assemblage is affected by plant development. *ISME J* 8: 790–803
- Cheynier V, Comte G, Davies KM, Lattanzio V, Martens S (2013)

- Plant phenolics: Recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiol Biochem* 72: 1–20
- Dakora FD, Joseph CM, Phillips DA (1993) Alfalfa (*Medicago sativa* L.) root exudates contain isoflavonoids in the presence of *Rhizobium meliloti*. *Plant Physiol* 101: 819–824
- Dardanelli MS, Fernandez de Cordoba FJ, Espuny MR, Rodriguez Carvajal MA, Soria Diaz ME, Gil Serrano AM, Okon Y, Megias M (2008) Effect of *Azospirillum brasilense* coinoculated with *Rhizobium* on *Phaseolus vulgaris* flavonoids and Nod factor production under salt stress. *Soil Biol Biochem* 40: 2713–2721
- Dardanelli MS, Fernandez de Cordoba FJ, Estevez J, Contreras R, Cubo MT, Rodriguez-Carvajal MA, Gil-Serrano AM, Lopez-Baena FJ, Bellogin R, Manyani H, et al. (2012) Changes in flavonoids secreted by *Phaseolus vulgaris* roots in the presence of salt and the plant growth-promoting rhizobacterium *Chryseobacterium balustinum*. *Appl Soil Ecol* 57: 31–38
- Dardanelli MS, Manyani H, Gonzalez-Barroso S, Rodriguez-Carvajal MA, Gil-Serrano AM, Espuny MR, Lopez-Baena FJ, Bellogin RA, Megias M, Ollero FJ (2010) Effect of the presence of the plant growth promoting rhizobacterium (PGPR) *Chryseobacterium balustinum* Aur9 and salt stress in the pattern of flavonoids exuded by soybean roots. *Plant Soil* 328: 483–493
- de Werra P, Huser A, Tabacchi R, Keel C, Maurhofer M (2011) Plant- and microbe-derived compounds affect the expression of genes encoding antifungal compounds in a pseudomonad with biocontrol activity. *Appl Environ Microbiol* 77: 2807–2812
- Dixon RA, Strack D (2003) Phytochemistry meets genome analysis, and beyond. *Phytochemistry* 62: 815–816
- Djordjevic MA, Redmond JW, Batley M, Rolfe BG (1987) Clovers secrete specific phenolic compounds which either stimulate or repress nod gene expression in *Rhizobium trifolii*. *EMBO J* 6: 1173–1179
- Drinkwater LE, Snapp SS (2007) Understanding and managing the rhizosphere in agroecosystems. Academic Press
- Estabrook EM, Sengupta-Gopalan C (1991) Differential expression of phenylalanine ammonia-lyase and chalcone synthase during soybean nodule development. *Plant Cell* 3: 299–308
- Falcone Ferreyra ML, Rius SP, Casati P (2012) Flavonoids: Biosynthesis, biological functions, and biotechnological applications. *Front Plant Sci* 3: 222
- Farag MA, Huhman DV, Dixon RA, Sumner LW (2008) Metabolomics reveals novel pathways and differential mechanistic and elicitor-specific responses in phenylpropanoid and isoflavonoid biosynthesis in *Medicago truncatula* cell cultures. *Plant Physiol* 146: 387–402
- Ferrer JL, Austin MB, Stewart C Jr, Noel JP (2008) Structure and function of enzymes involved in the biosynthesis of phenylpropanoids. *Plant Physiol Biochem* 46: 356–370
- Fraenkel GS (1959) The raison d'être of secondary plant substances; These odd chemicals arose as a means of protecting plants from insects and now guide insects to food. *Science* 129: 1466–1470
- Furubayashi A, Hiradate S, Fujii Y (2007) Role of catechol structure in the adsorption and transformation reactions of L-DOPA in soils. *J Chem Ecol* 33: 239–250
- Garcia-Lafuente A, Guillaumon E, Villares A, Rostagno MA, Martinez JA (2009) Flavonoids as anti-inflammatory agents: Implications in cancer and cardiovascular disease. *Inflamm Res* 58: 537–552
- Garcia JAL, Probanza A, Ramos B, Colon Flores JJ, Gutierrez Manero FJ (2004) Effects of plant growth promoting rhizobacteria (PGPRs) on the biological nitrogen fixation, nodulation, and growth of *Lupinus albus* L. cv. *multolupa*. *Eng Life Sci* 4: 71–77
- Grunewald W, van Noorden G, van Isterdael G, Beeckman T, Gheysen G, Mathesius U (2009) Manipulation of auxin transport in plant roots during *Rhizobium* symbiosis and nematode parasitism. *Plant Cell* 21: 2553–2562
- Guenoune D, Galili S, Phillips DA, Volpin H, Chet I, Okon Y, Kapulnik Y (2001) The defense response elicited by the pathogen *Rhizoctonia solani* is suppressed by colonization of the AM-fungus *Glomus intraradices*. *Plant Sci* 160: 925–932
- Guo ZY, Kong CH, Wang JG, Wang YF (2011) Rhizosphere isoflavones (daidzein and genistein) levels and their relation to the microbial community structure of mono-cropped soybean soil in field and controlled conditions. *Soil Biol Biochem* 43: 2257–2264
- Hartmann A, Rothballer M, Schmid M (2008) Lorenz Hiltner, a pioneer in rhizosphere microbial ecology and soil bacteriology research. *Plant Soil* 312: 7–14
- Hartwig UA, Phillips DA (1991) Release and modification of nod-gene-inducing flavonoids from alfalfa seeds. *Plant Physiol* 95: 804–807
- Hassan S, Mathesius U (2012) The role of flavonoids in root-rhizosphere signalling: Opportunities and challenges for improving plant-microbe interactions. *J Exp Bot* 63: 3429–3444
- Hocher V, Alloisio N, Auguy F, Fournier P, Dumas P, Pujic P, Gherbi H, Queiroux C, Da Silva C, Wincker P, et al. (2011) Transcriptomics of actinorhizal symbioses reveals homologs of the whole common symbiotic signaling cascade. *Plant Physiol* 156: 700–711
- Hoerger CC, Praplan AP, Becker L, Wettstein FE, Hungerbuhler K, Bucheli TD (2011a) Quantification of five isoflavones and coumestrol in various solid agroenvironmental matrices using C-13(3)-labeled internal standards. *J Agric Food Chem* 59: 847–856
- Hoerger CC, Wettstein FE, Bachmann HJ, Hungerbuhler K, Bucheli TD (2011b) Occurrence and mass balance of isoflavones on an experimental grassland field. *Environ Sci Technol* 45: 6752–6760
- Hooper AM, Tsanuo MK, Chamberlain K, Tittcomb K, Scholes J, Hassanali A, Khan ZR, Pickett JA (2010) Isochaftoside, a C-glycosylflavonoid from *Desmodium uncinatum* root exudate, is an allelochemical against the development of *Striga*. *Phytochemistry* 71: 904–908
- Jimenez-Gonzalez L, Alvarez-Corral M, Muñoz-Dorado M, Rodriguez-Garcia I (2008) Pterocarpanes: Interesting natural products with antifungal activity and other biological properties. *Phytochem Rev* 7: 125–154
- Jones JT, Furlanetto C, Phillips MS (2007) The role of flavonoids produced in response to cyst nematode infection of *Arabidopsis thaliana*. *Nematology* 9: 671–677
- Kato-Noguchi H, Peters RJ (2013) The role of momilactones in rice allelopathy. *J Chem Ecol* 39: 175–185
- Keel C, Schneider U, Maurhofer M, Voisard C, Laville J, Burger U, Wirthner P, Haas D, Defago G (1992) Suppression of root diseases by *Pseudomonas fluorescens* CHA0: Importance of the bacterial secondary metabolite 2,4-diacetylphloroglucinol. *Mol Plant Microbe Interact* 5: 4–13
- Khan ZR, Midega CAO, Bruce TJA, Hooper AM, Pickett JA (2010) Exploiting phytochemicals for developing a 'push-pull' crop protection strategy for cereal farmers in Africa. *J Exp Bot* 61: 4185–4196
- Kikuchi K, Matsushita N, Suzuki K, Hogetsu T (2007) Flavonoids induce germination of basidiospores of the ectomycorrhizal fungus *Suillus bovinus*. *Mycorrhiza* 17: 563–570

- Kosslak RM, Bookland R, Barkei J, Paaren HE, Appelbaum ER (1987) Induction of *Bradyrhizobium japonicum* common nod genes by isoflavones isolated from *Glycine max*. *Proc Natl Acad Sci USA* 84: 7428–7432
- Kremer RJ, Ben-Hammouda M (2009) Allelopathic plants. 19. Barley (*Hordeum vulgare* L.). *Allelopathy J* 24: 225–241
- Laplaze L, Gherbi H, Frutz T, Pawlowski K, Franche C, Macheix JJ, Auguy F, Bogusz D, Duhoux E (1999) Flavan-containing cells delimit *Frankia*-infected compartments in *Casuarina glauca* nodules. *Plant Physiol* 121: 113–122
- Leon-Barrios M, Dakora FD, Joseph CM, Phillips DA (1993) Isolation of *Rhizobium meliloti* nod gene inducers from alfalfa rhizosphere soil. *Appl Environ Microbiol* 59: 636–639
- Levizou E, Karageorgou P, Petropoulou Y, Grammatikopoulos G, Manetas Y (2004) Induction of ageotropic response in lettuce radical growth by epicuticular flavonoid aglycones of *Dittrichia viscosa*. *Biol Plant* 48: 305–307
- Lozovaya VV, Lygin AV, Zernova OV, Li S, Hartman GL, Widholm JM (2004) Isoflavonoid accumulation in soybean hairy roots upon treatment with *Fusarium solani*. *Plant Physiol Biochem* 42: 671–679
- Manero FJG, Probanza A, Ramos B, Colon Flores JJ, Garcia JAL (2003) Effects of culture filtrates of rhizobacteria isolated from wild lupine on germination, growth, and biological nitrogen fixation of lupine seedlings. *J Plant Nutr* 26: 1101–1115
- Mathesius U, Schlaman HRM, Spaijk HP, Of Sautter C, Rolfe BG, Djordjevic MA (1998) Auxin transport inhibition precedes root nodule formation in white clover roots and is regulated by flavonoids and derivatives of chitin oligosaccharides. *Plant J* 14: 23–34
- Maurhofer M, Keel C, Haas D, Defago G (1994) Pyoluteorin production by *Pseudomonas fluorescens* strain CHA0 is involved in the suppression of *Pythium* damping-off of cress but not of cucumber. *Eur J Plant Pathol* 100: 221–232
- Medina-Bolivar F, Condori J, Rimando AM, Hubstenberger J, Shelton K, O'Keefe SF, Bennett S, Dolan MC (2007) Production and secretion of resveratrol in hairy root cultures of peanut. *Phytochemistry* 68: 1992–2003
- Mehrshahi P, Stefano G, Andaloro JM, Brandizzi F, Froehlich JE, DellaPenna D (2013) Transorganellar complementation redefines the biochemical continuity of endoplasmic reticulum and chloroplasts. *Proc Natl Acad Sci USA* 110: 12126–12131
- Miadokova E (2009) Isoflavonoids—an overview of their biological activities and potential health benefits. *Interdiscip Toxicol* 2: 211–218
- Morandi D (1996) Occurrence of phytoalexins and phenolic compounds in endomycorrhizal interactions, and their potential role in biological control. *Plant Soil* 185: 241–251
- Morandi D, le Signor C, Gianinazzi-Pearson V, Duc G (2009) A *Medicago truncatula* mutant hyper-responsive to mycorrhiza and defective for nodulation. *Mycorrhiza* 19: 435–441
- Mougel C, Offre P, Ranjard L, Corberand T, Gamalero E, Robin C, Lemancau P (2006) Dynamic of the genetic structure of bacterial and fungal communities at different developmental stages of *Medicago truncatula* Gaertn. cv. Jemalong line J5. *New Phytol* 170: 165–175
- Nakabayashi R, Yonekura-Sakakibara K, Urano K, Suzuki M, Yamada Y, Nishizawa T, Matsuda F, Kojima M, Sakakibara H, Shinozaki K, et al. (2014) Enhancement of oxidative and drought tolerance in *Arabidopsis* by overaccumulation of antioxidant flavonoids. *Plant J* 77: 367–379
- Nakagawa T, Kaku H, Shimoda Y, Sugiyama A, Shimamura M, Takanashi K, Yazaki K, Aoki T, Shibuya N, Kouchi H (2011) From defense to symbiosis: Limited alterations in the kinase domain of LysM receptor-like kinases are crucial for evolution of legume-*Rhizobium* symbiosis. *Plant J* 65: 169–180
- Ng TB, Ye XJ, Wong JH, Fang EF, Chan YS, Pan W, Ye XY, Sze SC, Zhang KY, Liu F, Wang HX (2011) Glyceollin, a soybean phytoalexin with medicinal properties. *Appl Microbiol Biotechnol* 90: 59–68
- Noguchi A, Saito A, Homma Y, Nakao M, Sasaki N, Nishino T, Takahashi S, Nakayama T (2007) A UDP-glucose:Isoflavone 7-O-glucosyltransferase from the roots of soybean (*Glycine max*) seedlings. *J Biol Chem* 282: 23581–23590
- Parniske M (2008) Arbuscular mycorrhiza: The mother of plant root endosymbioses. *Nat Rev Microbiol* 6: 763–775
- Peck MC, Fisher RF, Long SR (2006) Diverse flavonoids stimulate NodD1 binding to nod gene promoters in *Sinorhizobium meliloti*. *J Bacteriol* 188: 5417–5427
- Peters NK, Frost JW, Long SR (1986) A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science* 233: 977–980
- Pichersky E, Lewinsohn E (2011) Convergent evolution in plant specialized metabolism. *Annu Rev Plant Biol* 62: 549–566
- Pillai BV, Swarup S (2002) Elucidation of the flavonoid catabolism pathway in *Pseudomonas putida* PML2 by comparative metabolic profiling. *Appl Environ Microbiol* 68: 143–151
- Pollock JL, Callaway RM, Thelen GC, Holben WE (2009) Catechin-metal interactions as a mechanism for conditional allelopathy by the invasive plant *Centaurea maculosa*. *J Ecol* 97: 1234–1242
- Popovici J, Comte G, Bagnarol E, Alloisio N, Fournier P, Bellvert F, Bertrand C, Fernandez MP (2010) Differential effects of rare specific flavonoids on compatible and incompatible strains in the *Myrica gale*-*Frankia* actinorhizal symbiosis. *Appl Environ Microbiol* 76: 2451–2460
- Popp C, Ott T (2011) Regulation of signal transduction and bacterial infection during root nodule symbiosis. *Curr Opin Plant Biol* 14: 458–467
- Pueppke SG, Vanetten HD (1974) Pisatin accumulation and lesion development in peas infected with *Aphanomyces euteiches*, *Fusarium solani* f. sp. pisi, or *Rhizoctonia solani*. *Phytopathology* 64: 1433–1440
- Rasmann S, Kollner TG, Degenhardt J, Hiltbold I, Toepfer S, Kuhlmann U, Gershenzon J, Turlings TCJ (2005) Recruitment of entomopathogenic nematodes by insect-damaged maize roots. *Nature* 434: 732–737
- Redecker D, Kodner R, Graham LE (2000) Glomalean fungi from the Ordovician. *Science* 289: 1920–1921
- Redmond JW, Batley M, Djordjevic MA, Innes RW, Kuempel PL, Rolfe BG (1986) Flavones induce expression of nodulation genes in *Rhizobium*. *Nature* 323: 632–635
- Remy W, Taylor TN, Hass H, Kerp H (1994) Four hundred-million-year-old vesicular arbuscular mycorrhizae. *Proc Natl Acad Sci USA* 91: 11841–11843
- Saito K, Yonekura-Sakakibara K, Nakabayashi R, Higashi Y, Yamazaki M, Tohge T, Fernie AR (2013) The flavonoid biosynthetic pathway in *Arabidopsis*: Structural and genetic diversity. *Plant Physiol Biochem* 72: 21–34
- Sasaki K, Mito K, Ohara K, Yamamoto H, Yazaki K (2008) Cloning and characterization of naringenin 8-prenyltransferase, a flavonoid-specific prenyltransferase of *Sophora flavescens*. *Plant Physiol* 146: 1075–1084
- Savoure A, Magyar Z, Pierre M, Brown S, Schultze M, Dudits D, Kondorosi A, Kondorosi E (1994) Activation of the cell cycle

- machinery and the isoflavonoid biosynthesis pathway by active *Rhizobium meliloti* Nod signal molecules in *Medicago microcallus* suspensions. *EMBO J* 13: 1093–1102
- Scervino JM, Ponce MA, Erra-Bassells R, Bompadre J, Vierheilig H, Ocampo JA, Godeas A (2007) The effect of flavones and flavonols on colonization of tomato plants by arbuscular mycorrhizal fungi of the genera *Gigaspora* and *Glomus*. *Can J Microbiol* 53: 702–709
- Scervino JM, Ponce MA, Erra-Bassells R, Vierheilig H, Ocampo JA, Godeas A (2005) Flavonoids exhibit fungal species and genus specific effects on the presymbiotic growth of *Gigaspora* and *Glomus*. *Mycol Res* 109: 789–794
- Schopfer CR, Kochs G, Lottspeich F, Ebel J (1998) Molecular characterization and functional expression of dihydroxypterocarpan 6a-hydroxylase, an enzyme specific for pterocarpanoid phytoalexin biosynthesis in soybean (*Glycine max* L.). *FEBS Lett* 432: 182–186
- Shaw LJ, Hooker JE (2008) The fate and toxicity of the flavonoids naringenin and formononetin in soil. *Soil Biol Biochem* 40: 528–536
- Shaw LJ, Morris P, Hooker JE (2006) Perception and modification of plant flavonoid signals by rhizosphere microorganisms. *Environ Microbiol* 8: 1867–1880
- Shimada N, Sato S, Akashi T, Nakamura Y, Tabata S, Ayabe S, Aoki T (2007) Genome-wide analyses of the structural gene families involved in the legume-specific 5-deoxyisoflavonoid biosynthesis of *Lotus japonicus*. *DNA Res* 14: 25–36
- Shitan N, Yazaki K (2013) New insights into the transport mechanisms in plant vacuoles. *Int Rev Cell Mol Biol* 305: 383–433
- Simon L, Bousquet J, Levesque RC, Lalonde M (1993) Origin and diversification of endomycorrhizal fungi and coincidence with vascular land plants. *Nature* 363: 67–69
- Siqueira JO, Safir GR, Nair MG (1991) Stimulation of vesicular-arbuscular mycorrhizae formation and growth of white clover by flavonoid compounds. *New Phytol* 118: 87–93
- Smith SE, Read DJ (1997) Mycorrhizal symbiosis. 2nd Edition. Academic Press, San Diego, CA
- Soderquist CJ (1973) Juglone and allelopathy. *J Chem Educ* 50: 782–783
- Steinkellner S, Lendzemo V, Langer I, Schweiger P, Khaosaad T, Toussaint JP, Vierheilig H (2007) Flavonoids and strigolactones in root exudates as signals in symbiotic and pathogenic plant-fungus interactions. *Molecules* 12: 1290–1306
- Sugiyama A, Shitan N, Yazaki K (2007) Involvement of a soybean ATP-binding cassette-type transporter in the secretion of genistein, a signal flavonoid in legume-*Rhizobium* symbiosis. *Plant Physiol* 144: 2000–2008
- Sugiyama A, Ueda Y, Zushi T, Takase H, Yazaki K (2014) Changes in the bacterial community of soybean rhizospheres during growth in the field. *PLoS ONE* 9: e100709
- Sugiyama A, Yazaki K (2012) Root exudates of legume plants and their involvement in interactions with soil microbes. In: Vivanco JM, Baluska F (eds) *Secretions and Exudates in Biological Systems*. Springer, pp 27–48
- Suzuki S, Koeduka T, Sugiyama A, Yazaki K, Umezawa T Microbial production of plant specialized metabolites. *Plant Biotechnol* (in press)
- Suzuki H, Nishino T, Nakayama T (2007) cDNA cloning of a BAHF acyltransferase from soybean (*Glycine max*): Isoflavone 7-O-glucoside-6"-O-malonyltransferase. *Phytochemistry* 68: 2035–2042
- Suzuki H, Takahashi S, Watanabe R, Fukushima Y, Fujita N, Noguchi A, Yokoyama R, Nishitani K, Nishino T, Nakayama T (2006) An isoflavone conjugate-hydrolyzing beta-glucosidase from the roots of soybean (*Glycine max*) seedlings: Purification, gene cloning, phylogenetics, and cellular localization. *J Biol Chem* 281: 30251–30259
- Toussaint JP, Pham TTM, Barriault D, Sylvestre M (2012) Plant exudates promote PCB degradation by a rhodococcal rhizobacteria. *Appl Microbiol Biotechnol* 95: 1589–1603
- Ueda H, Sugimoto Y (2010) Vestitol as a chemical barrier against intrusion of parasitic plant *Striga hermonthica* into *Lotus japonicus* roots. *Biosci Biotechnol Biochem* 74: 1662–1667
- Vacheron J, Desbrosses G, Bouffaud ML, Touraine B, Moenne-Loccoz Y, Muller D, Legendre L, Wisniewski-Dye F, Prigent-Combaret C (2013) Plant growth-promoting rhizobacteria and root system functioning. *Front Plant Sci* 4: 356
- Wang D, Yang S, Tang F, Zhu H (2012) Symbiosis specificity in the legume: rhizobial mutualism. *Cell Microbiol* 14: 334–342
- Wasson AP, Pellerone FI, Mathesius U (2006) Silencing the flavonoid pathway in *Medicago truncatula* inhibits root nodule formation and prevents auxin transport regulation by rhizobia. *Plant Cell* 18: 1617–1629
- Wasson AP, Ramsay K, Jones MGK, Mathesius U (2009) Differing requirements for flavonoids during the formation of lateral roots, nodules and root knot nematode galls in *Medicago truncatula*. *New Phytol* 183: 167–179
- Welle R, Grisebach H (1988) Induction of phytoalexin synthesis in soybean: Enzymatic cyclization of prenylated pterocarpan to glyceollin isomers. *Arch Biochem Biophys* 263: 191–198
- Werner D (2007) Molecular biology and ecology of the rhizobia-legume symbiosis. In: Pinton R, Varanini Z, Nannipiero P (ed) *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*. CRC Press, New York, pp 237–266
- Weston LA, Mathesius U (2013) Flavonoids: their structure, biosynthesis and role in the rhizosphere, including allelopathy. *J Chem Ecol* 39: 283–297
- Wissuwa M, Mazzola M, Picard C (2009) Novel approaches in plant breeding for rhizosphere-related traits. *Plant Soil* 321: 409–430
- Wuyts N, Swennen R, De Waele D (2006) Effects of plant phenylpropanoid pathway products and selected terpenoids and alkaloids on the behaviour of the plant-parasitic nematodes *Radopholus similis*, *Pratylenchus penetrans* and *Meloidogyne incognita*. *Nematology* 8: 89–101
- Yamamoto H, Senda M, Inoue K (2000) Flavanone 8-dimethylallyltransferase in *Sophora flavescens* cell suspension cultures. *Phytochemistry* 54: 649–655
- Yamamoto H, Yatou A, Inoue K (2001) 8-Dimethylallylnaringenin 2'-hydroxylase, the crucial cytochrome P450 mono-oxygenase for lavandulylated flavanone formation in *Sophora flavescens* cultured cells. *Phytochemistry* 58: 671–676
- Yazaki K, Sasaki K, Tsurumaru Y (2009) Prenylation of aromatic compounds, a key diversification of plant secondary metabolites. *Phytochemistry* 70: 1739–1745
- Yokota K, Hayashi M (2011) Function and evolution of nodulation genes in legumes. *Cell Mol Life Sci* 68: 1341–1351
- Zhang J, Subramanian S, Stacey G, Yu O (2009) Flavones and flavonols play distinct critical roles during nodulation of *Medicago truncatula* by *Sinorhizobium meliloti*. *Plant J* 57: 171–183
- Zhao J, Dixon RA (2010) The 'ins' and 'outs' of flavonoid transport. *Trends Plant Sci* 15: 72–80