Note

De novo transcriptome analysis of needles of *Thujopsis dolabrata* var. *hondae*

Shiro Suzuki^{1,2}, Hideyuki Suzuki³, Koji Tanaka⁴, Masaomi Yamamura¹,
Daisuke Shibata³, Toshiaki Umezawa^{1,2,5,*}

¹Research Institute of Sustainable Humanosphere, Kyoto University, Uji, Kyoto 611-0011, Japan; ²Institute of Sustainability Science, Kyoto University, Uji, Kyoto 611-0011, Japan; ³Kazusa DNA Research Institute, Kisarazu, Chiba 275-4242, Japan;

⁴Forestry and Forest Products Research Institute, Aomori Prefectural Industrial Research Center, Hiranai, Aomori 039-3321, Japan; ⁵Research Unit for Development of Global Sustainability, Kyoto University, Uji, Kyoto 611-0011, Japan

*E-mail: tumezawa@rish.kyoto-u.ac.jp Tel: +81-774-38-3625 Fax: +81-774-38-3682

Received December 11, 2018; accepted February 20, 2019 (Edited by T. Koezuka)

Abstract Podophyllotoxin is a starting material of the semisynthetic anticancer medicines etoposide, teniposide, and etopophos. The major plant source of podophyllotoxin is rhizomes of *Podophyllum hexandrum*, which is a Himalayan endangered species; therefore, alternative sources of podophyllotoxin or bioproduction systems have been pursued to avoid exploiting this limited natural resource. In this paper, we report de novo transcriptome analysis of *Thujopsis dolabrata* var. *hondae*, which accumulates the podophyllotoxin derivatives (deoxypodophyllotoxin and β -peltatin A methyl ether) in its needles. We analyzed transcriptomes of the *T. dolabrata* var. *hondae* young needles to obtain the sequences that putatively encode O-methyltransferases, cytochrome P450s, and a 2-oxoglutarate dependent dioxygenase because these protein families are responsible for podophyllotoxin-related compound formation in *P. hexandrum*. The resulting transcriptomes contained considerable numbers of coding sequences classified into the three protein families. Our results are a genetic basis for identifying genes involved in the biosynthesis of podophyllotoxin and related compounds and also for future metabolic engineering of podophyllotoxin in heterologous hosts.

Key words: 2-oxoglutarate-dependent dioxygenase, next-generation sequencer, O-methyltransferase, P450, podophyllotoxin.

Podophyllotoxin has been successfully used as the starting material of semisynthetic anticancer drugs, etoposide, teniposide, and etopophos. Of note, etoposide is on the World Health Organization's (WHO) list of essential medicines (Lau and Sattely 2015). Etoposide has been synthesized from podophyllotoxin via demethylation of 4'-methoxyl group on the aromatic E-ring and epimerization/acetonide-glucosylation of 7-hydroxyl group (Lata et al. 2009) (Figure 1). Currently, podophyllotoxin is mainly isolated from rhizomes of an endangered species, *Podophyllum hexandrum* (*Sinopodophyllum hexandrum*; Himalayan mayapple) (Hendrawati et al. 2012; Lata et al. 2009). In addition to *Podophyllum* spp., podophyllotoxin and related compounds are produced in other plant families including Linaceae, Apiaceae, Cupressaceae, and Hernandiaceae (Umezawa 2003a, 2003b). Therefore, much attention has been focused on biosynthesis and bioproduction of podophyllotoxin and related compounds in *Podophyllum* spp. and other plant species

(Bhattacharyya et al. 2016; Kumar 2017; Kumari et al. 2014; Sasheva and Ikonkova 2017; Seegers et al. 2017; Suzuki and Umezawa 2007).

The current biosynthetic pathway of podophyllotoxin-related compounds is illustrated in Figure 2. First, Dewick et al. revealed that yatein is converted to podophyllotoxin via deoxypodophyllotoxin (=desoxypodophyllotoxin or anthricin) (Jackson and Dewick 1984; Kamil and Dewick 1986a) in *Podophyllum* plants. They stated that matairesinol is likely the branch-point compound that affords deoxypodophyllotoxin and 4'-demethylpodophyllotoxin (Broomhead et al. 1991; Dewick 1993; Kamil and Dewick 1986b). However, Kawai et al. (1994, 1999) proposed 4'-demethylyatein formation via thujaplicatin in *Thuja occidentalis*. Xia et al. (2000) showed the formation of 6-O-methylpodophyllotoxin via 7-hydroxymatairesinol. Kranz and Petersen (2003) reported that β -peltatin 6-O-methyltransferase activity gave β -peltatin A methyl ether in *Linum nodiflorum*. In

Abbreviations: 2OGD, 2-oxoglutarate-dependent dioxygenase; IIX, Genome Analyzer IIX; CYP, cytochrome P450; CALDOMT1, coniferaldehyde OMT 1; CCoAOMT, caffeoyl-CoA OMT 1; FLX, Genome Sequencer FLX; OMT, O-methyltransferase.

This article can be found at <http://www.jspcmb.jp/>

Published online June 18, 2019

Figure 1. Chemical synthesis of etoposide from (-)-podophyllotoxin.

2003, Sakakibara et al. administered a series of isotope-labeled dibenzyl butylolactone lignans to *Anthriscus sylvestris* (Lim et al. 1999; Suzuki et al. 2002) and showed that yatein is produced from matairesinol via thujaplicatin, 4'-O-methylthujaplicatin, 4',5'-di-O-methylthujaplicatin in this species. In contrast, Marques et al. (2013) found that a cytochrome P450 enzyme (CYP) in *P. hexandrum*, CYP719A23, catalyzes the conversion of (-)-matairesinol to (-)-pulviatolide. In 2013, Ragamustari et al. found an

Figure 2. Current biosynthetic pathways of podophyllotoxin and related compounds.

O-methyltransferase (AsTJOMT) that catalyzes the conversion of thujaplicatin to 5'-O-methylthujaplicatin in *A. sylvestris*. Using *Agrobacterium*-mediated transient protein expression system in *Nicotiana benthamiana*, Lau and Sattely (2015) reported two O-methyltransferases (OMTs), named OMT3 and OMT1, for O-methylation of (–)-pulvitolide and (–)-5'-demethylatein, two CYPs (CYP71CU1 and CYP82D61) mediating the hydroxylation of (–)-burshehennin and (–)-4'-demethylpodophyllotoxin, one CYP (CYP71BE54) demethylating 4'-methoxyl group of (–)-deoxypodophyllotoxin, and one 2-oxoglutarate-dependent dioxygenase (2OGD) named deoxypodophyllotoxin synthase that catalyzes the conversion of yatein to deoxypodophyllotoxin.

Among Cupressaceae species, *Thujaopsis dolabrata* var. *hondae*, which is endemic to the northern Honshu and southern Hokkaido Islands of Japan (Kanetoshi et al. 1998), is a good source of podophyllotoxin derivatives; needles contain 1.1% of deoxypodophyllotoxin and 0.56% of β -peltatin-A methyl ether in the 70% acetone/water extract (Kanetoshi et al. 1999). In addition to podophyllotoxin derivative accumulation in the needles, because the wood contains the terpenoids β -thujaplicin (hinokitiol) and related compounds, which have high insecticidal and anti-wood-rot activities, the timber is highly valued (Inamori et al. 2006; Nakashima and Shimizu 1972). Therefore, transcriptomic analysis of *T. dolabrata* var. *hondae* could strengthen the genetic knowledgebase to unveil the biosynthetic mechanism of podophyllotoxin and related compounds and improve tree breeding.

Five-year-old *T. dolabrata* var. *hondae* trees originating from the Shimokita Peninsula, Aomori Prefecture, Japan, were obtained from Aomori Tree Seedling Commission (Aomori, Japan). The plants were planted in pots with a 2:3 ratio mixture of vermiculite and peat moss-based soil for horticulture (Hanasaki-monogatari; Akimoto-tensanbutsu, Iga, Japan). We then isolated total RNA with sufficient quality and yield from young *T. dolabrata* var. *hondae* needles near the shoot tip by using a modified cetyltrimethylammonium bromide method (Suzuki et al. 2008).

Next, we analyzed *T. dolabrata* transcriptome by using two different next-generation sequencers, Genome Sequencer FLX (abbreviated hereafter as FLX; Roche Diagnostics, Basel, Switzerland) and Illumina Genome Analyzer IIX (abbreviated hereafter as IIX; Illumina, San Diego, USA). While IIX can read only about 100bp from both 3' and 5' ends (paired-ends) of adapter-added cDNAs in DNA synthesis using luminescent substrates, FLX can read 200–300bp from 3' ends of cDNA using a pyrosequencing method. We anticipated that we could obtain complementary transcriptomic information by using two different next-generation sequencing methods.

We outsourced FLX pyrosequencing of total RNA from young *T. dolabrata* var. *hondae* needles to Takara Bio Inc., Kusatsu, Japan. The sequences were provided as a multi-FASTA (.fna) file, and the sequence quality was recorded in a .qual file from Takara Bio. Sequences were assembled using PHRED and PHRAP programs as previously described (Suzuki et al. 2011). An aliquot of RNA from *T. dolabrata* var. *hondae* young needles was used to construct a cDNA library according to the RNA-Seq protocol (Illumina). The sample was hybridized to flow cell surfaces, cluster-amplified using a cBot Cluster Generation system (Illumina), and sequenced on IIX sequencer with 100-bp paired-end reads. The obtained reads were assembled using CLC Genomics Workbench version 5.5.1 (CLC Bio Japan, Tokyo, Japan) with a minimum contig length of 200bp and performing scaffolding. In Illumina RNA-seq analysis, Reads Per Kilobase of exon per Million mapped reads (RPKM) values (Mortazavi et al. 2008) were calculated using CLC Genomics Workbench with scaling normalization and a median value as a reference. Thus, whereas we sequenced 24,806 reads with an average 225bp totaling approximately 5.58Mbp using FLX sequencing, we paired and trimmed 6,412,133 reads with an average 95bp totaling 610Mbp using IIX. These reads have been deposited in the DNA Data Bank of Japan Sequence Read Archive (accession number DRA007617).

The reads obtained by FLX were assembled to non-redundant 3,730 contigs. The reads generated by IIX were assembled by CLC genomics workbench to give 29,925 non-redundant contigs. These obtained contigs, which will be uploaded on a server of Research Institute for Sustainable Humanosphere, Kyoto University (RISH Data Server; <http://database.rish.kyoto-u.ac.jp>), were individually submitted to BLASTX (Altschul et al. 1990) search against NCBI non-redundant protein sequence (nr) and Arabidopsis protein sequence (TAIR10; <https://www.arabidopsis.org>) databases (Suzuki et al. 2011) at the cutoff value of E-value $\leq 1e-5$ and E-value ≤ 1 , respectively. This process allowed us to determine the best-matching proteins in these public databases with the contigs at amino acid levels (Supplementary Tables S1 and S4). We then performed a keyword search against the TAIR10 annotation because many annotations tagged by BLASTX search against nr gave little information of gene functions. Using the keywords “O-methyltransferase”, “CYP”, and “2-oxoglutarate (2OG) and Fe(II)-dependent oxygenase” enabled us to select candidate sequences encoding OMTs, CYPs, and 2OGDs (Supplementary Tables S2, S3, S5–S7).

Assembling FLX contigs and a keyword search for “O-methyltransferase” afforded 9 non-redundant contigs that putatively encode OMTs (Supplementary Table S2). Among them, three proteins, TdhFLX522, TdhFLX1251, and TdhFLX2512, share relatively high amino acid

identities (66%, 63%, and 75%) with a leucine carboxyl methyltransferase (AT5G42760), a protein-L-isoaspartate methyltransferase 2 (AT5G50240), and caffeoyl-CoA OMT 1 (CCoAOMT1; AT4G34050) (Do et al. 2007) in *Arabidopsis*, respectively, according to TAIR10 annotation. In contrast to FLX transcriptomic analysis, IIX reads were assembled to more contigs: 28 different IIX contigs putatively encoding OMTs (Supplementary Table S5). Among these contigs, amino acid sequences derived from *TdhIIX01261* and *TdhIIX19500* showed high identities (72%) with a putative *Arabidopsis* caffeoyl-CoA O-methyltransferase 6 (CCoAOMT6; AT1G67980) (Raes et al. 2003), and those derived from *TdhIIX09399* and *TdhIIX17089* shared identities with an *Arabidopsis* isoprenylcysteine carboxyl O-methyltransferase (AT5G59500) at 30% and 50% identities, respectively, based on TAIR10 annotation.

Plant S-adenosyl L-methionine dependent OMTs are roughly classified into two groups. While the group PI-OMT I (Joshi and Chiang 1998; also named Type 2 OMT by Noel et al. 2003 and Lineage A by Lam et al. 2007) includes CCoAOMT and carboxyl OMTs, the group PI-OMT II (Joshi and Chiang 1998; also named Type 1 OMT by Noel et al. 2003 and Lineage B by Lam et al. 2007) contains *Arabidopsis* coniferaldehyde OMT 1 (CaldOMT1) (Nakatsubo et al. 2008) and is more diverse. Previously identified lignan OMTs are all classified as PI-OMT I (Lau and Sattely 2015; Ragamustari et al. 2013, 2014; Umezawa et al. 2013, 2019). We checked which previously identified lignan OMTs shared the highest E-value with *T. dolabrata* var. *hondae* FLX contigs by using BLASTX search, and

found that *TdFLX0850* shared the highest E-value with previously identified lignan OMTs (Table 1). In IIX contigs, either of *TdhIIX05783* or *TdhIIX03595* shared the highest E-value with the previously identified lignan OMTs (Table 1). TAIR10 annotation indicated that 2 FLX contigs and 18 IIX contigs shared more or less identities with *Arabidopsis* CaldOMT1 in amino acid levels (Supplementary Tables S2 and S5). Collectively, *T. dolabrata* var. *hondae* lignan OMT(s) may be encoded by some of these contigs.

Several CYPs are involved in lignan biosynthesis. For example, CYP81Q1 catalyzes the methylenedioxy bridge formation in (+)-sesamin biosynthesis (Ono et al. 2006); CYP92B14 performs oxidative rearrangement of (+)-sesamin to (+)-sesamol and (+)-sesaminol (Murata et al. 2017); CYP719A11 mediates the conversion of (–)-matairesinol to (–)-pulvitolide (Marques et al. 2013); CYP71CU1, CYP71BE54, and CYP82D61 hydroxylate the B-ring of (–)-burshehnerin, demethylate the methoxyl group at the 4'-position of (–)-deoxypodophyllotoxin, and hydroxylate the 7-position of (–)-4-demethyldeoxypodophyllotoxin in *P. hexandrum* etoposide aglycone formation, respectively (Figure 2). Therefore, we anticipated that several CYP genes were involved in biosynthesis of yatein and its derivatives in *T. dolabrata* var. *hondae*.

Both FLX and IIX analyses afforded many contigs, 57 and 298 respectively, annotated as putative CYPs (Supplementary Tables S3 and S6). We performed a TBLASTN search against these contigs using the known lignan CYPs as queries. Among FLX contigs, *TdhFLX3464* was top-ranked in terms of E-value in

Table 1. The relationship between the previously identified lignan OMTs and putative *T. dolabrata* var. *hondae* OMTs.

OMT name	FLX Contig			IIX Contig			
	Name	Identity	E-value	Name	Identity	E-value	RPKM
CtMROMT	<i>TdFLX0850</i>	54%	4e-29	<i>TdhIIX05783</i>	37%	4e-79	76.42
FkMROMT	<i>TdFLX0850</i>	53%	4e-32	<i>TdhIIX03595</i>	55%	e-127	53.15
AsMROMT	<i>TdFLX0850</i>	59%	3e-30	<i>TdhIIX05783</i>	39%	4e-82	76.42
AsTJOMT	<i>TdFLX0850</i>	52%	3e-31	<i>TdhIIX03595</i>	53%	e-113	53.15
PhOMT1	<i>TdFLX0850</i>	42%	1e-23	<i>TdhIIX05783</i>	36%	2e-68	76.42
PhOMT3	<i>TdFLX0850</i>	52%	3e-32	<i>TdhIIX03595</i>	54%	e-122	53.15

Putative *T. dolabrata* var. *hondae* OMTs were selected based on the highest E-value.

Table 2. The relationship between the previously identified lignan CYPs and putative *T. dolabrata* var. *hondae* CYPs.

CYP name	FLX Contig			IIX Contig			
	Name	Identity	E-value	Name	Identity	E-value	RPKM
CYP81Q1	<i>TdhFLX3464</i>	40%	2e-42	<i>TdhIIX00903</i>	35%	1e-96	74.34
CYP92B14	<i>TdhFLX3464</i>	38%	5e-44	<i>TdhIIX00903</i>	48%	e-173	74.34
CYP719A11	<i>TdhFLX3464</i>	33%	5e-27	<i>TdhIIX03868</i>	33%	1e-91	153.57
CYP71CU1	<i>TdhFLX3464</i>	40%	3e-45	<i>TdhIIX00903</i>	36%	e-118	74.34
CYP71BE54	<i>TdhFLX3464</i>	42%	1e-51	<i>TdhIIX06243</i>	41%	e-123	29.17
CYP82D61	<i>TdhFLX3464</i>	42%	7e-44	<i>TdhIIX00981</i>	40%	e-109	83.69

Putative *T. dolabrata* var. *hondae* CYPs were selected based on the highest E-value.

comparison with the previously identified lignan CYPs. *TdIIx00903*, *TdhIIx00981*, *TdhIIx03868*, and *TdhIIx06243*, were ranked as the highest E-value CYPs (Table 2). It is noteworthy that *TdhFLX3464*, *TdhIIx00903*, *TdhIIx00981*, *TdhIIx03868*, and *TdhIIx06243* shared identities with any of CYP71B35, CYP75B1, CYP82C2, CYP71B37, and CYP75B1 in comparison with all *Arabidopsis* CYPs (Supplementary Tables S3 and S6). However, further analysis is required to reduce the number of candidate contigs, as shown in a previous study of a *P. hexandrum* (–)-podophyllotoxin inducible experimental system (Lau and Sattely 2015).

Considerable numbers of 2-oxoglutarate dependent dioxygenases (2OGDs) are involved in plant secondary metabolism (Kawai et al. 2014). In biosynthesis of podophyllotoxin-related compounds in *P. hexandrum*, a 2OGD (Phx30848; deoxypodophyllotoxin synthase) is responsible for the formation of aryltetralin structure in the enzymatic conversion of (–)-yatein to (–)-deoxypodophyllotoxin (Lau and Sattely 2015). Therefore, we searched putative 2OGDs expressing in young needles of *T. dolabrata* var. *hondae*. As a result, compared with the number of CYPs, fewer contigs were identified using the IIX assemble collection (105 contigs; Supplementary Table S7). However, the highest identity of *P. hexandrum* deoxypodophyllotoxin synthase with *T. dolabrata* var. *hondae* 2OGD (*Tdh13966*) was as low as 30%, which suggests that further analyses are required to narrow down the candidates.

In summary, we established a collection of candidate gene sequences that encode OMT, CYP, and 2OGD family proteins in young *Thujopsis dolabrata* var. *hondae* needles, by de novo assembly of reads generated from two different next-generation sequencers. Our analysis is the genetic basis for biochemical identification of lignan biosynthetic genes in *T. dolabrata* var. *hondae*.

Acknowledgements

We acknowledge Tomoko Kawabata and Atsushi Hosouchi for technical assistance of cultivation of *T. dolabrata* var. *hondae* and Illumina RNA-seq sequencing, respectively. We also thank Tohoku Regional Breeding Office, Forest Tree Breeding Center, Forest Research and Management Organization, Iwate, Japan, for technical assistance. This research was partly supported by Grants-in-Aid for Scientific Research from JSPS (no. 18H02255) and a Special Fund for Area Studies for Sustainability from the Institute of Sustainability Science, Kyoto University, Japan. This work also performed with the assistance of Development and Assessment System for Humanosphere (DASH) of the Research Institute for Sustainable Humanosphere and Research Center of Ecosystems, Kyoto University, on preliminary GC-MS and LC-MS analysis of lignans in *T. dolabrata* var. *hondae*.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic

- local alignment search tool. *J Mol Biol* 215: 403–410
- Bhattacharyya D, Hazra S, Banerjee A, Datta R, Kumar D, Chakrabarti S, Chattopadhyay S (2016) Transcriptome-wide identification and characterization of CAD isoforms specific for podophyllotoxin biosynthesis from *Podophyllum hexandrum*. *Plant Mol Biol* 92: 1–23
- Broomhead AJ, Rahman MMA, Dewick PM, Jackson DE, Lucas JA (1991) Matairesinol as precursor of *Podophyllum* lignans. *Phytochemistry* 30: 1489–1492
- Dewick PM (1993) The biosynthesis of shikimate metabolism. *Nat Prod Rep* 10: 233–263
- Do CTP, Pollet B, Thévenin J, Sibout R, Denoue D, Barrière Y, Lapierre C, Jouanin L (2007) Both caffeoyl coenzyme A 3-O-methyltransferase 1 and caffeic acid O-methyltransferase 1 are involved in redundant functions for lignin flavonoids and sinapoyl malate biosynthesis in *Arabidopsis*. *Planta* 226: 1117–1129
- Hendrawati O, Hille J, Woerdenbag HJ, Quax WJ, Kayser O (2012) In vitro regeneration of wild chervil (*Anthriscus sylvestris* L.). *In Vitro Cell Dev Biol Plant* 48: 355–361
- Inamori Y, Morita Y, Sakagami Y, Okabe T, Ishida N (2006) The excellence of Aomori Hiba (Hinokiasunaro) in its use as building materials of Buddhist temples and Shinto Shrines. *Biocontrol Sci* 11: 49–54
- Jackson DE, Dewick PM (1984) Biosynthesis of *Podophyllum* lignans-II: Interconversions of aryltetralin lignans in *Podophyllum hexandrum*. *Phytochemistry* 23: 1037–1042
- Joshi CP, Chiang VL (1998) Conserved sequence motifs in plant Sadenosyl-l-methionine-dependent methyltransferases. *Plant Mol Biol* 37: 663–674
- Kamil WM, Dewick PM (1986a) Biosynthesis of the lignans α - and β -peltatin. *Phytochemistry* 25: 2089–2092
- Kamil WM, Dewick PM (1986b) Biosynthetic relationship of aryltetralin lactone lignans to dibenzylbutyrolactone lignans. *Phytochemistry* 25: 2093–2102
- Kanetoshi A, Fujimoto T, Hayashi T, Hori Y, Aoyama M, Saito N, Tsuda M, Mori M (1998) Cell proliferation suppressive components from *Thujopsis dolabrata* SIEB. et ZUCC. var. *hondae* MAKINO leaves. *Natural Medicines* 52: 444–447
- Kanetoshi A, Fujimoto T, Hori Y, Aoyama M, Saito N, Tsuda M, Cho NS, Kim HE (1999) Cell proliferation suppressive components in cupressaceae leaf extracts and determination of the contents. *Report of the Hokkaido Institute of Public Health* 49: 134–136 (in Japanese)
- Kawai S, Hasegawa T, Gotoh M, Ohashi H (1994) 4-O-Demethylyatein from the branch wood of *Thuja occidentalis*. *Phytochemistry* 37: 1699–1702
- Kawai S, Sugishita K, Ohashi H (1999) Identification of *Thuja occidentalis* lignans and its biosynthetic relationship. *Phytochemistry* 51: 243–247
- Kawai Y, Ono E, Mizutani M (2014) Evolution and diversity of the 2-oxoglutarate-dependent dioxygenase superfamily in plants. *Plant J* 78: 328–343
- Kranz K, Petersen L (2003) β -peltatin 6-O-methyltransferase from suspension cultures of *Liunum nodiflorum*. *Phytochemistry* 64: 453–458
- Kumar P, Jaiswal V, Pal T, Singh J, Chauhan RS (2017) Comparative whole-transcriptome analysis in *Podophyllum* species identifies key transcription factors contributing to biosynthesis of podophyllotoxin in *P. hexandrum*. *Protoplasma* 254: 217–228
- Kumari A, Singh HR, Jha A, Swarnkar MK, Shankar R, Kumar S (2014) Transcriptome sequencing of rhizome tissue of

- Sinopodophyllum hexandrum* at two temperatures. *BMC Genomics* 15: 871
- Lam KC, Ibrahim RK, Behdad B, Dayanandan S (2007) Structure, function, and evolution of plant O-methyltransferases. *Genome* 50: 1001–1013
- Lata H, Mizuno CS, Moraes R (2009) The role of biotechnology in the production of the anticancer compound podophyllotoxin. In: Jain SM, Saxena PK (eds) *Protocols for In Vitro Cultures and Methods in Molecular Biology, Methods in Molecular Biology* 547. Humana Press, New York, pp 387–402
- Lau W, Sattely ES (2015) Six enzymes from mayapple that complete the biosynthetic pathway to the etoposide aglycone. *Science* 349: 1224–1228
- Lim YH, Leem MJ, Shin DH, Chang HB, Hong SW, Moon EY, Lee DK, Yoon SJ, Woo WS (1999) Cytotoxic constituents from the roots of *Anthriscus sylvestris*. *Arch Pharm Res* 22: 208–212
- Marques JV, Kim K-W, Lee C, Costa MA, May GD, Crow JA, Davin LB, Lewis NG (2013) Next generation sequencing in predicting gene function in podophyllotoxin biosynthesis. *J Biol Chem* 288: 466–479
- Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B (2008) Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat Methods* 5: 621–628
- Murata J, Ono E, Yoroizuka S, Toyonaga H, Shiraishi A, Mori S, Tera M, Azuma T, Nagano AJ, Nakayasu M, et al. (2017) Oxidative rearrangement of (+)-sesamin by CYP92B14 co-generates twin dietary lignans in sesame. *Nat Commun* 8: 2155
- Nakashima Y, Shimizu K (1972) Studies on an antitermitic activity of Hinokiasunaro (*Thujopsis dolabrata* Sieb. et Zucc. var. *hondai* Makino). III. The components with a termiticidal activity. *Miyazaki Daigaku Nogakubu Kenkyu Hokoku* 19: 251–259
- Nakatsubo T, Kitamura Y, Sakakibara N, Mizutani M, Hattori T, Sakurai N, Shibata D, Suzuki S, Umezawa T (2008) At5g54160 gene encodes *Arabidopsis thaliana* 5-hydroxyconiferaldehyde O-methyltransferase. *J Wood Sci* 54: 312–317
- Noel JP, Dixon RA, Pichersky E, Zubietta C, Ferrer JL (2003) Structural, functional, and evolutionary basis for methylation of plant small molecules. In: Romeo JT (ed) *Recent Advances in Phytochemistry Vol. 37*. Elsevier, Amsterdam, pp 37–58
- Ono E, Nakai M, Fukui Y, Tomimori N, Fukuchi-Mizutani M, Saito M, Satake H, Tanaka T, Katsuta M, Umezawa T, et al. (2006) Formation of two methylenedioxy bridges by a *Sesamum* CYP81Q protein yielding a furofuran lignan, (+)-sesamin. *Proc Natl Acad Sci USA* 103: 10116–10121
- Raes J, Rohde A, Christensen JH, de Peer YV, Boerjan W (2003) Genome-wide characterization of the lignification toolbox in *Arabidopsis*. *Plant Physiol* 133: 1051–1071
- Ragamustari SK, Yamamura M, Ono E, Hattori T, Suzuki S, Suzuki H, Shibata D, Umezawa T (2014) Substrate-enantiomer selectivity of matairesinol O-methyltransferases. *Plant Biotechnol* 31: 257–267
- Ragamustari SK, Nakatsubo T, Hattori T, Ono E, Kitamura Y, Suzuki S, Yamamura M, Umezawa T (2013) A novel O-methyltransferase involved in the first methylation step of yatein biosynthesis from matairesinol in *Anthriscus sylvestris*. *Plant Biotechnol* 30: 375–384
- Sakakibara N, Suzuki S, Umezawa T, Shimada M (2003) Biosynthesis of yatein in *Anthriscus sylvestris*. *Org Biomol Chem* 1: 2474–2485
- Sasheva S, Ikonkova I (2017) Small cells for big ideas: The cytotoxic podophyllotoxin and the long journey in discovering its biosynthetic pathway. In: Malik S (ed) *Biotechnology and Production of Anti Cancer Compounds*. Springer, Cham, pp 177–199
- Seegers CL, Setroikromo R, Quax WJ (2017) Towards metabolic engineering of podophyllotoxin production. In: Badria FA (ed) *Natural Products and Cancer Drug Discovery*. Intech Open, London, pp 287–306
- Suzuki S, Umezawa T (2007) Biosynthesis of lignans and norlignans. *J Wood Sci* 53: 273–284
- Suzuki S, Sakakibara N, Umezawa T, Shimada M (2002) Survey and enzymatic formation of lignans of *Anthriscus sylvestris*. *J Wood Sci* 48: 536–541
- Suzuki S, Suda K, Sakurai N, Ogata Y, Hattori T, Suzuki H, Shibata D, Umezawa T (2011) Analysis of expressed sequence tags in developing secondary xylem and shoot of *Acacia mangium*. *J Wood Sci* 57: 40–46
- Suzuki Y, Mae T, Makino A (2008) RNA extraction from various recalcitrant plant tissues with a cethyltrimethylammonium bromide-containing buffer followed by an acid guanidium thiocyanate-phenol-chloroform treatment. *Biosci Biotechnol Biochem* 72: 1951–1953
- Umezawa T (2003a) Diversity in lignan biosynthesis. *Phytochem Rev* 2: 371–390
- Umezawa T (2003b) Phylogenetic distribution of lignan producing plants. *Wood Res* 90: 27–110
- Umezawa T, Ragamustari SF, Nakatsubo T, Wada S, Li L, Yamamura M, Sakakibara N, Hattori T, Suzuki S, Chiang V (2013) A lignan O-methyltransferase catalyzing the regioselective methylation of matairesinol in *Carthamus tinctorius*. *Plant Biotechnol* 30: 97–109
- Umezawa T, Yamamura M, Ono E, Shiraishi A, Ragamustari SK (2019) Recent advances in lignan OMT studies. *Mokuzai Gakkaishi* 65: 1–12 (in Japanese)
- Xia ZQ, Costa MA, Proctor J, Davin LB, Lewis NG (2000) Dirigent-mediated podophyllotoxin biosynthesis in *Linum flavum* and *Podophyllum peltatum*. *Phytochemistry* 55: 537–549