20. Characteristics of Air Pollution Concentration in Jakarta and Its Relationship with Local Flow Field

ジャワ島西部の局地風がジャカルタの大気汚染に与え得る影響

アセップ ソフィアン*。北田敏廣*。倉田 学児 *
Asep SOFYAN*, Toshihiro KITADA*, Gakuji KURATA*

ABSTRACT: The obtained spatial distributions of NO_2 and SO_2 show that the highest concentration appears near the high traffic main highway, and then higher concentrations in business center and industrial areas. The highest “one-day” NO_2 , appeared at a site near toll highway in central Jakarta, was $446.1 \mu\text{g}/\text{m}^3$; 16 observation points including this site exceeded $92.5 \mu\text{g}/\text{m}^3$, the Jakarta ambient air quality standard (Jakarta AAQS). On the other hand, “one day” SO_2 concentrations were below the Jakarta AAQS $260 \mu\text{g}/\text{m}^3$, though these at two sites were over the WHO AAQS $125 \mu\text{g}/\text{m}^3$. Vertical profile, up to 110 m high, of air pollution was also studied using a high building in business center of Jakarta. The “one day” NO_2 concentration were 119.8, 101.3, 87.6, and $100.4 \mu\text{g}/\text{m}^3$ at 10, 30, 55, and 110 m high, respectively, and similarly those for SO_2 were 7.3, 1.8, 6.5, and $16.2 \mu\text{g}/\text{m}^3$. It is interesting that the SO_2 increases with height, suggesting huge point sources of power plants located along the coast in the north-eastern part of Jakarta may be resulted in this profile under the sea breeze condition. At the symposium, the results will be discussed together with CTM calculation.

KEYWORDS: Jakarta, air pollution, local flow, sea breeze

1. Introduction

Jakarta, Indonesia has experienced serious air pollution problems associated with the use of energy in the transport, domestic, and industrial sectors. Transportation sector is responsible for approximately 70% of the total emissions in Jakarta. Large scale stationary sources such as power plant and metal industries also degrade air quality in Jakarta.

For planning of long term better air quality, to know quantitative relationship between air quality and emission sources' distribution in the area is the first but the most important step, since air flow and other meteorological factors which are characteristic in the area largely affect the relationship in complex manner. As a “first” step to this purpose, we conducted field observation for air quality over Jakarta city using a number of passive samplers for NO_2 and SO_2 in August 2004. The samplers were distributed in 50 locations to measure both 1 day and 1 week averaged concentration distribution. Using the observed values, contours of NO_2 and SO_2 were plotted together with the model simulated surface wind fields.

To understand the characteristics and the development mechanism of complex local flows, numerical simulation over western Java area was performed using the Fifth-Generation Pennsylvania State University-National Center for Atmospheric Research Mesoscale Model Version 3.6 (hereafter will be abbreviated as MM5; see Dudhia et al., 2003 for the detail of the software). The simulations were done for 8-13 February 2001 in “rainy” season and for 6-19 August 2004 in “dry” season. Latitude and longitude at the southwest and northeast corners of the outermost domain were 8°S and 105°E , and 5°S and 109°E , respectively. The domain system used in this calculation is a triply nested two-way interacting mesh. The domain 1 has 50×37 horizontal grids points, the domain 2 has 73×73 grids point, and the domain 3 has 82×82 grids points. The horizontal grid size was constant at 9, 3, and 1 km for the domain 1, 2, and 3, respectively (see Fig 1). Each domain has 23 vertical grid points for the depth from the earth's surface to 100 hPa. The framework of meteorology in larger scale was provided by ECMWF data with a resolution of $0.5^\circ \times 0.5^\circ$.

---

*Department of Ecological Engineering, Toyohashi University of Technology, Toyohashi, Japan

Fig.1 The domain 1, 2, and 3 have 9 km, 3 km, and 1 km grid size, respectively.

2. NO₂ and SO₂ distribution over Jakarta

For planning of long term better air quality, to know quantitative relationship between air quality and emission sources' distributions in the area is the first but the most important step, since air flow and other meteorological factors which are characteristic in the area largely affect the relationship in complex manner. As a "first" step to this purpose, we conducted field observation for air quality over Jakarta city using a number of passive samplers for NO₂ and SO₂ in August 2004. The samplers were distributed in 50 locations to measure both 1 day and 1 week averaged concentration distributions. Using the observed values, contours of NO₂ and SO₂ were plotted together with the model simulated surface wind fields (see Figs. 2 and 3). The obtained spatial distributions of NO₂ and SO₂ show that the highest concentration appears near the high traffic main highway, and then higher concentrations in business center and industrial areas. The highest "one-day" NO₂ was 446.1 $\mu\text{g}/\text{m}^3$, appeared at a site near toll highway in central Jakarta; 16 observation points including this site exceeded 92.5 $\mu\text{g}/\text{m}^3$, which is the Jakarta ambient air quality standard (Jakarta AAQS). On the other hand, "one-day" SO₂ concentrations were below the Jakarta AAQS 260 $\mu\text{g}/\text{m}^3$, though these at two sites exceeded the WHO AAQS 125 $\mu\text{g}/\text{m}^3$. Vertical profile, up to 110 m high, of air pollution was also studied using a high building in business center of Jakarta. The "one-day" NO₂ concentrations were 119.8, 101.3, 87.6, and 100.4 $\mu\text{g}/\text{m}^3$ at 10, 30, 55, and 110 m high, respectively, and similarly those for SO₂ were 7.3, 1.8, 6.5, and 16.2 $\mu\text{g}/\text{m}^3$. It is interesting that the SO₂ increases with height, suggesting huge point sources of power plants located along the coast in the north-eastern part of Jakarta may have resulted in this profile under the sea breeze condition (see Fig. 4).

3. Local Flow Field over Jakarta area

Characteristics of local flow in Jakarta area were numerically investigated for wet (rainy) and dry seasons. In the "rainy" season of 6-13 Feb, 2001, strong synoptic scale WSW wind regularly existed, and it suppressed penetration of NW sea breeze from the Java Sea on sunny days by forming a convergence with the sea breeze. Subsidence behind the convergence line (the sea breeze front) generated stable thin-layer above the sea breeze and thus part of the Jakarta area, indicating possible trap of the air pollutants below the layer and increase of air pollution potential. In the "dry" season of 6-19 Aug, 2004, though synoptic scale wind in upper layer was constantly southeasterly, it was very weak in the plain area of Jakarta because the synoptic SE wind was blocked by the mountains along the south coast of the Java Island. Hence, local winds of sea breeze from the Java Sea and valley wind over the northern slope of the southern mountains fully developed, forming one large scale combined local flow from the Java Sea to the mountains. This situation might give better ventilation of polluted air mass over Jakarta, though the "better ventilation" means export of air pollutants into the rural area.

Fig. 2 Observed spatial distribution of NO₂; measured with 50 passive samplers distributed over the greater Jakarta during short and long term measurements in August 2004.

[illegible]

1. Toll Road (11)   2. Near Toll Road (4)   3. Near Powerplant (3)   4. Industrial Area (7)
5. High Building (1)   6. Urban Area (15)   7. Sub Urban (3)   8. Rural (5)

The map displays the following PM2.5 concentrations across various locations in Indonesia:

- Sumatra:** 28.9 ppb (North Sumatra), 8.5 ppb (Aceh), 2.1 ppb (Sumatra Utara), 0.1 ppb (Sumatra Barat), 2.2 ppb (Sumatra Barat), 0.4 ppb (Sumatra Barat), 1.1 ppb (Sumatra Barat), 1.0 ppb (Sumatra Barat).
- Java:** 2.1 ppb (Jawa Barat), 2.5 ppb (Jawa Barat), 2.1 ppb (Jawa Barat), 4.4 ppb (Jawa Barat), 4.5 ppb (Jawa Barat), 3.5 ppb (Jawa Barat), 1.4 ppb (Jawa Barat), 2.5 ppb (Jawa Barat), 4.0 ppb (Jawa Barat), 1.6 ppb (Jawa Barat), 3.5 ppb (Jawa Barat), 3.1 ppb (Jawa Barat), 3.9 ppb (Jawa Barat), 4.9 ppb (Jawa Barat), 3.8 ppb (Jawa Barat), 1.2 ppb (Jawa Barat), 3.8 ppb (Jawa Barat), 1.3 ppb (Jawa Barat), 1.5 ppb (Jawa Barat), 1.1 ppb (Jawa Barat), 0.4 ppb (Jawa Barat), 1.1 ppb (Jawa Barat), 1.0 ppb (Jawa Barat).
- Central Java:** 2.1 ppb (Central Java), 2.5 ppb (Central Java), 2.1 ppb (Central Java), 4.4 ppb (Central Java), 4.5 ppb (Central Java), 3.5 ppb (Central Java), 1.4 ppb (Central Java), 2.5 ppb (Central Java), 4.0 ppb (Central Java), 1.6 ppb (Central Java), 3.5 ppb (Central Java), 3.1 ppb (Central Java), 3.9 ppb (Central Java), 4.9 ppb (Central Java), 3.8 ppb (Central Java), 1.2 ppb (Central Java), 3.8 ppb (Central Java), 1.3 ppb (Central Java), 1.5 ppb (Central Java), 1.1 ppb (Central Java), 0.4 ppb (Central Java), 1.1 ppb (Central Java), 1.0 ppb (Central Java).
- East Java:** 2.1 ppb (East Java), 2.5 ppb (East Java), 2.1 ppb (East Java), 4.4 ppb (East Java), 4.5 ppb (East Java), 3.5 ppb (East Java), 1.4 ppb (East Java), 2.5 ppb (East Java), 4.0 ppb (East Java), 1.6 ppb (East Java), 3.5 ppb (East Java), 3.1 ppb (East Java), 3.9 ppb (East Java), 4.9 ppb (East Java), 3.8 ppb (East Java), 1.2 ppb (East Java), 3.8 ppb (East Java), 1.3 ppb (East Java), 1.5 ppb (East Java), 1.1 ppb (East Java), 0.4 ppb (East Java), 1.1 ppb (East Java), 1.0 ppb (East Java).
- South Java:** 2.1 ppb (South Java), 2.5 ppb (South Java), 2.1 ppb (South Java), 4.4 ppb (South Java), 4.5 ppb (South Java), 3.5 ppb (South Java), 1.4 ppb (South Java), 2.5 ppb (South Java), 4.0 ppb (South Java), 1.6 ppb (South Java), 3.5 ppb (South Java), 3.1 ppb (South Java), 3.9 ppb (South Java), 4.9 ppb (South Java), 3.8 ppb (South Java), 1.2 ppb (South Java), 3.8 ppb (South Java), 1.3 ppb (South Java), 1.5 ppb (South Java), 1.1 ppb (South Java), 0.4 ppb (South Java), 1.1 ppb (South Java), 1.0 ppb (South Java).
- West Java:** 2.1 ppb (West Java), 2.5 ppb (West Java), 2.1 ppb (West Java), 4.4 ppb (West Java), 4.5 ppb (West Java), 3.5 ppb (West Java), 1.4 ppb (West Java), 2.5 ppb (West Java), 4.0 ppb (West Java), 1.6 ppb (West Java), 3.5 ppb (West Java), 3.1 ppb (West Java), 3.9 ppb (West Java), 4.9 ppb (West Java), 3.8 ppb (West Java), 1.2 ppb (West Java), 3.8 ppb (West Java), 1.3 ppb (West Java), 1.5 ppb (West Java), 1.1 ppb (West Java), 0.4 ppb (West Java), 1.1 ppb (West Java), 1.0 ppb (West Java).
- South Sumatra:** 2.1 ppb (South Sumatra), 2.5 ppb (South Sumatra), 2.1 ppb (South Sumatra), 4.4 ppb (South Sumatra), 4.5 ppb (South Sumatra), 3.5 ppb (South Sumatra), 1.4 ppb (South Sumatra), 2.5 ppb (South Sumatra), 4.0 ppb (South Sumatra), 1.6 ppb (South Sumatra), 3.5 ppb (South Sumatra), 3.1 ppb (South Sumatra), 3.9 ppb (South Sumatra), 4.9 ppb (South Sumatra), 3.8 ppb (South Sumatra), 1.2 ppb (South Sumatra), 3.8 ppb (South Sumatra), 1.3 ppb (South Sumatra), 1.5 ppb (South Sumatra), 1.1 ppb (South Sumatra), 0.4 ppb (South Sumatra), 1.1 ppb (South Sumatra), 1.0 ppb (South Sumatra).
- Other Regions:** 0.8 ppb (Kalimantan), 1.5 ppb (Kalimantan), 41.8 ppb (Kalimantan), 13.6 ppb (Kalimantan), 16.7 ppb (Kalimantan), 11.2 ppb (Kalimantan), 11.3 ppb (Kalimantan), 2.2 ppb (Kalimantan), 0.1 ppb (Kalimantan).

1. Toll Road (11)	2. Near Toll Road (4)	3. Near Powerplant (3)	4. Industrial Area (7)
5. High Building (1)	6. Urban Area (15)	7. Sub Urban (3)	8. Rural (6)

— 126 —

Fig. 4. Vertical profiles of measured NO₂ and SO₂ at business center of Jakarta: “black” for short term (1-day) and “red” for long term (1-week) term measurements.

3. Conclusion

Field observation for air quality over Jakarta city using a number of passive samplers for NO₂ and SO₂ in August 2004. The samplers were distributed in 50 locations to measure both 1 day and 1 week averaged concentration distributions. The obtained spatial distributions of NO₂ and SO₂ show that the highest concentration appears near the high traffic main highway, and then higher concentrations in business center and industrial areas.

Characteristics of local flow in Jakarta area were numerically investigated for wet (rainy) and dry seasons. In the “rainy” season of 6-13 Feb, 2001, strong synoptic scale WSW wind regularly existed, and it suppressed penetration of NW sea breeze from the Java Sea on sunny days by forming a convergence with the sea breeze. In the “dry” season of 6-19 Aug, 2004, though synoptic scale wind in upper layer was constantly southeasterly, it was very weak in the plain area of Jakarta because the synoptic SE wind was blocked by the mountains along the south coast of the Java Island. Hence, local winds of sea breeze from the Java Sea and valley wind over the northern slope of the southern mountains fully developed, forming one large scale combined local flow from the Java Sea to the mountains. This situation might give better ventilation of polluted air mass over Jakarta, though the “better ventilation” means export of air pollutants into the rural area.

References

References

1. Dudhia J and Gill D (2003), PSU/NCAR Mesoscale Modeling System Tutorial Class Notes and User's Guide: MM5 Modeling System Version 3, Mesoscale and Microscale Meteorology Division, National Center for Atmospheric Research.
2. Kitada T., Okamura K, and Tanaka S (1998) Effect of Topography and Urbanization on Local Winds and Thermal Environment in the Nohbi Plain, Coastal Region of Central Japan: A Numerical Analysis by Mesoscale Meteorological Model, *Journal of Applied Meteorology*, **37**, 1026-1046.
3. McGregor, G. R. and Nieuwolt, S. (1998), *Tropical Climatology*, Second Edition, John Wiley & Sons Ltd, 339pp.
4. Regmi, R.P., Kitada, T., and Kurata G.(2003) Numerical Simulation of Late Wintertime Local Flows in Kathmandu Valley, Nepal: Implication for Air Pollution Transport, *Journal of Applied Meteorology*, **42**, 389-402.
5. Riehl, H. (1979) *Climate and Weather in the Tropics*, Academic Press, 611p.