NOTE
 =====

(39)

book⁵⁾, which has been cited in the appendix of the previous paper⁶⁾. Sr was co-precipitated by calcium oxalate on the basis of the method recommended in the text book⁴⁾, by which the yield of Sr has been examined in the previous paper²⁾.

To examine cross contamination of radioactive Sr and Cs, special chemical procedures were designed. For Sr fraction, as illustrated in Fig. 1, after removal of Cs by co-precipitation with ammonium phosphomolybdate, Cs, possibly remaining in solution, was co-precipitated according to the second step of the procedure. For Cs fraction, as illustrated in Fig. 2, after removal of radioactive Sr by co-precipitation with calcium oxalate, Sr, possibly remaining in solution, was co-precipitated according to the second step of the procedure. The final precipitates obtained from both the procedures were subjected to the radioactivity measurement, for the examination of cross contamination. The experiment was triplicated.

Prior to this study, an examination was made for the effect of ammonium chloride concentration on the co-precipitation yield of Sr as described in Appendix 1.

Fig. 1 Chemical procedure for examination of contamination of Sr fraction with ¹³⁷Cs.

Fig. 2 Chemical procedure for examination of contamination of Cs fraction with ⁸⁹Sr or ⁹⁰Sr + ⁹⁰Y.

2. Results

2.1 Contamination of Sr fraction with ¹³⁷Cs

The results of this experiment are shown in Table 1. Without phosphomolybdate treatment, that is in the case of the direct measurement of Sr fraction, the degree of contamination* was in the order of 10⁻⁴. The degree decreased down to 10⁻⁶ by the first treatment of phosphomolybdate. The repetition of phosphomolybdate treatment reduced slightly the degree, whereas the addition of Cs as a hold back carrier had no effect.

2.2 Contamination of Cs fraction with ⁹⁰Sr

As shown in Table 2, the direct measurement of Cs fraction gave a high degree of

* "Degree of contamination" is the reciprocal of "decontamination factor".

$$\text{Degree of contamination} = \frac{\text{Final activity (cpm)}}{\text{Initial activity (cpm)}}$$

contamination with ^{90}Sr (the order of 10^{-1}).

The contaminants were identified to be composed of ^{90}Y mainly and ^{90}Sr in part, from decay curves. Most of contaminants was removed by the first treatment of oxalate precipitate, down to 10^{-3} in terms of the

degree of contamination. The degree decreased down to 10^{-4} by the repetition of oxalate treatment. The use of Sr as a collector for the oxalate precipitation instead of Ca did not give any profit.

Table 1 Contamination of Sr fraction with ^{137}Cs

No.	Ca added (mg)	Cs carrier added (mg)	Phosphomolybdate treatment	cpm*	Degree of contamination
1	15	—	—	170 $\pm$ 63	$(2.0 \pm 0.7) \times 10^{-4}$
2	"	1	—	318 $\pm$ 50	$(3.8 \pm 0.6) \times 10^{-4}$
3	"	—	0.1 g, single	4.6 $\pm$ 1.5	$(5.4 \pm 1.8) \times 10^{-6}$
4	"	1	"	5.5 $\pm$ 0.7	$(6.5 \pm 0.8) \times 10^{-6}$
5	"	—	0.1 g, repeated	1.8 $\pm$ 0.3	$(2.1 \pm 0.3) \times 10^{-6}$
6	"	1	"	1.6 $\pm$ 0.3	$(1.9 \pm 0.4) \times 10^{-6}$

* By the use of an anti-coincidence LBC counter, background=1.2 cpm

Table 2 Contamination of Cs fraction with ^{90}Sr

No.	Ca or Sr added (mg)	Oxalate treatment	cpm*	Degree of contamination
11	—	—	$(537 \pm 100) \times 10^2$	$(1.4 \pm 0.3) \times 10^{-1}$
12	15—Ca	—	$(463 \pm 112) \times 10^2$	$(1.2 \pm 0.3) \times 10^{-1}$
13	30—Sr	—	$(445 \pm 39) \times 10^2$	$(1.1 \pm 0.2) \times 10^{-1}$
14	15—Ca	single	1,067 $\pm$ 87	$(2.7 \pm 0.3) \times 10^{-3}$
15	30—Sr	"	2,040 $\pm$ 30	$(5.2 \pm 0.2) \times 10^{-3}$
16	15—Ca	repeated	37 $\pm$ 12	$(0.9 \pm 0.3) \times 10^{-4}$
17	30—Sr	"	99 $\pm$ 26	$(2.5 \pm 0.7) \times 10^{-4}$

* By the use of an end-window GMLB counter, background=7 cpm

Table 3 Contamination of Cs fraction with ^{89}Sr

No.	Ca or Sr added (mg)	Oxalate treatment	cpm*	Degree of contamination
21	—	—	5,489 $\pm$ 880	$(7.5 \pm 1.2) \times 10^{-3}$
22	15—Ca	—	3,852 $\pm$ 616	$(5.3 \pm 0.8) \times 10^{-3}$
23	30—Sr	—	3,641 $\pm$ 604	$(5.0 \pm 0.8) \times 10^{-3}$
24	15—Ca	single	355 $\pm$ 98	$(4.9 \pm 1.3) \times 10^{-4}$
25	30—Sr	repeated	165 $\pm$ 120	$(2.3 \pm 1.6) \times 10^{-4}$

* By the use of an end-window GM counter, background=37 cpm

2-3 Contamination of Cs fraction with ^{90}Sr

To avoid the contribution of ^{90}Y , which was identified as the main contaminant for Cs fraction, ^{90}Sr was used instead of ^{90}Sr . As shown in Table 3, the direct measurement of Cs fraction gave the degree of contamination less than 10^{-2} . This value decreased down to 10^{-4} by the oxalate treatment. The use of Sr instead of Ca for precipitation of radioactive fraction had no effect.

3. Conclusion

The degree of contamination of Sr fraction with ^{137}Cs was 10^{-4} for the direct measurement of ^{90}Sr , and 10^{-6} after the removal of Cs fraction by phosphomolybdate treatment. The addition of Cs as a hold back carrier had no effect in this experiment. Moreover a large amount of Cs carrier inhibits very much the co-precipitation yield of Cs (see Appendix 2).

The degree of contamination of Cs fraction with radioactive Sr was 10^{-1} when ^{90}Sr coexisted; this contamination was mainly caused by daughter ^{90}Y . The degree decreased down to 10^{-3} by single application of oxalate treatment, and down to 10^{-4} by repetition of the treatment. ^{90}Sr having no radioactive daughter gave the value less than 10^{-2} without oxalate treatment; it gave 10^{-4} after first oxalate treatment, that is the same value as that for ^{90}Sr after second treatment. The use of Sr instead of Ca for precipitation of radioactive fraction had no effect.

Acknowledgment

The authors wish to thank Dr. H. Kobayashi for many helpful suggestions during this work.

References

- 1) Kodaira, K. and Kitayama, K.: A method of emission-spectrographic determination of strontium for soil analysis, *Bunko Kenkyu (J. Spectroscopical Soc. Japan)*, (in Japanese), **13**, 18-23 (1964)

- 2) Kodaira, K. and Shinohara, R.: Simultaneous radiochemical determination of ^{85}Sr and ^{45}Ca by oxalate precipitation method, *Radioisotopes*, **14**, 78-82 (1965)
- 3) Kodaira, K., Tsumura, A. and Kobayashi, H.: Studies on the mechanism of radiocontamination in rice plants (Part 1), submitted to *J. Radiation Research*
- 4) Science and Technology Agency, Japan: Determination of radioactive strontium, Edited by the Committee on radioactivity measurement, Radiation council, (in Japanese), (1963)
- 5) Science and Technology Agency, Japan: Determination of cesium-137, Edited by the Committee on radioactivity measurement, Radiation council, (in Japanese), (1963)
- 6) Yamagata, N., Kodaira, K. and Matsuda, S.: Cesium-137 in Japanese soil and rice, *J. Radiation Research*, **6**, 32-44 (1965)

Appendix 1.

Effect of ammonium chloride on the yield of Sr co-precipitated by oxalate method

In the above-mentioned study, a relatively large amount of hydrochloric acid was added to complete the co-precipitation of Cs together with ammonium phosphomolybdate. This gives rise to a high concentration of ammonium chloride, when the solution is neutralized for preparation of oxalate precipitate. The effect of ammonium chloride concentration on the yield of Sr was examined by the use of ^{90}Sr as shown in the table. Although a slight decrease in the yield of Sr was observed at the concentration of 3.6 N ammonium chloride, it did not disturb the outline of examination of cross contamination.

NH_4Cl (N)	0.36	0.6	3.6
Yield of Sr (%)	100	100*	94

* Taken as 100

Appendix 2.**Effect of various ions on the yield of Cs co-precipitated together with ammonium phosphomolybdate**

The effect of monovalent and divalent cations on the yield of Cs was examined, under such a condition as 50 ml in total volume, 0.6 *N* in hydrochloric acid concen-

tration and 0.1 g of ammonium phosphomolybdate. The yield of ^{137}Cs was expressed in terms of percentage in the table below.

Monovalent cations gave low values of co-precipitation yield of ^{137}Cs , whereas divalent cations did not.

In another experiment without any interfering ions, $99.0 \pm 0.15\%$ of ^{137}Cs was recovered by this co-precipitation method.

Ion species		NH_4^+	Na^+	K^+	Rb^+	Cs^+	$\text{Mg}^{\#}$	$\text{Ca}^{\#}$	$\text{Sr}^{\#}$	$\text{Ba}^{\#}$
Amounts added	0.5 m.e.	79	94	95	67	14	95	—	95	104
	1.0 m.e.	77	83	88	54	8	97	101	99	101